

SUPPLEMENTARY INFORMATION

Novel star-shaped viologens containing phenyl and triphenylamine moieties for electrochromic applications

Radosław Banasz¹ and Monika Wałęsa-Chorab^{1,*}

¹*Faculty of Chemistry, Adam Mickiewicz University in Poznań, Uniwersytetu Poznańskiego 8,
61-614 Poznań, Poland*

Table of content

Table S1. Solubility of 1 and 2 in different organic solvents	2
Figure S1. ^1H NMR spectra of compound (TPA) Br_3 in CDCl_3	3
Figure S2. ^{13}C NMR spectra of compound (TPA) Br_3 in CDCl_3	3
Figure S3. ^1H NMR spectra of compound A in CDCl_3	4
Figure S4. ^{13}C NMR spectra of compound A in CDCl_3	4
Figure S5. ^1H NMR spectra of compound B in CDCl_3	5
Figure S6. ^{13}C NMR spectra of compound B in CDCl_3	5
Figure S7. ^1H NMR spectra of compound 1 in d_6 -DMSO.....	6
Figure S8. ^{13}C NMR spectra of compound 1 in d_6 -DMSO.....	6
Figure S9. ^1H NMR spectra of compound 2 in d_6 -DMSO.....	7
Figure S10. ^{13}C NMR spectra of compound 2 in d_6 -DMSO.....	7
Figure S11. HR-ESI-MS spectra of compound (TPA) Br_3	8
Figure S12. HR-ESI-MS spectra of compound A	8
Figure S13. HR-ESI-MS spectra of compound B	9
Figure S14. HR-ESI-MS spectra of compound 1	9
Figure S15. HR-ESI-MS spectra of compound 2	10
Figure S16. Normalized spectra of 2 in solvents of different polarities: ethyl acetate (\blacktriangle green), tetrahydrofuran (\blacktriangledown red), dichloromethane (\blacklozenge yellow), methanol (\bullet orange), acetonitrile (\blacksquare black).....	10

Table S1. Solubility of **1** and **2** in different organic solvents

Solvent	1 PF_6^-	1 I^-	2 PF_6^-	2 I^-	
Ethyl acetate	+	-	+	-	
Tetrahydrofuran	+	-	+	-	
Dichloromethane	+	-	+	-	
Methanol	+	+	+	+	
Acetonitrile	+	+	+	+	




Figure S1. ¹H NMR spectra of compound (TPA)Br₃ in CDCl₃




Figure S2. ¹³C NMR spectra of compound (TPA)Br₃ in CDCl₃




Figure S3. ^1H NMR spectra of compound A in CDCl_3




Figure S4. ^{13}C NMR spectra of compound A in CDCl_3




Figure S5. ^1H NMR spectra of compound **B** in CDCl_3




Figure S6. ^{13}C NMR spectra of compound **B** in CDCl_3




Figure S7. ^1H NMR spectra of compound **1** in $\text{d}_6\text{-DMSO}$




Figure S8. ^{13}C NMR spectra of compound **1** in $\text{d}_6\text{-DMSO}$




Figure S9. ^1H NMR spectra of compound **2** in $\text{d}_6\text{-DMSO}$




Figure S10. ^{13}C NMR spectra of compound **2** in $\text{d}_6\text{-DMSO}$




Figure S11. HR-ESI-MS spectra of compound $(\text{TPA})\text{Br}_3$




Figure S12. HR-ESI-MS spectra of compound A




Figure S13. HR-ESI-MS spectra of compound **B**




Figure S14. HR-ESI-MS spectra of compound **1**




Figure S15. HR-ESI-MS spectra of compound 2




Figure S16. Normalized spectra of 2 in solvents of different polarities: ethyl acetate (\blacktriangle green), tetrahydrofuran (\blacktriangledown red), dichloromethane (\blacklozenge yellow), methanol (\bullet orange), acetonitrile (\blacksquare black).