

Table S1. The ophthalmic characteristics of patients positive for SARS-CoV-2 infection and diagnosed with herpes simplex virus-1 keratitis.

	Factor	Total cases
The patients positive for SARS-CoV-2 by RT-PCR with herpes simplex-1 keratitis		18
BCVA of the affected eye		
Before treatment	20/20 - 20/25 20/32 - 20/50 20/63 - 20/100 20/125 - 20/200 Counting fingers Hand motion Light perception No light perception	4 6 3 3 1 0 1 0
After treatment	20/20 - 20/25 20/32 - 20/50 20/63 - 20/100 20/125 - 20/200 Counting fingers Hand motion Light perception No light perception	12 3 1 1 0 0 1 0
Length of treatment		
	< 7 days 8-14 days 15-21 days 22- 28 days > 28 days	0 2 5 8 3
COVID-19 RT-PCR test	Prior to HSV-1 keratitis diagnosis	> 14 days 14 - 8 days 7 - 3 days 2- 0 days
		2 3 4 7
		1-2 days
		2

Factor		Total cases
After HSV-1 keratitis diagnosis	3 - 7 days	0
	7 - 14 days	0
	> 14 days	0

Note: BCVA, best-corrected visual acuity; HSV-1, herpes simplex virus-1