

## Labour Observation form

Start of observation:			Initials:		ID	
age:		pregnancy:		labour:		week:
height:		Weight before pregnancy:		Weight before labour:		

<b>induction:</b>	YES	NO	<b>Indication:</b>	
-------------------	-----	----	--------------------	--

On the admission:		
	USG	Leopold's manoeuvres:
Foetal positioning	<input type="checkbox"/> right <input type="checkbox"/> left <input type="checkbox"/> unable to determine	<input type="checkbox"/> right
		<input type="checkbox"/> left
Placenta location	<input type="checkbox"/> anterior <input type="checkbox"/> posteri <input type="checkbox"/> funda <input type="checkbox"/> right <input type="checkbox"/> left	<input type="checkbox"/> unable to determine

[illegible]

# Labour Observation form

Progression of foetal head in relations to the interspinal line												
epidural (+/- )												

Pelvis assessment:	
	<input type="checkbox"/> normal <input type="checkbox"/> abnormal

## Assessment of uterine contractions: (more than one answer possible)

- ☐ PROM (no uterine contractions)
 ☐ primary hypotonic uterine contractions
- ☐ secondary uterine inertia
 ☐ primary hypertonic uterine contractions
- ☐ discoordination of uterine contractions
 ☐ contractions of normal strenght, but shorter
- ☐ normal uterine contractions
- ☐ normal uterine contractions after oxytocine
- ☐ normal uterine contractions after amiotomy
- ☐ normal uterine contractions after spontaneous rupture of the membranes

How long was the uterine contractions (how many hours)							
irregular		every 10 minutes		every 5 minutes		every 2-3 minutes or more frequent	

Phases of labour (in minutes)			
Latent phase of the first stage (< 4cm)		First stage (from contractions every 5 min)	
Second stage		Active pushing in the second stage	

- ☐ Lack of progress in the first stage of labour (dilation)..... hours without progress.....
- ☐ Very slow progress

Lack of progress in the second stage of labour (more than 1 hours)

- ☐ yes, progression of the foetal head .....
- ☐ no

When the woman experiences the urge to push?

- ☐ no urge to push
- ☐ in the first stage of labour (dilation?) .....
- ☐ in the second stage of labour ( after how many minutes?) .....

If epidural was used, what were the indications?
--------------------------------------------------

# Labour Observation form

<input type="checkbox"/> woman's request as initially said in the birth plan		
Woman's request because: <input type="checkbox"/> pain too intense <input type="checkbox"/> woman not feeling well psychologically <input type="checkbox"/> lack of progress <input type="checkbox"/> suggested by the personnel		
Does epidural changes the engagement and the rotation of foetal head?		YES      NO
Where the pain was mostly located	<input type="checkbox"/> sacrum <input type="checkbox"/> back <input type="checkbox"/> thighs <input type="checkbox"/> lower abdomen <input type="checkbox"/> other.....	
<b>Delivery:</b>	<input type="checkbox"/> vaginal <input type="checkbox"/> vacuum <input type="checkbox"/> forceps <input type="checkbox"/> CS	
<b>Indications for instrumental or surgical delivery:</b>	<input type="checkbox"/> abnormal foetal heart pattern <input type="checkbox"/> lack of labour progress <input type="checkbox"/> prolonged second stage of labour <input type="checkbox"/> woman's exhaustion <input type="checkbox"/> lack of cooperation between woman and medical personnel <input type="checkbox"/> lack of personnel's patience <input type="checkbox"/> other.....	

Birth:						
Sex		weight		Apgar score		ICU
Does the external foetal head rotation was spontaneous?				YES		
				NO		
Assessment of perineal injury :				<input type="checkbox"/> First-degree tear <input type="checkbox"/> Second-degree tear <input type="checkbox"/> Third-degree tear <input type="checkbox"/> laceration of the vaginal wall <input type="checkbox"/> laceration of the vaginal mucosa <input type="checkbox"/> laceration of the periurethral labia <input type="checkbox"/> minor abrasion <input type="checkbox"/> other .....		
Assessment of perineal injury by a midwife:				<input type="checkbox"/> small <input type="checkbox"/> avarage <input type="checkbox"/> big		
<b>Blood loss</b>				<input type="checkbox"/> < 350 ml <input checked="" type="checkbox"/> 400-500 ml <input type="checkbox"/> >500 ml		
<b>Woman's attitude before labour</b>				<input type="checkbox"/> very positive <input type="checkbox"/> positive <input type="checkbox"/> neutral <input type="checkbox"/> negative <input type="checkbox"/> very negative		