

In vivo, in vitro and in silico Anticancer Activity of Ilama Leaves. An Edible and Medicinal Plant in Mexico.

Jesica Ramírez-Santos ^{1,2}, Fernando Calzada ^{2,*}, Rosa María Ordoñez-Razo ³, Jessica Elena Mendieta-Wejebe ¹, José Antonio Velázquez-Domínguez⁴, Raúl Argüello-García⁵, Claudia Velázquez⁶, and Elizabeth Barbosa¹

Supplementary material

Figure S1. Molecular Docking of the terpenoids bound to Bcl-2, (A) geranylgeraniol, (B) phytol, (C) farnesyl acetate and (D) navitoclax.

Figure S2. Molecular Docking of the terpenoids bound to Mcl-1, (A) geranylgeraniol, (B) phytol, (C) farnesyl acetate and (D) 9EA

Figure S3. Molecular Docking of the terpenoids bound to VEGFR-2, (A) geranylgeraniol, (B) farnesyl acetate, (C) phytol and (D) axitinib.




Figure S1. Molecular Docking of the terpenoids bound to Bcl-2, (A) geranylgeraniol, (B) phytol, (C) farnesyl acetate and (D) navitoclax




Figure S2. Molecular Docking of the terpenoids bound to Mcl-1, (A) geranylgeraniol, (B) phytol, (C) farnesyl acetate and (D) 9EA.




Figure S3. Molecular Docking of the terpenoids bound to VEGFR-2, (A) geranylgeraniol, (B) farnesyl acetate, (C) phytol and (D) axitinib.