

**Supplemental Table 2** Molecular indices for individual markers of *F. grandis* by sample site. *N*, No. of Sequences; *M*, No. of haplotypes, *h*, haplotypic diversity;  $\pi$ , nucleotide diversity; SD, standard deviation; *S*, No. of segregating (polymorphic) sites; *D*, Tajima's *D* neutrality test with probability value (*P*).

Locus	Site	M	<i>h</i> (SD)	$\pi$ (SD)	S	T <sub>s</sub>	T <sub>v</sub>	I/D
<b>CR-1</b>								
	Corpus Christi	11	0.849 (0.050)	0.0193 (0.011)	25	20	3	2
	Matagorda	13	0.954 (0.034)	0.0209 (0.012)	19	16	3	2
	Reitan Marsh	18	0.956 (0.022)	0.0242 (0.013)	31	28	2	2
	Sportsman's	21	0.977 (0.017)	0.0220 (0.012)	29	24	5	2
	Venice	13	0.757 (0.069)	0.0068 (0.004)	18	12	6	0
	Ocean Springs	17	0.843 (0.061)	0.0097 (0.006)	23	17	6	1
<b>ND-2</b>								
	Corpus Christi	11	0.895 (0.034)	0.0153 (0.009)	22	19	3	0
	Matagorda	13	0.954 (0.034)	0.0199 (0.011)	24	20	4	0
	Reitan Marsh	14	0.929 (0.025)	0.0187 (0.010)	23	22	1	0
	Sportsman's	17	0.960 (0.019)	0.0199 (0.011)	24	22	2	1
	Venice	8	0.768 (0.044)	0.0042 (0.003)	8	7	2	0
	Ocean Springs	17	0.928 (0.024)	0.0064 (0.004)	16	11	7	0
<b>ND-5</b>								
	Corpus Christi	14	0.935 (0.052)	0.017 (0.009)	21	19	1	1
	Matagorda	16	0.909 (0.040)	0.016 (0.009)	23	22	1	0
	Reitan Marsh	16	0.949 (0.025)	0.017 (0.009)	25	23	2	0
	Sportsman's	8	0.598 (0.084)	0.003 (0.002)	11	11	0	0
	Venice	12	0.722 (0.080)	0.004 (0.003)	15	15	0	0
	Ocean Springs	8	0.769 (0.059)	0.007 (0.004)	17	16	1	0