

Article

Transport Model of Rare Earth Elements in Weathering Crusts during Electrokinetic Mining

Gaofeng Wang ^{1,2,3}, Bowen Ling ^{4,5}, Xiaoliang Liang ^{1,2,3}, Jie Xu ^{1,2,3}, Shichang Kang ^{1,2,3}, Jingming Wei ^{1,2,3}, Wei Tan ^{1,2,3}, Runliang Zhu ^{1,2,3}, Jianxi Zhu ^{1,2,3,*} and Hongping He ^{1,2,3}

¹ CAS Key Laboratory of Mineralogy and Metallogeny/Guangdong Provincial Key Laboratory of Mineral Physics and Material, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China; wanggaofeng@gig.ac.cn (G.W.); liangxl@gig.ac.cn (X.L.); xujie777@gig.ac.cn (J.X.); kangshichang@gig.ac.cn (S.K.); weijm@gig.ac.cn (J.W.); tanwei@gig.ac.cn (W.T.); zhurl@gig.ac.cn (R.Z.); hehp@gig.ac.cn (H.H.)

² CAS Center for Excellence in Deep Earth Science, Guangzhou 510640, China

³ University of Chinese Academy of Sciences, Beijing 100049, China

⁴ Key Laboratory for Mechanics in Fluid Solid Coupling Systems, Institute of Mechanics, Chinese Academy of Sciences, Beijing 100190, China; lingbowen@imech.ac.cn

⁵ School of Engineering Science, University of Chinese Academy of Sciences, Beijing 100049, China

* Correspondence: zhujx@gig.ac.cn

Supplementary Material




Figure S1. (a) Variation of LREE/HREE ratio in the collected leachate with increasing EKM treatment time. (b) LREE/HREE ratio in weathering crust after the EKM treatment.




Figure S2. Electromigration velocities of REEs in weathering crust calculated based on the (a) conventional and (b) newly proposed models.