

Supplementary Materials

Table S1. Score frequency of lack of Accessibility - Availability in Health Care Services (LAAHCS) in relation to characteristics of 46,105 Europeans adults aged 50+ years.

			Score frequency of lack of Accessibility - Availability in Health Care Services ^a			
			0 n=40,728	1 to 24 n=4874	25+ n=503	
n			weighted %			p-value
Gender	♂	19,641	88.8	10.3	0.9	<0.001
	♀	26,464	85.5	13.3	1.1	
Age, years	50-59	4696	85.1	14.2	0.7	0.088
	60-69	16,208	87.4	11.6	1.0	
	70-79	15,601	88.2	10.6	1.2	
	80-89	8164	87.5	11.4	1.1	
	90-104	1436	84.6	14.3	1.1	
Education, years	0	1549	83.5	15.4	1.1	0.015
	1-7	6721	84.4	14.1	1.4	
	8-12	24,766	87.4	11.5	1.1	
	13+	13,069	88.5	11.0	0.5	
Family status	unmarried, divorced, widow	5911	83.5	14.2	2.3	<0.001
	married, living with partner	40,194	87.6	11.6	0.8	
Occupation	employed	9619	87.1	12.3	0.6	0.017
	unemployed, retired, housemaker	36,486	87.0	11.8	1.2	
Chronic conditions or diseases	none	11,921	89.4	10.0	0.6	<0.001
	1	12,890	87.0	11.8	1.2	
	2	10,085	87.2	12.0	0.8	
	3+	11,209	83.5	14.9	1.6	

^aScore range from 0 to 100, with a higher score indicating higher lack of Accessibility - Availability in Health Care Services.

χ^2 tests.