

Two *Ascophyllum nodosum* Fucoidans with Different Molecular Weights Inhibit
Inflammation via Blocking of TLR/NF- κ B Signaling Pathway Discriminately

Lilong Wang¹, Linlin Wang¹, Chunhong Yan¹, Chunqing Ai¹, Chengrong Wen¹, Xiaoming Guo²,
Shuang Song^{1,*}

¹ National Engineering Research Center of Seafood, School of Food Science and Technology,
Collaborative Innovation Center of Seafood Deep Processing, School of Food Science and
Technology, Dalian Polytechnic University, Dalian 116034, P. R. China

² Shenzhen Key Laboratory of Food Nutrition and Health, Institute for Advanced Study, Shenzhen
University, Shenzhen, 518060, P. R. China

* Corresponding author at School of Food Science and Technology, Dalian Polytechnic University,
No.1 Qinggongyuan, Ganjingzi District, Dalian 116034, P. R. China. Tel: +86-411-86323262, Fax:
+86-411-86323262. E-mail address: songs1008@163.com.

Table S1. Primer sequences used for quantitative RT-PCR reactions.

Gene symbol	Forward primer sequence	Reverse primer sequence
TLR2	CAGCTGGAGAACTCTGACCC	CAAAGAGCCTGAAGTGGGAG
TLR4	CAACATCATCCAGGAAGGC	GAAGGCGATACAATTCCACC
iNOS	ATGTCCGAAGCAAACATCAC	TAATGTCCAGGAAGTAGGTG
COCX2	CAGCAAATCCTTGCTGTTCC	TGGGCAAAGAATGCAAACATC
TNF- α	GATCGGTCCCCAAAGGGATG	GGCTACAGGCTTGTCACCTG
IL-1 β	TTCATCTTTGAAGAAGAGCCCAT	TCGGAGCCTGTAGTGCAGTT
IL-6	TGGAAATGAGAAAAGAGTTGTGC	CCAGTTTGGTAGCATCCATCA
IL-10	TACTCGGCAAACCTAGTGCG	GTGTCCCAACATTCATAATTGTCAGT
β -actin	TCAGCAAGCAGGAGTACGATG	AACGCAGCTCAGTAACAGTCC