

Article

Acuminatol and Other Antioxidative Resveratrol Oligomers from the Stem Bark of *Shorea acuminata*

Norhayati Muhammad ¹, Laily B. Din ¹, Idin Sahidin ², Siti Farah Hashim ³, Nazlina Ibrahim ³, Zuriati Zakaria ⁴ and Wan A. Yaacob ^{1,*}

¹ School of Chemical Sciences and Food Technology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, UKM Bangi 43600, Selangor D.E., Malaysia; E-Mails: norhayatimuhammad@ymail.com (N.M.); lbdin@ukm.my (L.B.D.)

² Faculty of Mathematics and Natural Sciences, Haluoleo University, Kendari 93232, Sulawesi Tenggara, Indonesia; E-Mail: sahidin02@yahoo.com

³ School of Biosciences and Biotechnology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, UKM Bangi 43600, Selangor D.E., Malaysia; E-Mails: farah_samura@yahoo.com (S.F.H.); nazlina@ukm.my (N.I.)

⁴ Malaysia-Japan International Institute of Technology, Universiti Teknologi Malaysia Kuala Lumpur, Jalan Semarak, Kuala Lumpur 54100, Malaysia; E-Mail: zuriz@ic.utm.my

* Author to whom correspondence should be addressed; E-Mail: wanyaa@ukm.my; Tel.: +603-8921-5424; Fax: +603-8921-5410.

Received: 1 June 2012; in revised form: 10 July 2012 / Accepted: 18 July 2012 /

Published: 30 July 2012

Abstract: A new resveratrol dimer, acuminatol (**1**), was isolated along with five known compounds from the acetone extract of the stem bark of *Shorea acuminata*. Their structures and stereochemistry were determined by spectroscopic methods, which included the extensive use of 2D NMR techniques. All isolated compounds were evaluated for their antioxidant activity using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity (RSA) and the β -carotene-linoleic acid (BCLA) assays, and compared with those of the standards of ascorbic acid (AscA) and butylated hydroxytoluene (BHT). All compounds tested exhibited good to moderate antioxidant activity in the DPPH assay (IC₅₀s 0.84 to 10.06 mM) and displayed strong inhibition of β -carotene oxidation (IC₅₀s 0.10 to 0.22 mM). The isolated compounds were evaluated on the Vero cell line and were found to be non-cytotoxic with LC₅₀ values between 161 to 830 μ M.


Keywords: Dipterocarpaceae; *Shorea acuminata*; resveratrol oligomer; acuminatol; antioxidant activity; cytotoxicity

1. Introduction

Plants from the Dipterocarpaceae, Gnetaceae and Vitaceae families are known as rich sources of resveratrol oligomers [1]. There have been reports of 275 new resveratrol oligomers from these species between 1995 and 2008 [2]. Many studies have suggested that these groups of constituents exhibited a range of biological activities [3] which included antioxidant [4–8], antimicrobial [9], anti-inflammatory [10], anti-hepatotoxicity [11], anti-tumor [12], cytotoxic effects [6] and other activities [2]. Previous antioxidative studies conducted on oligomers of vitisinols B, C, D; (+)- ϵ -viniferin, (–)-viniferol, ampelopsin C and (+)-vitisin C from the roots of *Vitis thunbergii* (Vitaceae) have shown that they have strong free RSA with IC₅₀ values between 2.8 and 6.6 μ M [1]. New resveratrol trimers and tetramer of wilsonols A, B, C and diviniferin B from *V. wilsonae* also exhibited potent antioxidant activities towards DPPH with IC₅₀ values of 103.5, 195.4, 182.2 and 175.3 μ M respectively [13]. Structure-activity relationship studies revealed that the DPPH RSA of resveratrol dimers isolated from *Cyperus longus* (Cyperaceae) of longusols A, B, C; longusone A and *trans*-scirpusins A, B was stronger (IC₅₀s 2.8 to 9.3 μ M) than those of the monomers piceatanol and resveratrol (IC₅₀s 11 and 24 μ M, respectively) [5]. The DPPH RSA study previously conducted on resveratrol oligomers isolated from *Parthenocissus laetevirens* (Vitaceae) of laetevirenols A and B containing an unusual phenanthrene moiety which exhibited much stronger antioxidant activities (38.4 and 37.3 μ M) compared to those without that moiety of laetevirenols C-E (110.8, 128.0 and 158.2 μ M) [7]. An unusual resveratrol hexamer of chunganenol isolated from *V. chunganensis*, which was composed of more than five monomers, exhibited much stronger DPPH RSA than two resveratrol trimers from the same species of (+)-gnetin H and (+)-amurensin G (with respective values of 37.3; 251.0 and 138.0 μ M) [8].

Shorea is the largest genus in the Dipterocarpaceae family. To date, about 26 resveratrol oligomers have been successfully isolated from this genus [1]. *Shorea acuminata* Dyer is a timber tree which is classified as light Red Meranti and locally known as Meranti Rambai Daun. The distribution of the species ranges from the Malay Peninsula to Sumatra and up to the Lingga Archipelago. It occurs on low-lying and well-drained land, but it is more abundant in hilly areas up to 300 m [14]. A previous phytochemical study on *Shorea acuminata* resin had resulted in the characterization of 2 α ,3 α -dihydroxyolean-12-en-28-oic, mangiferonic, 2 α -hydroxyursolic and asiatic acids [15]. In our ongoing search for resveratrol oligomers, we now report the isolation and structural elucidation of a new resveratrol oligomer derivative named acuminatol (**1**) and the five known compounds laevifonol (**2**) [16], (+)- α -viniferin (**3**) [17], shoreaketone (**4**) [18], vaticanol B (**5**) [19] and (–)-hopeaphenol (**6**) [20] (Figure 1). This is the first phytochemical report on resveratrol oligomers isolated from *S. acuminata* that incorporates their antioxidant activity against DPPH radical and BCLA, and also their cytotoxic property against Vero cells.

Figure 1. Structures of 1–6.



2. Results and Discussion

2.1. Structure Elucidation

Compound **1** was isolated as a yellow amorphous solid, mp 186–188 °C, $[\alpha]_D^{19} -42^\circ$ (c 0.00024, MeOH), which exhibited a molecular ion peak in the negative-ion high resolution ESIMS at $[M-H]^-$ m/z 469.1275 (calcd. 469.1293) attributable to the molecular formula $C_{28}H_{22}O_7$, which corresponded to a resveratrol dimer. This assumption was reinforced by the UV and IR absorption data, together with the 1H - and ^{13}C -APT NMR data, which was assigned by the interpretation of the HMQC, HMBC, 1H - 1H COSY and NOESY spectra (Table 1). UV absorption λ_{max} nm (MeOH): 282; IR (KBr) ν_{max} cm^{-1} : 3366 (OH), 2922 (C–H aliphatic), 1602, 1516 and 1449 (C=C aromatic), 1249 and 1175 (C–O oxyaryl), and 835 (*p*-disubstituted benzene).

Table 1. ^{13}C - and ^1H -NMR spectral data * of **1**.

Position	δ_{C}	δ_{H} mult. (J in Hz)	^{13}C - ^1H HMBC	^1H - ^1H COSY	^1H - ^1H NOESY
1a	129.9	-	3a(5a), 8a	-	
2a/6a	129.3	7.09 d (8.6)	7a	3a(5a)	8a, 14a, 7a, 3a(5a)
3a/5a	115.2	6.76 d (8.6)	OH _{4a}	2a(6a)	2a(6a), OH _{4a}
4a	158.9	-	3a(5a), OH _{4a}	-	-
7a	87.5	5.67 d (11.7)	2a(6a), 8a	8a	8a, 14a, 2a(6a)
8a	49.1	4.19 d (11.7)	7a, 14a	7a	7a, 2a(6a), 2b(6b)
9a	141.3	-	7a, 8a, 8b	-	-
10a	120.3	-	7b, 14a, 12a, OH _{11a}	-	-
11a	156.5	-	7b, 12a, OH _{11a}	-	-
12a	100.5	6.41 d (1.5)	14a, OH _{13a}	14a	14a, OH _{11a}
13a	157.7	-	OH _{13a}	-	-
14a	104.6	6.18 br s	OH _{13a}	12a	7a, 12a, 2a(6a), OH _{13a}
1b	131.9	-	7b, 3b(5b)	-	-
2b/6b	129.1	7.13 d (8.2)	2b(6b)	3b/5b	7b, 3b(5b), 8a
3b/5b	114.2	6.57 d (8.2)	OH _{4b}	2b(6b)	2b(6b)
4b	155.0	-	2b(6b), OH _{4b}	-	-
7b	45.0	5.46 br s	-	8b	8b, 2b(6b)
8b	71.0	5.09 br s	7b, 14b, 9a	7b	7b, 14b
9b	141.6	-	7b, 8a	-	-
10b	116.7	-	8a, 12b, 14b	-	-
11b	158.1	-	-	-	-
12b	95.4	6.11 d (1.8)	14b, OH _{13b}	14b	14b, OH _{13b}
13b	156.0	-	-	-	-
14b	106.4	6.94 d (1.8)	OH _{13b}	12b	8b, 12b, OH _{13b}
OH _{8b}	-	3.63		-	
OH _{11a}	-	8.57		-	
OH _{4a}	-	8.54		-	
OH _{13b}	-	8.23		-	
OH _{13a}	-	8.22		-	
OH _{4b}	-	8.05		-	

* ^{13}C - and ^1H -NMR spectra were obtained at 100 and 400 MHz (acetone- d_6), respectively.

The ^1H -NMR spectral data of **1** exhibited the presence of two sets of *ortho*-coupled aromatic protons which can be assigned to two 4-hydroxyphenyl groups [ring A1: δ 7.09 (2H, d, J = 8.6 Hz, H-2a and 6a), δ 6.76 (2H, d, J = 8.6 Hz, H-3a and 5a), and ring B1: δ 7.13 (2H, d, J = 8.2 Hz, H-2b and 6b), δ 6.57 (2H, d, J = 8.2 Hz, H-3b and 5b)], two sets of *meta*-coupled aromatic protons on two 1,2,3,5-tetrasubstituted benzene rings [ring A2: δ 6.18 (1H, br s, H-14a), δ 6.41 (1H, d, J = 1.5 Hz, H-12a), and ring B2: δ 6.94 (1H, d, J = 1.8 Hz, H-14b), δ 6.11 (1H, d, J = 1.8 Hz, H-12b)], five phenolic hydroxyl groups (δ 8.05, 8.22, 8.23, 8.54 and 8.57), and one aliphatic hydroxyl group (δ 3.63). The ^1H -NMR spectrum also showed two pairs of aliphatic methine protons coupled successively: H-7a (δ 5.67, d, J = 11.7) and H-8a (δ 4.19, d, J = 11.7); H-7b (δ 5.46, br s) and H-8b (δ 5.09, br s) as shown in Table 2. The large coupling constant of H-7a and H-8a (11.7 Hz) in compound **1** indicated that the protons were in *trans* orientation [21], which is also exhibited by its

stereoisomers, namely (+)-ampelopsin A (11.7 Hz) [22], (–)-hemsleyanol A (9.8 Hz) [17] and (+)-balanocarpol (9.3 Hz) [23]. On the other hand, H-7b and H-8b of **1** gave two broad singlets as those in (+)-balanocarpol [23], which proved that the two protons were in *cis* orientation. However, *trans*-H-7b~H-8b in (+)-ampelopsin A [22] and (–)-hemsleyanol A [17] gave coupling constants of 5.0 and 5.9 Hz for H-7b whereas H-8b produced broad singlet and broad doublet respectively.


Table 2. Chemical shifts of aliphatic proton pairs of H-7a~H-8a and H-7b~H-8b for stereoisomers **1**, (+)-ampelopsin A [22], (–)-hemsleyanol A [17] and (+)-balanocarpol [23].

Compound	δ_{H} mult. (J in Hz)			
	7a	8a	7b	8b
1 ^a	β : 5.67 (d, 11.7)	α : 4.19 (d, 11.7)	β : 5.46 (br s)	β : 5.09 (br s)
(+)-Ampelopsin A ^b	α : 5.77 (d, 11.7)	β : 4.17 (br d, 11.7)	α : 5.45 (d, 5.0)	β : 5.42 (br s)
(–)-Hemsleyanol A ^a	β : 5.75 (d, 9.8)	α : 5.41 (d, 9.8)	α : 5.07 (d, 5.9)	β : 4.76 (br d)
(+)-Balanocarpol ^a	β : 5.69 (d, 9.3)	α : 5.16 (br d, 9.3)	α : 4.90 (br s)	α : 5.40 (br s)

^a Measured in acetone-*d*₆ (400 MHz); ^b measured in acetone-*d*₆ (500 MHz).

Relative configuration of methine protons at C-8a, C-7b and C-8b on a cycloheptane ring in compound **1** can be further verified by the results of NOESY experiment. The absence of NOEs between H-8a and either H-7b or H-8b revealed that H-8a and H-7b~H-8b were in opposite sides of the cycloheptane ring. This also confirmed that both H-7b and H-8b were in the *cis* position. (+)-Balanocarpol where all three hydrogens were on the same side of the cycloheptane ring had significant NOEs between H-8a/H-7b and H-8a/H-8b [23]. Since H-7a and H-8a of **1** were *trans* based on the above ¹H-NMR spectral data, the relative configuration of methine protons at C-7a, C-8a, C-7b and C-8b were β , α , β and β . Other significant NOEs in support of these observations were between β H-7a/H-14a, β H-7b/ β H-8b and β H-8b/H-14b. In the HMBC spectrum (Table 1, Figure 2a), significant correlations were observed between C-2a(6a)/H-7a, C-8a/H-7a, C-9a/H-7a, C-9a/H-8a and C-10b/H-8a, indicated that a pair of benzylic methine protons of β H-7a and α H-8a was assigned to the protons on a dihydrofuran ring.

Figure 2. Key HMBC (a) and NOE (b) correlations for **1**.




There were significant correlations observed between C-9a/H-8b, C-9b/H-7b, C-1b/H-7b, C-11a/H-7b, C-10a/H-7b and C-8b/H-7b. These correlations proved that there was one more pair of methine protons

attached to C-7b (β H-7b) and C-8b (β H-8b). The relative stereo structure for compound **1** was confirmed as shown in Figure 1. It should be noted that the above three resveratrol dimers, which were stereoisomers to compound **1**, were previously isolated from *Ampelopsis brevipedunculata* var. *hancei* (Vitaceae) [(+)-ampelopsin A] [22], *H. parvifolia* (Dipterocarpaceae) [(+)-balanocarpol] [23] and *Shorea hemsleyana* (Dipterocarpaceae) [(-)-hemsleyanol A] [17].

All of the four stereoisomers shared a same basic planar structure. However, the chemical shifts for aliphatic protons of H-7a, H-8a, H-7b and H-8b of the four stereoisomers were markedly different as shown in Table 2. These differences have been suggested to be mainly due to the A1 and B1 rings in their energy-optimized conformations as in Figure 3.

Figure 3. Energy-optimized stereo structures of **1** (a), (+)-ampelopsin A (b), (-)-hemsleyanol A (c) and (+)-balanocarpol (d).



The NOE interactions between H-8a/H-2a(6a), H-7a/H-14a, H-8b/H-14b and H-8a/H-2b(6b) in compound **1** (Figure 3a) and (+)-ampelopsin A (Figure 3b) [22] lead to *cis* and *skew-cis* conformations, while the absence of NOE interaction between H-8a/H-2b(6b) in (+)-balanocarpol (Figure 3d) [23] would suggest that it was in a *trans* conformation. The absence of NOE interaction between H-8a/H-2b(6b) in the *skew-cis* conformation of (-)-hemsleyanol A (Figure 3c) [17], unlike that in the *skew-cis* conformation of (+)-ampelopsin A, was due to the relative positions of H-8a and B1 ring which were farther apart in the actual energy-optimized 3D-model of the molecule. It should be noted that the NOE correlations between H-2a(6a)/H-14a and H-7a/H-8a in compound **1** seem to suggest that the H-7a and H-8a were *cis* to each other with H-7a in α position. However, this contradicted with the large coupling constants between H-7a and H-8a of the *trans* in compound **1** (11.7 Hz) as well as in its stereoisomers of (+)-ampelopsin A (11.7 Hz) [22], (-)-hemsleyanol A (9.8 Hz) [17] and (+)-balanocarpol (9.3 Hz) [23]. The chemical shift for H-8a in compound **1** was more upfield (δ 4.19) like the one found in (+)-ampelopsin A (δ 4.17) [22] when compared to that in (-)-hemsleyanol A (δ 5.41) [17] and (+)-balanocarpol (δ 5.16) [23]. This strongly suggested that the A1 and B1 rings in compound **1** and (+)-ampelopsin A were in *cis* configuration. The chemical shifts for H-8a in (-)-hemsleyanol A and (+)-balanocarpol which appeared 1.23 and 0.98 ppm more downfield indicated that the A1 and B1 rings for the latter were in *trans* configuration (Table 2). Although the A1 and B1 rings for (-)-hemsleyanol A were in *cis* configuration, its B1 ring did not give any deshielding effect like in (+)-balanocarpol since those rings were in the same orientation as shown in Figure 3c,d. The H-8a in (-)-hemsleyanol A was more deshielded than that in (+)-balanocarpol because the former lay within the deshielding area up front to the A1 ring whereas this ring in the latter was a little tilted. Moreover NOE correlations [Table 1, Figure 2b] between H-8a/H-2a(6a) and

H-8a/H-2b(6b) in compound **1** positively suggested that its A1, H-8a and B1 were all on the same side of the plane. The results from the energy-optimized conformations as calculated by GaussView 5.0 and Gaussian 09W using DFT-6-31G-(d,p) method, have confirmed the above deductions (Figure 3). Therefore, **1** is a new stereoisomer of these three compounds.

2.2. Antioxidant Activity

2.2.1. DPPH Assay

As shown in Table 3, the scavenging activity of compounds **1** to **6** towards DPPH free radicals was expressed in terms of IC_{50} values. Since lower IC_{50} values indicated stronger ability of the compounds to act as DPPH radical scavengers, it was obvious that the positive controls were excellent DPPH radical scavengers with AscA and BHT exhibited average 6- and 4-fold higher scavenging activities ($IC_{50} = 0.68$ and 0.95 mM, respectively) compared to compounds **1** to **6** (IC_{50} s = 0.84 to 10.06 mM). Compound **5** exhibited excellent RSA with IC_{50} having no significant difference ($p > 0.05$) compared to positive controls of AscA and BHT. The antioxidant activity of all compounds and positive controls was in the following order (Table 3): AscA ~ **5** ~ BHT > **4** > **6** > **2** > **3** > **1**.

Table 3. Antioxidant and cytotoxic activities of compounds **1** to **6**.

Compound	DPPH radical scavenging activities (IC_{50} , mM)	BCLA method (IC_{50} , mM)	Cytotoxic activities on Vero cell lines (LC_{50} , μ M)
1	10.06 ± 0.05^f	0.18 ± 0.01^a	400
2	4.21 ± 0.23^d	0.22 ± 0.02^a	597
3	6.29 ± 0.05^e	0.18 ± 0.00^a	208
4	1.54 ± 0.10^b	0.11 ± 0.00^a	830
5	0.84 ± 0.02^a	0.10 ± 0.01^a	759
6	2.78 ± 0.16^c	0.10 ± 0.01^e	161
AscA ^z	0.68 ± 0.00^a	25.19 ± 1.74^b	-
BHT ^z	0.95 ± 0.05^a	0.09 ± 0.00^a	-

^{a-g} Mean within each column with different letters differ significantly ($p < 0.05$). Each value is presented as mean \pm SD ($n = 3$). ^z Positive reference standards; IC_{50} , 50% inhibition concentration.

These results, especially for the compounds **5**, **4**, **3** and **1**, were in agreement with the recent study on antioxidant activity which revealed that the RSA of the resveratrol oligomers was related to their structures. The multiple phenolic hydroxyl groups of 10, 8, 6 and 5 for the compounds **5**, **4**, **3** and **1** with four, three, three and two *para*-hydroxy groups, respectively, contributed to their being better hydrogen donors. Furthermore, the presence of the extensive double bond conjugation within the compounds which was responsible for electron delocalization has made them good as radical targets [3]. Compounds **6** and **2** with 10 and five aromatic-OHs, and four and two *para*-OHs of did not fit very well into the order. Other related studies by He *et al.* in 2009 [8] on eight stilbene oligomers from *Vitis chunganensis* (Vitaceae), that is, hexamer of chunganenol, tetramers of vitisin A and hopeaphenol (**6**), four trimers and one monomer revealed that chunganenol was the most active, as expected, followed by vitisin A, with hopeaphenol ranking fifth in the order.

2.2.2. BCLA Assay

The antioxidant activity of the compounds **1** to **6** as well as the positive controls BHT and AscA, as measured by the bleaching of β -carotene, are presented in Table 3. It was noted that in this assay, AscA exhibited low antioxidant activity (IC_{50} 25.19 mM) compared to BHT (IC_{50} 0.09 mM) and compounds **1** to **6** (IC_{50} 0.10 to 0.22 mM). These results suggest that AscA is a weak antioxidant despite the fact that it is a well-known, polar antioxidant. Our results are in agreement with the previous report which pointed out that AscA did not show its antioxidant activity under similar assay [24]. In this assay, all compounds exhibited antioxidant activity with no significant difference ($p > 0.05$) than the positive control BHT. These results agree with previous report [1], which indicated that most stilbenoids possess antioxidant activity because they had polyphenol functions in the molecules.

2.3. Cytotoxicity Assay

The results of cytotoxicity evaluation of compounds **1** to **6** as LC_{50} (mM) are shown in Table 3. The compounds were considered safe when their LC_{50} s are higher than 100 μ M [25]. All compounds tested possessed LC_{50} values more than 100 μ M (161 to 830 μ M).

3. Experimental

3.1. General

IR spectra were recorded on a Perkin Elmer GX FT-IR spectrophotometer (Waltham, MA, USA). UV spectra were measured on Shimadzu UV-160 (200–400 nm, Kyoto, Japan). 1H and ^{13}C -APT NMR spectra were recorded in acetone- d_6 using JEOL ECP400 spectrometer (400 and 100 MHz for 1H and ^{13}C ; Akishima, Japan). Mass spectra were measured in electron spray ionization mode on Bruker (micro TOF-Q; Bremen, Germany) LC-MS spectrometer (ESI-MS in negative mode, Dionex, Sunnyvale, CA, USA). Melting points were determined by Stuart SMP10 melting point apparatus (Burlington, VT, USA) and were uncorrected. Optical rotations were recorded on Jasco Polarimeter P-1020 (Easton, MD, USA) in MeOH. Vacuum liquid chromatography (VLC) was carried out on Si-gel 60 GF₂₅₄ (Merck, Darmstadt, Germany), radial chromatography (RC) was done on Si-gel 60 PF₂₅₄ (Merck) and TLC was performed on pre-coated silica gel (Merck, Kieselgel 60 F₂₅₄ 0.25 mm), and detected by UV light (254 nm) or by CeSO₄ spraying reagent followed by heating. All solvents used were of analytical grades. Absorbance values for BCLA and cytotoxic assays were measured on microplate reader (Labsystem Multiskan Multisoft, Basingstoke, UK) and DPPH assay on Shimadzu UVmini-1240 spectrophotometer (Kyoto, Japan). Dulbecco's Modified Eagle's Medium (DMEM) and fetal bovine serum (FBS) used to maintain and supplement the cell, respectively, were from Flowlab (North Ryde, Australia). Samples emulsifying was done on Branson 5200 (Los Angeles, CA, USA) sonicator.

3.2. Plant Material

The stem bark of *Shorea acuminata* was collected from Universiti Kebangsaan Malaysia (UKM) Forest Reserve, Bangi, in September 2009. A voucher specimen (UKMB 23520) has been deposited in the UKM Herbarium, and identified by Mr. Sani Miran.

3.3. Extraction and Isolation

The dried powder of stem bark of *Shorea acuminata* (1 kg) was macerated with acetone (3 × 5 L, 3 days each) at room temperature. The extract was concentrated using rotary evaporator to yield a brownish acetone extract (49.6 g, 4.96%) that was fractionated by VLC eluted with mixtures of *n*-hexane-EtOAc of increasing polarity. The eluates that showed similar profile on TLC chromatogram were combined to give five fractions A–E. Fraction C (1.5 g) was subjected to RC by eluting with CHCl₃-MeOH (9:1) to afford compound **3** (4.9 mg). Purification of fraction D (400 mg) by RC using CHCl₃-MeOH (8.4:1.6) followed by preparative TLC (CHCl₃-MeOH, 8.2:1.8) gave compounds **1** (12.0 mg), **2** (4.0 mg) and **4** (7.6 mg). Purification of fraction E (600 mg) by RC using CHCl₃-MeOH (8:2) followed by preparative TLC afforded compounds **5** (22.5 mg) and **6** (5.2 mg).

Compound 1: Yellow amorphous solid, m.p. 186–188 °C, $[\alpha]_D^{19} -42^\circ$ (c 0.00024, MeOH), UV (MeOH) λ_{\max} 282 nm; IR (KBr) ν_{\max} cm⁻¹: 3366, 2922, 1602, 1516 and 1449, 1249 and 1175, 835. ESIMS (neg): m/z 469.1275 [M-H]⁻, calcd: 469.1293; ¹H- and ¹³C-NMR data, see Table 1.

3.4. Antioxidant Assays

3.4.1. DPPH Assay

The antioxidant activity of all six compounds was determined by the DPPH radical scavenging method according to the previous procedure [26] with modification. Each compound was diluted two-fold in a series of five starting from 5.00 mg/mL. The solution of DPPH in methanol (6 × 10⁻⁵ M) was prepared daily before UV measurements. A 3 mL aliquot of this solution was mixed with 77 μL of the compound solution in a 1-cm path length microcuvette. The mixture was shaken vigorously. The mixtures were then kept in the dark for 15 min at room temperature and the decrease in absorption was measured immediately in a UV-Vis spectrophotometer at 515 nm. The blank solution containing the same amount of methanol and DPPH solution was prepared and its absorption was measured daily. The experiment was carried out in triplicate. AscA and BHT were used as positive controls. The RSA of samples, expressed as percentage inhibition of DPPH, was calculated according to the formula: Inhibition percentage (Ip) = $[(A_B - A_A)/A_B] \times 100$, where A_B and A_A are the absorbance values—checked after 15 min—of the blank sample and of the tested sample solutions, respectively. The IC₅₀ values, which represented the concentrations of the tested samples and standards that caused 50% RSA of DPPH, were calculated from the plots of inhibition percentages against concentrations.

3.4.2. BCLA Assay

This test was carried out according to the reported method of literature [27] with modification. Approximately 4 mL of a solution of β -carotene in chloroform (1 mg/mL) were pipetted into a flask containing 40 mg of linoleic acid and 400 mg of Tween-40. The chloroform was removed using a rotary evaporator at 40 °C for 5 min, and to the residue, 100 mL of distilled water were added, slowly with vigorous agitation, to form an emulsion. A 96-well micro-titer plate was added with 50 μ L of the Part 2.4.1 solution of the test compound and 200 μ L of the emulsion, and the absorbance was measured at 450 nm, immediately, against a blank consisting of the emulsion without β -carotene. The plate was allowed to stand at room temperature (20–23 °C), and the absorbance measurements were conducted again at 30 min intervals up to 120 min. All tests were carried out in triplicate. Stable antioxidants of AscA and BHT were used as positive controls. The antioxidant activity (AA) of the test samples was evaluated in terms of bleaching of β -carotene using the formula; $AA = [1 - (A_0 - A_t)/(A_0' - A_t')] \times 100$, where A_0 and A_0' are the absorbance values measured at zero time of the sample and the blank, respectively, and A_t and A_t' are the absorbance measured in the test sample and the blank, respectively, at times up to 120 min. The IC_{50} values, which represented the concentrations of the tested samples and standards that caused 50% bleaching of BCLA, were calculated from the plots of inhibition percentages against concentrations.

3.5. Cytotoxicity Assay

The isolated compounds were tested for *in-vitro* cytotoxicity using Vero cells by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay [28]. The Vero cell line was initiated from kidney of a normal adult African green monkey, *Cercopithecus aethiops*, obtained from Virology Laboratory, School of Biosciences and Biotechnology, Faculty of Science and Technology, UKM. Vero cells were maintained in DMEM, supplemented with 10% FBS and cultured at 37 °C in a humidified 5% CO₂ incubator. The concentration of stock compound solution was 1.0 mg/mL prepared by dissolving 1 mg compound in 50 μ L methanol and 950 μ L 5% FBS-DMEM. Sonicator was used to emulsify the compounds for 40 min before the two-fold dilutions made in 5% FBS-DMEM to produce a compound solution at concentrations of 0.5, 0.25, 0.125 and 0.0625 mg/mL. Briefly, a total of 50 mL of cell suspension with different concentrations were added to each well in the 96-well micro-titer plates. As a positive control, each well from first to third wells was added with cells of 2.5×10^5 , 1.25×10^5 and 0.625×10^5 cells/mL. For the negative control (100% cell death, LC_{100}), 50 mL of DMEM without cells were added to the twelfth well. Cells of 2.5×10^5 cells/mL were added into the fourth to eleventh wells. The plates were incubated for 48 h at 37 °C with 5% CO₂ until a monolayer is formed. A total of 50 μ L of test compound solution were placed in each well containing the monolayer cells and 50 μ L of phosphate buffered saline was added to each control cell well and negative control well (LC_{100}), and incubated for 24 h at 37 °C with 5% CO₂. After the incubation period, MTT (20 μ L, 5 mg/mL) was added into each well and the cells incubated for 2 to 4 h, until a purple precipitate was clearly visible under a microscope, the medium together with MTT (190 μ L) was aspirated off from the wells, DMSO (100 μ L) was added and the plate shaken for 5 min. The absorbance for each well was measured at 540 nm in a micro-titer plate reader and percentage of

cell viability (CV) was calculated manually using the formula: $CV = [(Ab_{S_{sample}} - Ab_{S_{negative\ control}}) / (Ab_{S_{cell}} - Ab_{S_{negative\ control}})] \times 100$. A dose-response curve was plotted to enable the calculation of the concentration that kills 50% of the Vero cells (LC_{50}).

3.6. Statistical Analysis

Values expressed are means of the three replicate determinations \pm standard deviation. All statistical analyses were carried out using SPSS 16.00 for Windows. To determine whether there were any differences between activities of samples, variance analysis (one-way ANOVA) was applied to the results. Values of $p < 0.05$ were considered as significant different ($\alpha = 0.05$).

4. Conclusions

The compounds **1** to **6** were found to be potent antioxidants, comparable in activity to the widely used synthetic antioxidant BHT in both assays. The activity was found to mostly increase with the number of phenolic units in the oligomer molecules. The non-toxic nature of all compounds **1** to **6** against normal (Vero) cells should merit further investigation to assess the effectiveness of these compounds in other biological activities including against other cell lines.

Supplementary Materials

Supplementary materials can be accessed at: <http://www.mdpi.com/1420-3049/17/8//9043/s1>.

Acknowledgments

We would like to thank the Ministry of Education Malaysia for PhD study leave of Norhayati Muhammad, Ministry of Higher Education Malaysia and Universiti Kebangsaan Malaysia (UKM) for financial support (Grant Codes: UKM-GUP-2011-205, UKM-OUP-KPB-31-156/2011 and UKM-DLP-2012-033). We are also very thankful to Centre for Research and Innovation Management, UKM for the technical support on LC-MS instrumentation. Thanks are also due to A. Hamid A. Hadi from the Chemistry Department, Faculty of Science, University of Malaya for kindly allowing us to use the polarimeter.

References

1. Xiao, K.; Zhang, H.-J.; Xuan, L.-J.; Zhang, J.; Xu, Y.-M.; Bai, D.-L. Stibenoids: Chemistry and bioactivities. *Stud. Nat. Prod. Chem.* **2008**, *34*, 453–646.
2. Shen, T.; Wong, X.-N.; Lou, H.-X. Natural stilbenes: An overview. *Nat. Prod. Rep.* **2009**, *26*, 916–935.
3. Lin, M.; Yao, C.-S. Natural oligostilbenes. *Stud. Nat. Prod. Chem.* **2006**, *33*, 601–644.
4. Quideau, S.; Deffieux, D.; Douat-Casassus, C.; Pouysegu, L. Plant polyphenols: Chemical properties, biological activities, and synthesis. *Angew. Chem. Int.* **2011**, *50*, 586–621.

5. Morikawa, T.; Xu, F.; Matsuda, H.; Yoshikawa, M. Structures of novel norstilbene dimer, longusone A, and three new stilbene dimers, longusols A, B, and C, with antiallergic and radical scavenging activities from Egyptian natural medicine *Cyperus longus*. *Chem. Pharm. Bull.* **2010**, *58*, 1379–1385.
6. Ito, T.; Tanaka, T.; Nakaya, K.; Iinuma, M.; Takahashi, Y.; Naganawa, H.; Ohyama, M.; Nakanishi, Y.; Bastow, K.F.; Lee, K.-H. A new resveratrol octamer, vateriaphenol A, in *Vateria indica*. *Tetrahedron Lett.* **2001**, *42*, 5909–5912.
7. He, S.; Wu, B.; Pan, Y.; Jiang, L. Stilbene oligomers from *Parthenocissus laetevirens*: Isolation, biomimetic synthesis, absolute configuration, and implication of antioxidative defense system in the plant. *J. Org. Chem.* **2008**, *73*, 5233–5241.
8. He, S.; Jiang, L.; Wu, B.; Li, C.; Pan, Y. Chunganenol: An unusual antioxidative resveratrol hexamer from *Vitis chunganensis*. *J. Org. Chem.* **2009**, *74*, 7966–7969.
9. Nitta, T.; Arai, T.; Takamatsu, H.; Inatomi, Y.; Murata, H.; Iinuma, M.; Tanaka, T.; Ito, T.; Asai, F.; Ibrahim, I.; *et al.* Antibacterial activity of extracts preprepared from tropical and subtropical plants on Methicillin-Resistant *Staphylococcus aureus*. *J. Health Sci.* **2002**, *48*, 273–276.
10. Chung, E.Y.; Roh, E.; Kwak, J.A.; Lee, H.S.; Lee, C.K.; Han, S.B.; Kim, Y. Alpha-viniferin suppresses the signal transducer and activation of transcription-1 (STAT-1)-inducible inflammatory genes in interferon-gamma-stimulated macrophages. *J. Pharmacol. Sci.* **2010**, *112*, 405–414.
11. Oshima, Y.; Namao, K.; Kamijou, A.; Matsuoko, S.; Nakano, M.; Terao, K.; Ohizumi, Y. Powerful hepatoprotective and hepatotoxic plant oligostilbenes, isolated from the oriental medicinal plant *Vitis coignetiae* (Vitaceae). *Experientia* **1995**, *51*, 63–66.
12. Ito, T.; Akao, Y.; Yi, H.; Ohgucgi, K.; Matsumoto, K.; Tanaka, T.; Iinuma, M.; Nozawa, Y. Antitumor effect of resveratrol oligomers against human cancer cell lines and the molecular mechanism of apoptosis induced by vaticanol C. *Carcinogenesis* **2003**, *24*, 1489–1497.
13. Jiang, L.; He, S.; Sun, C.; Pan, Y. Selective $^1\text{O}_2$ quenchers, oligostilbenes, from *Vitis wilsonae*: Structural identification and biogenetic relationship. *Phytochemistry* **2012**, *77*, 294–303.
14. Symington, C.F. *Foresters' Manual of Dipterocarps: Malayan Forest Records No. 16*, 2nd ed.; Forest Research Institute and Malaysian Nature Society: Kuala Lumpur, Malaysia, 2004; pp. 175–197.
15. Cheung, H.T.; Yan, T.C. Constituents of Dipterocarpaceae resins IV. Triterpenes of *Shorea acuminata* and *Shorea resina-nigra*. *Aust. J. Chem.* **1972**, *25*, 2003–2012.
16. Hirano, Y.; Kondo, R.; Sakai, K. Novel stilbenoids isolated from the heartwood of *Shorea laevifolia*. *J. Wood Sci.* **2003**, *49*, 53–58.
17. Ito, T.; Tanaka, T.; Ido, Y.; Nakaya, K.-I.; Iinuma, M.; Riswan, S. Stilbenoids isolated from stem bark of *Shorea hemsleyana*. *Chem. Pharm. Bull.* **2000**, *48*, 1001–1005.
18. Ito, T.; Furusawa, M.; Iliya, I.; Tanaka, T.; Nakaya, K.-I.; Sawa, R.; Kubota, Y.; Takahashi, Y.; Riswan, S.; Iinuma, M. Rotational isomerism of a resveratrol tetramer, shoreaketone, in *Shorea uliginosa*. *Tetrahedron Lett.* **2005**, *46*, 3111–3114.
19. Tanaka, T.; Ito, T.; Nakaya, K.; Iinuma, M.; Riswan, S. Oligostilbenoids in stem bark of *Vatica rassak*. *Phytochemistry* **2000**, *54*, 63–69.
20. Kawabata, J.; Fukushi, E.; Hara, M.; Mizutani, J. Detection of connectivity between equivalent carbons in a C₂ molecule using isotopomeric asymmetry: Identification of hopeaphenol in *Carex pumila*. *Magn. Reson. Chem.* **1992**, *30*, 6–10.

21. Karplus, M. Vicinal proton coupling in nuclear magnetic resonance. *J. Am. Chem. Soc.* **1963**, *85*, 2870–2871.
22. Oshima, Y.; Ueno, Y.; Hikino, H.; Yang, L.L.; Yen, K.Y. Ampelopsins A, B and C, new oligostilbenes of *Ampelopsis brevipedunculata* var. *hancei*. *Tetrahedron* **1990**, *46*, 5121–5126.
23. Tanaka, T.; Ito, T.; Ido, Y.; Son, T.-K.; Nakaya, K.; Iinuma, M.; Ohyama, M.; Chelladurai, V. Stilbenoids in the stem bark of *Hopea parviflora*. *Phytochemistry* **2000**, *53*, 1015–1019.
24. Siramon, P.; Ohtani, Y. Antioxidative and antiradical activities of *Eucalyptus camaldulensis* leaf oils from Thailand. *J. Wood Sci.* **2007**, *53*, 498–504.
25. Das, A.; Trousdale, M.D.; Ren, S.; Lien, E.J. Inhibition of herpes simplex virus type 1 and adenovirus type 5 by heterocyclic Schiff bases of aminohydroxyguanidine tosylate. *Antivir. Res.* **1999**, *44*, 201–208.
26. Miliauskas, G.; Venskutonis, P.R.; van Beek, T.A. Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food Chem.* **2004**, *85*, 231–237.
27. Jayaprakasha, G.K.; Singh, R.P.; Sakariah, K.K. Antioxidant activity of grape seed (*Vitis vinifera*) extracts on peroxidation models *in vitro*. *Food Chem.* **2001**, *73*, 285–290.
28. Fazari, G.M.; Azilawaty, A.; Nazlina, I.; Yaacob, W.A. Cytotoxic effect and anti-MRSA activity of methanolic extracts of *Phyllanthus gracilipes* and *Phyllanthus columnaris*. *Sains Malays.* **2011**, *40*, 457–466.

Sample Availability: Samples of the compounds **1–6** are available from the authors.

© 2012 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).