Molecules 1997, 2, M20

## [3S\*,5S\*,6S\*]-1,7-Dioxaspiro[5.5]undecane-3,5-diol

## Margaret A. Brimble\* and Andrew D. Johnston

Division of Organic Chemistry, School of Chemistry F11, The University of Sydney, N.S.W 2006, Australia. phone +61-2-9351-2750, fax +61-2-9351-6650, e-mail brimble\_m@alf.chem.su.oz.au, http://www.chem.usyd.edu.au/~brimble

Received: 18 June 1997 / Published: 20 June 1997

## **Scheme**

The reaction and the product ratio shown in the scheme was reported [1]. The two products (2 [1] and 3) were separated by flash chromatography using hexane-ethyl acetate (1:1) as eluent to afford the title compound 3 in the scheme) as a colourless glass (33 mg, 6%).

M.p. 63-64 deg.C.

High Res. MS calc. for C<sub>9</sub>H<sub>16</sub>O<sub>4</sub> M<sup>+</sup>H (CI, NH<sub>3</sub>) 189.1128, found: M<sup>+</sup> 189.1129.

IR (Nujol) cm<sup>-1</sup> 3610-3140 (br, s, OH), 1101 (s, C-O).

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>) 1.37 (1H, ddd,  $J_{11ax,11eq}$  12.7,  $J_{11ax}$ ,  $I_{10ax}$  12.7 and  $J_{11ax,10eq}$  4.7 Hz, 11ax-H), 1.49-1.70 (4H, m, 9-CH<sub>2</sub> and 10-CH<sub>2</sub>), 1.85-1.98 (3H, m, 11eq-H and 4-CH<sub>2</sub>), 2.51 (2H, m, 2 x O<u>H</u>), 3.39 (1H, dd,  $J_{2ax,2eq}$  10.4 and  $J_{2ax}$ ,  $I_{3ax}$  10.4 Hz,  $I_{3ax}$  10.4

<sup>13</sup>C-NMR (50 MHz, CDCl<sub>3</sub>) 18.3, 24.9, 29.9, 35.6 (CH<sub>2</sub>, C-4, C-9, C-10 and C-11), 60.8 (CH<sub>2</sub>, C-8), 61.9 (CH, C-3), 64.3 (CH<sub>2</sub>, C-2), 71.5 (CH, C-5), 95.7 (quat, C-6).

CI-MS 189 (M<sup>+</sup>H, 8%), 171 (M<sup>+</sup>H-H<sub>2</sub>O, 100), 153 (M<sup>+</sup>H-2H<sub>2</sub>O, 15), 111 (13), 101 (6).

*Acknowledgment*: The authors gratefully acknowledge financial support from the Australian Research Council and The University of Sydney.

## References

1. Brimble, M. A.; Johnston, A. D. Molecules 1997, M19.

Sample Availability: Available from Author, 0.1g and from MDPI, MDPI 11866.

© 1997 MDPI. All rights reserved.

1 von 1 08.05.2009 14:25