

Review

# The Rationale for Insulin Therapy in Alzheimer's Disease

Samo Ribarič

Institute of Pathophysiology, Faculty of Medicine, University of Ljubljana, Zaloška 4, SI-1000 Ljubljana, Slovenia; samo.ribaric@mf.uni-lj.si; Tel.: +386-1-543-7053; Fax: +386-1-543-7021

Academic Editors: Michael Decker and Diego Muñoz-Torrero

Received: 14 March 2016; Accepted: 19 May 2016; Published: 26 May 2016

**Abstract:** Alzheimer's disease (AD) is the most common form of dementia, with a prevalence that increases with age. By 2050, the worldwide number of patients with AD is projected to reach more than 140 million. The prominent signs of AD are progressive memory loss, accompanied by a gradual decline in cognitive function and premature death. AD is the clinical manifestation of altered proteostasis. The initiating step of altered proteostasis in most AD patients is not known. The progression of AD is accelerated by several chronic disorders, among which the contribution of diabetes to AD is well understood at the cell biology level. The pathological mechanisms of AD and diabetes interact and tend to reinforce each other, thus accelerating cognitive impairment. At present, only symptomatic interventions are available for treating AD. To optimise symptomatic treatment, a personalised therapy approach has been suggested. Intranasal insulin administration seems to open the possibility for a safe, and at least in the short term, effective symptomatic intervention that delays loss of cognition in AD patients. This review summarizes the interactions of AD and diabetes from the cell biology to the patient level and the clinical results of intranasal insulin treatment of cognitive decline in AD.

**Keywords:** Alzheimer's disease; diabetes; insulin; cognition

---

## 1. Amyloidogenesis in Alzheimer's Disease and Diabetes

The increase in life expectancy in the developed world is accompanied with an increased number of patients suffering from two age-related diseases, Alzheimer's disease (AD) and diabetes mellitus. AD is the most common form of dementia, with a prevalence that increases with age from 3% in people aged 65–74 to about 50% in people aged 85 or older. The worldwide number of patients with AD is projected to increase from 48 million in 2015 to more than 140 million in 2050 [1]. AD is associated with progressive memory loss, a gradual decline in cognitive function and a premature death 3–9 years after diagnosis [2]. Typical pathological features in the brain of AD patients are (a) intra-neuronal neurofibrillary tangles (NFT) of predominantly hyper phosphorylated tau protein; (b) extracellular deposition of senile plaques mainly composed of aggregated amyloid  $\beta$  (A $\beta$ ) peptide and (c) neuronal atrophy, starting in the entorhinal region and the temporal lobe, and progressing to the limbic system and major areas of the neocortex, as reviewed in [3–7]. Most patients with AD have a sporadic, late onset form, where the major risk factors are aging, type 2 diabetes (T2D) and apolipoprotein E  $\varepsilon 4$  (APOE- $\varepsilon 4$ ) [8–13]. The minority of AD patients have the early onset, genetic, familial form of AD due to the presence of autosomal dominant mutations in three genes: amyloid  $\beta$  precursor protein (A $\beta$ PP), presenilin-1 (PS1), and presenilin-2 (PS2) [14]. Therefore, the cause of disease in most AD patients is not known, what is known are the factors that may increase (e.g., aging, T2D) or decrease (e.g., mental and physical exercise) the risk and progression of AD.

Diabetes mellitus is one of the most prevalent metabolic disorders, with the total number of affected people projected to rise to 366 million in 2030 [15]. Diabetes mellitus is characterized by chronic hyperglycaemia and associated with long-term damage, dysfunction, and failure of various

organs, including the brain [16]. Diabetes has been associated with brain atrophy, white matter abnormalities, cognitive impairment, and is also a risk factor for dementia [17–21]. About 90% of diabetes patients have type T2D. Several studies concluded that T2D patients have an increased risk to develop dementia and AD [13,22,23]. It has been suggested that diabetes accelerates the progression of AD, rather than increasing the risk for AD [24]. This view was supported by *post-mortem* studies concluding that, compared to control subjects, senile plaques were less frequent and cerebrovascular pathology was more frequent in T2D patients [23,25]; in animal studies of diet-induced obesity with T2D cognitive impairment, brain atrophy, brain insulin resistance, neuro-inflammation, oxidative stress, and deficits in cholinergic function were relatively mild compared to the expected AD related pathology [26–28].

AD and T2D share the process of amyloidogenesis where a soluble protein forms insoluble fibrillary protein aggregates [29]. In AD, extracellular senile plaques in the brain are formed by abnormal protein processing of peptides of 30–51 amino acid residues by the proteolytic cleavage of amyloid  $\beta$  precursor protein (A $\beta$ PP) by  $\beta$ - and  $\gamma$ -secretases [30]. The most common soluble amyloid  $\beta$  peptides are A $\beta$ 42 and A $\beta$ 40; the former is produced by cleavage in the endoplasmic reticulum, the latter by cleavage in the trans-Golgi network [31–33]. It has been suggested that soluble amyloid  $\beta$  peptides present a negative feedback loop regulating synaptic plasticity and neuronal survival since low concentrations of A $\beta$  are present in the central nervous system of non-demented individuals [34]. Also, in cell culture studies, low concentrations of A $\beta$  were neurotrophic to undifferentiated hippocampal neurons and neurotoxic to mature neurons at higher concentrations [35]. Under pathological conditions, A $\beta$ 42 and A $\beta$ 40 form toxic soluble oligomers (A $\beta$ Os) that lead to cell death; A $\beta$ 42 is more susceptible to conformational changes than A $\beta$ 40 [36]. Studies of patients with AD, and of AD animal models, have linked A $\beta$ O with synaptic dysfunction, cognitive decline, inhibition of hippocampal long-term potentiation (LTP) component in memory, and learning and memory impairment [37–46]. In these studies, A $\beta$ Os were better correlated with dementia and synaptic loss than A $\beta$  in extracellular amyloid plaques [37,38]; however, toxicity mechanisms of amyloid and A $\beta$ Os were reported to be similar [47–49]. Several factors contribute to the progression of AD. What is not clear is the sequence of events that initiate the transition from a disease free state to an irreversible, progression of AD. Therefore, one of the key questions “*In a normal human brain, what causes the increase in soluble amyloid  $\beta$  peptides and their transition to A $\beta$ Os?*” has yet to be answered.

In human pancreatic islets of T2D patients, there is an accumulation of fibrillary protein aggregates of amylin—human islet amyloid polypeptide (hIAPP), a 37 residue peptide hormone, that is secreted from pancreatic  $\beta$ -cells in conjunction with insulin [50–52]. Human amylin forms oligomers that bind to A $\beta$ Os antibodies suggesting a common conformation [53–55]. The molecular structure and morphology of amylin fibrils resemble A $\beta$  fibrils of AD [56]. Amylin oligomers, the building blocks of amylin fibrils, induced  $\beta$ -cell apoptosis [57–59] and their toxic effect on cultured  $\beta$ -cells was similar to the A $\beta$ Os’ toxic effect in neurons [48,59–62]. Amylin accumulation in the pancreas is associated with a reduced  $\beta$ -cell volume and is present in the pancreatic islets of 90 % T2D patients [56,63]. Although several contributing factors have been suggested, the mechanism of transition from soluble amylin to toxic amylin aggregates is not known [64–71]. A step towards understanding the development of fibrillary protein aggregates of hIAPP was made by the study in isolated islets from hIAPP transgenic mice where amylin accumulated in a time- and glucose-concentration-dependent fashion and was associated with decreased  $\beta$ -cell areas and increased  $\beta$ -cell apoptosis [72,73]. Amylin oligomeric and plaque-like accumulations in brain parenchyma and cerebral vasculature were detected in T2D patients and in nondiabetic patients with late onset AD where amylin plaques were usually not co-localized with A $\beta$  plaques. The authors suggested that amylin amyloid formation in the wall of cerebral blood vessels could contribute to a reduced elimination of A $\beta$  from the brain, thus contributing to the progression of AD [74]. In an animal rat model, overexpressing human amylin in the pancreas, the elicited hyperamylinemia promoted accumulation of oligomerized amylin in the rat’s brain, associated

with an amylin-mediated brain inflammatory response, a reduced exploratory drive and a poor vestibulomotor performance on the rotarod test [75].

To summarize, T2D and AD are amyloid-forming diseases with insoluble protein aggregates in a fibrillary conformation that are caused by amylin deposition in pancreas and A $\beta$  deposition in brain, respectively. Amylin aggregation is associated with pancreatic  $\beta$ -cells loss, while A $\beta$  aggregation is associated with neuronal and synaptic dysfunction.

## 2. Interactions between Pathological Mechanisms of Alzheimer's Disease and Diabetes

The pathological mechanisms of AD and diabetes interact and tend to reinforce each other at the level of reduced cerebral blood flow and altered glucose metabolism, impaired insulin signalling, mitochondrial dysfunction, oxidative stress, advanced glycation end products, altered cholesterol metabolism, inflammation and cognitive impairment [3,5–7,16,76].

### 2.1. Cerebral Blood Flow and Glucose Metabolism

A decreased cerebral blood flow and a reduced brain glucose uptake occur during normal ageing of the human brain. These changes are more pronounced and occur sooner in life in patients with AD, in T1D and T2D patients. A reduced cerebral blood flow, by itself, promotes A $\beta$  accumulation in the brain and increased A $\beta$  brain levels promote local vasoconstriction, thus further promoting A $\beta$  accumulation.

#### 2.1.1. Cerebral Blood Flow and Glucose Metabolism in Alzheimer's Disease

**Human studies.** The normal ageing process of the brain is characterised by a progressive increase of morphologically abnormal capillaries, a decreased cerebral blood flow (CBF) and a lower brain glucose uptake and metabolism [77–80]. These changes develop sooner and to a greater degree in patients with AD [78–83]. PET studies associated impaired brain glucose metabolism with AD pathology, suggesting a causal link between impaired brain glucose metabolism and cognitive symptoms in AD patients [84]. A reduced brain glucose metabolism in AD patients in the early stages is most prominent in the posterior cingulate and parieto-temporal regions, and spreads to the prefrontal cortex in the advanced stages of the disease [85]. The CBF of AD patients was progressively reduced by 20% in the early stages of the disease and later by 55%–65% in the advanced stages of the disease [86]. Of interest is the finding that young and middle aged APOE- $\epsilon$ 4 allele carriers have normal cognition but also abnormally low rates of glucose metabolism, in the same brain regions as patients with probable AD, decades before the possible onset of dementia [87]. The glucose transporter GLUT-1 is significantly reduced in aged humans and in AD transgenic mice, coinciding with hippocampal atrophy [88]. In brains of AD patients, there was a negative correlation between on the one hand decreased GLUT-1 and -3 levels, decreased hypoxia-inducible factor 1-alpha (HIF-1 $\alpha$ ) and decreased O-GlcNAcylation and on the other hand hyper phosphorylation of tau protein and increased density of NFTs; concomitantly, the level of GLUT-2 was increased due to astrocyte activation [89].

**Animal model studies.** surgical reduction of CBF in an AD knock-in mouse model precipitated a positive-feedback cycle between brain A $\beta$  accumulation, cerebral amyloid angiopathy, amyloid plaque deposition and cognitive impairment on the one hand and CBF reduction on the other [90]. In a mouse model, the binding of A $\beta$  with the receptor for advanced glycation end products (RAGE) on the blood-brain barrier (BBB) triggered the release of vasoconstrictor endothelin-1 and pro-inflammatory factors thus reducing CBF [91].

#### 2.1.2. Cerebral Blood Flow and Glucose Metabolism in Diabetes

**Human studies.** A reduced cerebral glucose metabolism and insulin resistance were associated with memory deficits in pre-diabetic and T2D patients [92]. In human, insulin resistance was associated with a reduced cerebral glucose metabolism in frontal, temporo-parietal and cingulate regions in cognitively intact adults with prediabetes or T2D [92]. T1D patients, treated with insulin for several


years, had increased cerebrospinal fluid (CSF) concentrations of soluble low density lipoprotein receptor-related protein (LRP1) [93] that promoted the removal of A $\beta$  from the brain. T1D patients with recurrent hypoglycaemia and chronic hyperglycaemia were at an increased risk of cognitive decline [3].

**Animal model studies.** The expression of LRP1, a BBB transporter of A $\beta$  from the CSF into the blood, was down regulated in brain capillaries of streptozotocin-injected mice [94]. In streptozotocin injected mice, insulin reduced the concentration of RAGE in isolated brain micro vessels [95]. An animal model of diabetic AD mice, for investigating the links between T2D and AD, could be created with the combination of feeding a high-fat diet (HFD) to mice overexpressing A $\beta$ PP [5].

## 2.2. Impaired Insulin Signalling Links Systemic and Brain Oxidative Stress, Inflammation, Impaired Memory and Insulin Resistance in Diabetes and Alzheimer's Disease

Insulin modulates neurotransmitter release and synaptic plasticity, the basis for cognition, learning and memory [96–101]. Animal models and studies on patients have extensively documented impaired insulin signalling and degradation in AD and diabetes. Animal and cell culture models correlated reduced insulin signalling with increased activation of glycogen synthase kinase 3 beta (GSK3 $\beta$ ), hyper phosphorylation of tau protein, increased levels of A $\beta$  and cognitive deficiencies. In animal models, the dysregulated GSK3 activity contributed to both diabetes [102,103] and to AD [104]. Soluble A $\beta$  (40) is a competitive inhibitor of insulin binding to the insulin receptor (IR), and increased levels of this A $\beta$  could contribute to impaired insulin signalling and cognitive impairment in patients with AD. Both increased or decreased insulin blood levels can have detrimental effects on the progression of AD. In T2D, increased insulin blood levels promoted A $\beta$  accumulation by insulin competing with A $\beta$  for insulin degrading enzyme (IDE) [6,105]. IDE degrades both insulin [106] and A $\beta$  *in vivo* and *in vitro* [107,108]. Insulin deficiency, (in T1D or in the latter stages of T2D) attenuates insulin's and insulin-like growth factor-1's (IGF-1's) inhibition of A $\beta$ Os' binding to insulin receptor, thus reducing their protection of synapses from A $\beta$ Os' toxic effects [109,110].

Chronic, low-intensity systemic inflammation, for example during the ageing process or in T2D is characterised by increased blood levels of tumour necrosis factor  $\alpha$  (TNF $\alpha$ ) and interleukins 1 $\beta$  and 6. These peripherally released inflammatory mediators cross the blood–brain barrier and contribute to brain inflammation. Brain inflammation is further exacerbated by advanced glycation end products (AGE) and amyloid  $\beta$ -peptides (A $\beta$ s) in the brain that bind to RAGE and elicit the release of TNF $\alpha$ , IL1 $\beta$  and IL6 from microglia thus further increasing the brain levels of these inflammatory mediators. TNF $\alpha$  binds to its receptor on neurons and activates c-Jun N-terminal kinase (JNK). This kinase phosphorylates insulin receptor substrate 1 (IRS-1) at serine residues, which leads to the dissociation of IR from IRS1 and prevents further tyrosine phosphorylation of IRS-1 by the insulin-activated IR. The attenuated insulin signalling leads to reduced phosphoinositide 3-kinase (PI3K), protein kinase B (AKT) and mammalian target of rapamycin complex 1 (mTORC1) activities, which inhibit the development of synaptic plasticity and memory formation directly and also indirectly by an increased activity of glycogen synthase kinase 3 (GSK3). Brain insulin resistance in AD, due to chronic, low-intensity systemic inflammation, if further compounded by the presence of increased levels of A $\beta$ s and A $\beta$ Os in the brain. A $\beta$ s bind not only to RAGE on microglia, further promoting brain inflammation, but also to the IR, diminishing insulin signalling by preventing insulin binding. A $\beta$ Os' accumulation in the brain was associated with (a) increased brain levels of TNF $\alpha$ ; (b) removal of IRs from the synapses and their redistribution to the neuron's cell body; (c) decoupling of IRS-1 from IR in neurons that is mediated by I $\kappa$ B $\alpha$  kinase (IKK) and double-stranded RNA-dependent protein kinase (PKR) signalling and (d) aberrant binding to the N-methyl-D-aspartate receptors (NMDARs) that stimulated excessive Ca $^{2+}$  influx, increased oxidative stress and possibly activated protein tyrosine phosphatases that could further inhibit IRS-1 signalling [111–116]. The interactions between AD and T2D signalling pathways in the brain are summarized in Figure 1.



**Figure 1.** Interactions between AD and T2D signalling pathways in the brain. Abbreviations: A $\beta$  (amyloid  $\beta$  peptide); A $\beta$ O (toxic, soluble oligomer); A $\beta$ PP (amyloid  $\beta$  precursor protein); AGE (advanced glycation end product); AKT (protein kinase B); GSK3 (glycogen synthase kinase 3); IDE (insulin degrading enzyme); IKK ( $I\kappa B\alpha$  kinase); IL1 $\beta$  (interleukin 1 $\beta$ ); IL6 (interleukin 6); IR (insulin receptor); IRS-1 (insulin receptor substrate 1); JNK (c-Jun N-terminal kinase); LTP (long-term potentiation); mTORC1 (mammalian target of rapamycin complex 1); NMDAR (N-methyl-D-aspartate receptor); PI3K (phosphoinositide 3-kinase); PKR (double-stranded RNA-dependent protein kinase); RAGE (receptor for advanced glycation end products); SER (serine residue of IRS-1); T2D (type 2 diabetes); TNF $\alpha$  (tumour necrosis factor  $\alpha$ ); TNF $\alpha$ R (tumour necrosis factor  $\alpha$  receptor).

## 2.2.1. Impaired Insulin Signalling and Degradation in Alzheimer's Disease

**Human studies.** In AD patients, the insulin dose response curve for memory had an inverse U shaped function, with beneficial effects observed at the apex and null or negative effects when levels were too low or too high [117]. AD patients had reduced brain insulin receptor (IR) activity, lower CSF insulin levels and peripheral blood hyperinsulinemia [85,118] and an attenuated expression of IR and IGF receptors [119]. mRNA levels of insulin, IGF and their receptors were reduced in *post-mortem* human AD brains compared with controls [119] and this reduction was progressive with increasing severities of AD Braak Stage [120]. Insulin treatment, without changing fasting plasma glucose level, enhanced memory performance in AD patients [121,122].

**Animal model studies.** In a transgenic mouse model of AD, attenuated IR signalling reduced signalling through the phosphoinositide 3-kinase-protein kinase B (PI3K-Akt) pathway, increased activation of GSK3 $\beta$  and hyper phosphorylation of tau protein [123–125]. Overexpression of GSK3 $\beta$  in

the brain of transgenic mice was associated with an increased level of hyper phosphorylated tau and cognitive deficiencies [126]. Inhibition of GSK3 $\beta$  reduced A $\beta$  and hyperphosphorylated tau-associated neurodegeneration both *in vivo* and *in vitro* [127–130]. Cell culture models demonstrated that soluble A $\beta$  (40) is a competitive inhibitor of insulin binding to the IR [131,132].

## 2.2.2. Impaired Insulin Signalling and Degradation in Diabetes

**Animal model studies.** T1D animal model studies demonstrated: (1) an association between impaired cognitive performance and reduced hippocampal plasticity [133]; (2) a progressive impairment of cognitive function with an impairment of insulin and IGF-1 actions and neuronal apoptosis in hippocampus [134]; (3) mitigation of cognitive dysfunction and hippocampal apoptosis by proinsulin C-peptide with no concomitant effect on glucose levels [135]; (4) a loss of pancreatic  $\beta$  cells and long-term cognitive behaviour deficits in intra-cerebro-ventricularly streptozotocin treated rats [125,133,136]; (5) insulin treatment prevented streptozotocin induced deficits in the rat's cognition [99]; (6) the major contributing factor of T1D to AD was insulin deficiency that promoted increased tau protein phosphorylation in the cortex and hippocampus [137]. T2D animal model studies demonstrated that: (1) neuronal loss and neurite degeneration, associated with altered A $\beta$ PP metabolism, hyper phosphorylation of tau protein, and impaired signalling of insulin and IGF-1, were more severe in rat models of T2D than in rat models of T1D [138]; and (2) in a T2D animal model, the major contributing factor to AD was hyperglycaemia-mediated proteolytic tau cleavage since the cleaved tau served as a nucleation centre for the pathological assembly of tau filaments [137].

## 2.3. Mitochondrial Dysfunction in Alzheimer's Disease and Diabetes

Mitochondrial (MITO) dysfunction has a key role in the pathogenesis of AD, T1D and T2D [139,140]. Mitochondrial dysfunction precedes and sustains A $\beta$  accumulation in AD patients. In isolated human, rat and mouse MITO models, A $\beta$  (40 or 42) inhibited MITO enzymes cytochrome c oxidase and  $\alpha$ -ketoglutarate dehydrogenase, leading to MITO impairment [141–143]. Also A $\beta$  and A $\beta$ Os accumulated in mitochondria of transgenic mice overexpressing mutant A $\beta$ PP and in *post-mortem* brains from AD patients [142–145]. Studies on transgenic mice and AD patients confirmed that A $\beta$  can directly interact with MITO A $\beta$ -binding alcohol dehydrogenase (ABAD) leading to increased ROS generation, MITO dysfunction and cell death [146]; inhibition of the ABAD-A $\beta$  interaction in a mouse model attenuated A $\beta$  accumulation, conserved MITO function and improved spatial learning in an AD animal model [147]. In summary, the pathological basis of MITO dysfunction in AD, T1D and T2D is a combination of oxidative modifications of key MITO enzymes (e.g., pyruvate dehydrogenase, isocitrate dehydrogenase,  $\alpha$ -ketoglutarate dehydrogenase and cytochrome c oxidase), reduced antioxidant defences and an increased production of ROS.

### 2.3.1. Mitochondrial Dysfunction in Alzheimer's Disease

**Human studies.** The key findings on MITO dysfunction in AD were: (1) MITO dysfunction was observed in platelets and *post-mortem* brains of AD patients [148–154]; (2) MITO enzymes pyruvate dehydrogenase, isocitrate dehydrogenase, and  $\alpha$ -ketoglutarate dehydrogenase isolated from fibroblasts and brain tissue of AD patients were more susceptible to oxidative modification [155,156] and were altered by exposure to several pro-oxidants [157] and (3) in normal astrocytes, from primary human cortical foetal cell cultures, the inhibition of MITO metabolism by a MITO uncoupler induced amyloidogenic A $\beta$ PP processing and A $\beta$  accumulation seen in Down's syndrome of the brain [158].

**Animal model and cell culture studies.** The key findings on MITO dysfunction in AD were: (1) cells depleted of endogenous and repopulated with platelet MITO DNA from AD patients expressed MITO dysfunction, *i.e.*, reduced cytochrome c oxidase activity and enhanced ROS production [159]; (2) MITO dysfunction in cortices of transgenic AD mice preceded formation of amyloid plaque and NFT [160]; (3) in a transgenic AD mouse model, knockout of manganese superoxide dismutase, a major MITO antioxidant enzyme, increased A $\beta$  levels and amyloid plaque formation in the brain [161];

and (4) full-length A $\beta$ PP accumulated in MITO of cortical neurons and this accumulation precipitated MITO dysfunction [162].

### 2.3.2. Mitochondrial Dysfunction in Diabetes

**Human studies, animal model and cell culture studies.** Patients with T2D have altered MITO morphology and deficiency in bioenergetics and antioxidant capacity [163,164]. MITO dysfunction was demonstrated in animal models of diabetes [165–169]. In a T1D animal model, acute insulin-induced hypoglycaemia potentiated the detrimental effects of chronic hyperglycaemia in cortical and hippocampal MITO: the increase in ROS levels and decrease antioxidant defences [140]. Also, nerve damage, observed in an animal model of T1D, was causally linked to an increased, Ca<sup>2+</sup>-independent release of the excitatory amino acid glutamate during acute insulin-induced hypoglycaemia or during chronic hyperglycaemia [170].

## 2.4. Oxidative Stress

Increased oxidative stress is present in AD, T1D and T2D. In AD, oxidative stress precedes and coincides with A $\beta$  plaque formation, suggesting its role in initiating and sustaining AD-related changes in the brain. Oxidative stress promotes A $\beta$  production by upregulating  $\beta$ -secretase and  $\gamma$ -secretase expression [171–176]. A $\beta$  interacts with MITO proteins, disrupting the electron transport chain and promoting MITO dysfunction and an increased generation of ROS [177]. Oxidative stress also enhances tau hyper phosphorylation and subsequent NFT formation [178].

### 2.4.1. Oxidative Stress in Alzheimer's Disease

**Human studies and animal models.** Oxidative damage is increased in brain tissue samples from patients with AD [179–186]. Human autopsy brain samples from patients with AD and animal models of AD imply that oxidative damage occurs before A $\beta$  plaque formation [187–189]. In a transgenic AD mouse model, an increase in reactive nitrogen species coincided with the onset of A $\beta$  deposition [190].

### 2.4.2. Oxidative Stress in Diabetes

**Human studies, animal models and cell culture studies.** Clinical trials, as well as animal and cell culture models of diabetes, demonstrate that hyperglycaemia promotes an excessive and generalised production of free radicals in T1D and T2D [191–193].

## 2.5. Advanced Glycation End Products (AGEs)

The production of AGEs is enhanced by chronic hyperglycaemia in diabetes and by chronic oxidative stress present in diabetes and AD. A $\beta$  and AGEs bind to RAGE on microglial cells thus stimulating the release of proinflammatory mediators, (*i.e.*, free radicals and cytokines) and promoting the development of amyloid plaques and NFT [194–196]. AGEs and oxidative stress independently cause pathological changes on macromolecules and also act synergistically thus potentiating protein damage [197,198].

### 2.5.1. Advanced Glycation End Products in Alzheimer's Disease

**Human studies and cell culture studies.** In brain tissue from patients with AD, AGEs co-localized with NFT and amyloid plaques [199]. Also, AD plaques have more AGEs than healthy, age-matched controls [200]. It has been suggested that AGEs' accelerated the aggregation of both soluble A $\beta$  and tau thus facilitating development of NFTs and amyloid plaques [200,201]. In cultured neuroblastoma cells, AGEs promote neuronal oxidative stress and inflammation by nuclear factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa$ B) activation, increased cytokine IL6 gene expression and increased A $\beta$  release [202].

### 2.5.2. Advanced Glycation End Products in Diabetes

**Human studies.** In T2D patients, the formation and accumulation of AGEs, present in normal aging, is accelerated thus leading to diabetes associated damage on the retinal, renal, cardiovascular and peripheral nervous tissue [203–207]. Vascular AGEs staining is associated with cognitive impairment and a history of diabetes [208,209].

### 2.6. Cholesterol Metabolism

Increased blood cholesterol levels stimulate  $\beta$ - and  $\gamma$ -secretase proteolytic activity thus promoting A $\beta$  production and amyloid accumulation in the brain [210–212]. Increased blood cholesterol was also identified as an independent risk factor for AD [213]. In the brain, cholesterol is transported to the neurons by apolipoprotein E- $\epsilon$  (APOE- $\epsilon$ ), locally synthesized by astrocytes. The normal function of APOE- $\epsilon$  is essential not only for cholesterol catabolism but also for preventing AD-related brain changes. The isoform encoded by the ApoE- $\epsilon$ 4 allele promotes tau phosphorylation and inflammation thus contributing to the development of AD [214–217].

#### 2.6.1. Cholesterol Metabolism in Alzheimer's Disease

**Human studies.** Caucasian heterozygous and homozygous ApoE- $\epsilon$ 4 allele carriers have a three to eight-fold increased risk of AD compared to non ApoE- $\epsilon$ 4 carriers [8,218,219]. Also, MITO dysfunction in AD patients with ApoE- $\epsilon$ 4 allele correlate better with cognitive dysfunction, than in AD patients carrying the ApoE- $\epsilon$ 3 allele [220].

**Animal model studies.** The conclusions of cholesterol metabolism studies are: (1) the reduction of cholesteryl-ester levels, by inhibiting Acyl-CoA cholesterol acyltransferase (ACAT), reduces A $\beta$  production in cultured cells [221]; (2) in cell culture and whole animal mouse models of AD, ACAT1 inhibition reduces amyloid and tau deposition by enhancing autophagy [222–224]; (3) genetic ablation of ACAT in an AD mouse model increases cholesterol and 24(S)-hydroxycholesterol contents in the endoplasmic reticulum of mouse brain cells, attenuated human A $\beta$ PP harbouring the Swedish mutation and 3-hydroxy-3-methylglutaryl-CoA reductase protein contents and ameliorated A $\beta$  pathology [225] and (4) cholesterol depletion in rat hippocampal neurons decreases generation of A $\beta$  [226]. The implication of ApoE- $\epsilon$  in the development of AD was demonstrated by the following studies: (1) in transgenic mice, C-terminal truncated ApoE- $\epsilon$ 4 removed A $\beta$  with low efficiency and also acted synergistically with A $\beta$  to contribute to neuronal and behavioural deficits [227]; (2) carboxyl-terminal-truncated apolipoprotein  $\epsilon$ 4 caused Alzheimer's disease-like neurodegeneration and behavioural deficits in transgenic mice [228]; (3) ApoE contributed to the clearance of soluble A $\beta$  from the brain interstitial fluid of transgenic mice; the clearance was considerably lower in mice that expressed ApoE- $\epsilon$ 4 than in mice that expressed ApoE- $\epsilon$ 2 or ApoE- $\epsilon$ 3 [229]; in murine ApoE knockout mice, the expression of human ApoE- $\epsilon$ 4, but not of ApoE- $\epsilon$ 3, lead to deficits in learning and spatial memory, that increased with age and were seen primarily in females, although the A $\beta$  levels in female brains were comparable to male [230,231].

#### 2.6.2. Cholesterol Metabolism in Diabetes

**Human studies.** T2D patients with the ApoE- $\epsilon$ 4 allele are two-fold more likely to develop AD than nondiabetic ApoE- $\epsilon$ 4 carriers [232].

### 2.7. Inflammation

Inflammation is present in AD, T1D and T2D. Inflammation sustains insulin resistance, results in compensatory increase of insulin levels in T2D, and contributes to the destruction of pancreatic  $\beta$  cells in T1D [233–239]. In AD, inflammation promotes A $\beta$  deposition and tau hyper phosphorylation thus contributing to the progression of disease [240–251].

## Inflammation in Alzheimer's Disease

**Human studies.** *Post-mortem* human AD brains have an increased activation of inflammatory and immune pathways with upregulated levels of pro-inflammatory cytokines, chemokines and complement proteins [240]. This is consistent with microarray studies of brain samples from humans with AD or from animal models of AD that have identified an increased expression of genes involved in inflammation [242,243]. The intensity of inflammation varies over the course of AD. In patients with AD, cytokines in ventricular fluid were activated early but not late in the clinical course of AD; the observed reduced cytokine activation overlapped with reduced expressions of trophic factor and mediators of insulin signalling/responsiveness, and was concomitant with the increased brain levels of A $\beta$ , tau, and AGEs [241]. Treatment with nonsteroidal anti-inflammatory drugs (NSAIDs) was epidemiologically associated with a reduction in AD risk [246]; however, randomized trials failed to validate the benefit of several anti-inflammatory drugs in patients with AD [248–251].

**Animal model studies.** AD animal models have increased microglial activation and inflammation in the brain [244]. An AD animal model of A $\beta$ PP transgenic mice, demonstrated that inflammation promoted A $\beta$  deposition; an increased expression of  $\beta$ -site amyloid precursor protein cleaving enzyme 1 (BACE1) was associated with inflammation and this inflammation preceded A $\beta$  deposition [245]. Chronic administration of lipopolysaccharide lead to increased IL-1 levels and tau hyper phosphorylation in a triply transgenic AD mouse model [246].

### 2.8. Cognitive Impairment and Brain Insulin Sensitivity

The hippocampus is rich with insulin receptors that are localized mainly to nerve synapses [111,252]. It was demonstrated that insulin receptor signalling contributed to long-term memory consolidation and improved spatial learning in an animal model [253–255]. These observations are consistent with the suggested insulin involvement in the acquisition, consolidation and retrieval of memories [101]. Cognitive impairment is present in AD, T1D (due to insulin deficiency) and T2D (due to impaired insulin sensitivity of the brain) [256–258]. In patients with AD or mild cognitive impairment a single or repetitive administration of intranasal insulin improved memory and cognitive function [259–263]. When compared to the general population, T1D and T2D patients have a more pronounced brain atrophy [264–267]. The neuroprotective effects of insulin are mediated by (1) insulin binding to IGF receptors 1 and 2 [268] that attenuate GSK3 $\alpha$  activity [269] and reduce the neurotoxic effects of A $\beta$ PP [270–273]; (2) attenuation of A $\beta$ Os binding to neurons thus protecting synapses against the toxic effects of A $\beta$ Os [111] and (3) increasing transcription of antiamyloidogenic proteins insulin-degrading enzyme and  $\alpha$ -secretase and decreasing the transcription of pro-amylodogenic proteins A $\beta$ PP,  $\beta$ -secretase, and glycogen synthase kinase 3  $\alpha$  (Gsk3 $\alpha$ ) [274].

**Human studies.** Insulin infusion in normal older adults improves declarative memory and also increases CSF AB42 levels thus inhibiting A $\beta$ 42 intracellular accumulation by stimulating its extracellular secretion; improvement of declarative memory was attenuated in subjects with the highest, pre-insulin treatment levels of CSF A $\beta$ 42 [275]. Impaired insulin sensitivity of the brain was associated with (1) cognitive decline and brain atrophy in healthy elderly men and women [276]; (2) a positive correlation with the extent of cognitive impairment in AD patients [277,278] and (3) an increased risk for AD in women with elevated plasma insulin levels and decreased connectivity between the prefrontal cortex and hippocampus [279]. In *post-mortem* human brain samples of patients with AD the neuronal loss and impaired insulin/IGF signalling mechanisms correlates with the reduced expression of choline acetyltransferase [120].

## 3. Treatment of Alzheimer's Disease

The results of developing AD therapies on the basis of amyloid and NFTs hypotheses have been disappointing, even when the effective clearing of A $\beta$  deposits in AD brain was demonstrated [128,280–286]. Alternative therapies with antioxidants [287], anti-inflammatory

agents [288,289] were also ineffective. Current therapies of AD are symptomatic using NMDAR antagonists and cholinesterase inhibitors [7]. A personalised therapy approach for AD, based on clinical trials that selectively target different stages of AD, has been suggested as a way forward for promoting effective treatment. For example, at the pre-disease and pre-clinical stages of AD, treatment should be focused on managing factors that contribute to the onset of the disease (e.g., insulin resistance). For treating the pre-dementia stage, different combinations of preventive and curative drugs were suggested to stop the disease progression, and at the dementia stage, drugs targeting multiple pathogenic mechanisms of AD should be employed [7]. Combining different types of drugs, for personalised treatment of patients with AD, opens new challenges since it can lead to side effects and lower efficacy, as in the case of combined therapy with memantine and cholinesterase inhibitors [290].

#### 4. Antidiabetic Drugs for Treatment of AD

Insulin contributes to normal brain function, and insulin-signalling dysfunction accelerates the progression of AD as discussed in chapters 1 and 2.2. Therefore, therapeutic agents developed for the treatment of T2DM could be useful for treating AD. Drugs for treating T2D may affect the progression of AD brain changes either indirectly, by modifying the systemic blood concentrations of glucose, insulin, inflammatory markers and AGEs or directly in the brain, provided they pass the blood-brain barrier. Recent drug development for treating AD has focused on diabetes drugs that have a direct effect in the brain tissue since brain insulin resistance is often associated with AD [291].

##### 4.1. Peroxisome Proliferator-Activated Receptor- $\gamma$ Agonists

Thiazolidinediones (TZDs) diabetes drugs increase insulin sensitivity by activating the nuclear receptor peroxisome proliferator-activated receptor- $\gamma$ , thus increasing the expression of the glucose transporter GLUT-4. Two drugs, rosiglitazone and pioglitazone, are on the market for treatment of T2D; however, their use is restricted by their adverse side effects that include fluid retention, bone fractures and cardiovascular events [291,292]. Although initial studies in humans reported improved biomarkers of AD, improved memory and cognition these findings were not confirmed by larger clinical studies [293–296]. It was suggested that the potential long term beneficial effects of rosiglitazone, to ameliorate neuronal insulin resistance, were attenuated by its low blood-brain barrier penetration and a pronounced sensitizing effect on peripheral tissues to insulin, with a concomitant decrease in blood insulin levels that leads to a short term decrease in brain insulin signalling and worsening of cognitive impairment [297]. Two pilot studies are in progress and registered at ClinicalTrials.gov to evaluate pioglitazones efficacy for treating mild cognitive impairment due to AD: NCT01931566 with an estimated primary completion date of July 2019; and NCT02284906 with an estimated primary completion date of April 2021.

##### 4.2. Metformin

Metformin's glucose-lowering actions include increased glucose uptake in peripheral tissues and decreased liver gluconeogenesis by activating AMP-activated protein kinase (AMPK) in liver and other tissues. Activation of AMPK could increase insulin sensitivity through interactions with mTOR, P38 mitogen-activated protein kinases (p38 MAPK), and protein kinase C [291]. Animal studies support the suggestion that metformin penetrates the BBB and activates AMPK in the brain [298,299]. In isolated neuronal cells metformin sensitized neurons to insulin and prevented AD pathology in neurons chronically exposed to a hyperinsulinemic environment [300] but also increased  $\beta$ -secretase 1 (BACE1) transcription and generation of amyloid- $\beta$  [301]. Two pilot studies are registered at ClinicalTrials.gov to evaluate metformin's efficacy in humans: NCT00620191 that was completed but has not published the results and NCT01965756 with an estimated primary completion date of December 2015.

#### 4.3. Glucagon-Like Peptide-1 Receptor Agonists

Glucagon-like peptide-1 receptor agonists (GLP-1) is a hormone structurally unrelated to insulin. It circumvents the upstream stages of the insulin signalling pathway by binding to the G-protein-coupled GLP-1 receptor (GLP-1R) and activating a signalling pathway that converges with the downstream stages of the insulin signalling pathway thus facilitating insulin signalling [291]. GLP-1 stimulates adenylyl cyclase and modulates the activities of protein kinase A (PKA), phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K), mitogen-activated protein kinase (MAPK), protein kinase C (PKC), and protein kinase B (AKT) [291]. GLP-1 stimulates insulin secretion, decreases glucagon secretion and increases insulin sensitivity without significant hypoglycaemia [292]. GLP-1R is widely expressed in the brain and GLP-1R agonists with prolonged half-lives have been developed and approved for treatment of T2D (e.g., exenatide and liraglutide). The conclusion of preclinical studies of GLP-1R agonists in cell culture and animal models of AD was that GLP-1R agonists promoted synaptogenesis and neurogenesis, protected against oxidative injury, reduced A $\beta$ O and A $\beta$  plaque load, decreased microglial activation, and improved memory [302,303]. Two ongoing pilot studies are registered at ClinicalTrials.gov to evaluate GLP-1 agonists' efficacy in humans: NCT01255163 with an estimated study completion date of December 2018 and NCT01843075 with an estimated study completion date of January 2017.

#### 4.4. Leptin Analogues

Leptin receptors are abundant in areas of the brain involved in learning and memory and leptin deficiency was linked to cognitive impairment in human studies and in animal models [304,305]. Leptin reduced BACE1 activity, extracellular AB levels and tau phosphorylation in AD mouse models [306] and in humans, increased levels of circulating leptin were associated with a reduced incidence of AD [307]. At the time of writing no studies on the efficacy of leptin analogues for treatment of AD are posted on the ClinicalTrials.gov website.

#### 4.5. Amylin Analogues

Amylin readily crosses the BBB and the wide distribution of amylin brain receptors implicates amylin in a variety of brain functions, including memory, mood, anxiety, and satiety [291]. Amylin promotes normal blood glucose levels by delayed gastric emptying, decreased glucagon secretion and increased satiety. However, amylin oligomeric and plaque-like accumulations in brain parenchyma and cerebral vasculature were more frequent in patients with diabetes and nondiabetic patients with AD than in normal, nondiabetic control subjects. Amylin plaques were not co-localized with A $\beta$  plaques in most cases [74]. Pramlintide, an amylin analogue, is available as an adjunctive therapy for treatment of T1D and T2D and combines the beneficial glucose lowering effects of amylin without amylin's propensity to aggregate and form amylin oligomers and plaques and causes only a minimal hypoglycemia. Chronic infusion of pramlintide in an accelerated aging animal model of AD improved memory performance and decreased oxidative stress and inflammatory markers in the hippocampus [308]. Amylin and leptin activate overlapping signalling pathways that converge on the insulin-signalling pathway by activating AKT thus increasing insulin sensitivity. It was suggested that amylin sensitizes neurons to the effects of leptin, because amylin pre-treatment of neurons augmented leptin signalling [309]. Therefore, the synergy of amylin and leptin effects on brain cells could be used to develop combined therapies with amylin and leptin analogues for treatment of AD. At the time of this writing no studies on the efficacy of amylin analogues for treatment of AD are posted on the ClinicalTrials.gov website.

#### 4.6. Treatment of Alzheimer's Disease with Intranasal Insulin Application

The rationale for treating AD with intranasal insulin application was justified by the following research results: (1) insulin receptors were identified in several brain regions, with the highest

concentration in the olfactory bulb, hippocampus and hypothalamus [101,310]; (2) with intranasal application, insulin by-passes the BBB and directly enters the brain [311]; (3) insulin was detected in biologically relevant concentrations in the CSF in 30–40 min after intranasal application [312]; (4) intranasal insulin administration was associated with only minor side effects like dizziness or mild rhinitis [313–316]; (5) AD is associated with brain insulin resistance and insulin deficiency (reduced brain and CSF levels), with or without concomitant systemic insulin resistance or T2DM; (6) patients with diabetes, who were successfully managed with insulin, had a significantly improved memory, an attenuated progression of AD and lower densities of AD-related lesions; and (7) insulin therapy improved memory and cognition in patients with AD [317,318]. The positive effects of insulin therapy diminish with the progression of AD when increased A $\beta$  levels promote brain insulin resistance [319].

Repeated elevations of brain insulin concentrations, by intranasal administration, were associated with reduced A $\beta$  [122], no changes in blood pressure [320], and attenuated hypothalamus-pituitary-adrenal (HPA) secretory activity [259]. Intranasal insulin treatment, for early AD or mild cognitive impairment (MCI), improved memory and attention abilities in four phase 2 clinical trials without significant adverse effects or changes in blood levels of insulin or glucose [117,122,260,263,321–325]. Intranasal insulin administration elicited changes in peripheral glucose metabolism, but no significant change in blood insulin levels were detected [326–329]. There is some concern, that the efficacy of long-term treatment of AD patients with intranasal insulin application could be attenuated by long-term brain insulin receptor desensitisation or decreased efficacy of peripheral mechanisms (*i.e.*, the “sink hypothesis” for A $\beta$  lowering strategies) that contribute to insulin-induced AB clearance from the brain [119,268,277,330,331].

Most of the clinical trials of insulin treatment of AD used the regular, short half-life insulin [122]. An alternative is the insulin analogue detemir with a slower absorption and a higher lipophilicity compared to regular insulin [332–334]. Compared to regular insulin, insulin detemir was equal or more effective at reducing hyperglycaemia and nocturnal hypoglycaemic episodes [335] and more effective in eliciting insulin-signalling in the hypothalamus and cerebro-cortical tissue [336] thus generally affecting brain functions to a greater extent than regular insulin [337].

#### Clinical Trials of Alzheimer’s Disease Treatment with Intranasal Insulin Application

All three clinical trials presented below were randomised, blind and placebo-controlled. No treatment-related severe adverse effects were reported in any of these trials.

##### (a) Single dose trial with regular insulin

The participants were 59 normal adults and 33 memory impaired patients with AD, divided in two subgroups: apolipoprotein E- $\epsilon$ 4 allele carriers (ApoE- $\epsilon$ 4+) and apolipoprotein E- $\epsilon$ 4 allele non-carriers (ApoE- $\epsilon$ 4–). Participants received either a placebo or a single dose of 10, 20, 40 or 60 IU of insulin. The changes in plasma insulin, glucose, A $\beta$ 42 and A $\beta$ 40 levels after insulin treatment were: (1) plasma A $\beta$ 40 in normal adults or AD patients was not affected by insulin dosing; (2) plasma insulin and glucose levels in normal adults or AD patients were not influenced by the intervention; (3) plasma A $\beta$ 42 levels increased with the increase in insulin dose in ApoE- $\epsilon$ 4– patients; (4) in ApoE- $\epsilon$ 4+ patients, there was no significant change in A $\beta$ 42 levels with increased insulin concentrations; (5) in normal ApoE- $\epsilon$ 4– adults, the plasma A $\beta$ 42 levels showed a U shaped response curve to increasing insulin doses; (6) in normal ApoE- $\epsilon$ 4+ adults increasing insulin concentrations had no effect on plasma A $\beta$ .

The effects of insulin treatment on cognition were ApoE- $\epsilon$ 4(+ / –) dependent: (1) ApoE- $\epsilon$ 4– AD patients showed improved verbal memory at the optimal concentration of 20 IU; the insulin-dose response curve for memory had an inverse U shaped function, with beneficial effects observed at the apex and null or negative effects when levels were too low or too high and (2) ApoE- $\epsilon$ 4+ AD patients had a decline in verbal memory [117].

(b) Four-months treatment with 20 or 40 IU of regular insulin per day

The participants were 64 adults with amnestic mild cognitive impairment (MCI) and 40 adults with mild to moderate AD. Participants received daily either a placebo, or 20, or 40 IU of insulin for 4 months. The observed changes in A $\beta$ 42 cerebrospinal fluid (CSF) levels after insulin treatment were: (1) in both groups of adults, the CSF A $\beta$ 42 levels were slightly but not significantly lowered; (2) increased CSF A $\beta$ 42 levels were associated with improved delayed memory and preserved caregiver-rated functional ability; (3) decreased tau protein to A $\beta$ 42 ratios during the study period were correlated with improved delayed story recall and better daily function; (4) participants receiving 20 or 40 IU dose insulin showed a reduced progression of hypo metabolism in discrete areas of the cerebral cortex compared with the placebo group.

The effect of insulin treatment on cognition was partially insulin-dose dependent: (1) compared to placebo, treatment with 20 IU, but not with 40 IU, improved delayed memory in both groups of patients and (2) treatment with both insulin doses preserved caregiver-rated functional ability and general cognition in both groups of patients [321].

(c) Twenty-one-days treatment with 20 IU or 40 IU of insulin detemir twice daily

The participants were 60 adults with MCI and 20 adults with mild to moderate AD. Both groups had either ApoE- $\epsilon$ 4+ or ApoE- $\epsilon$ 4– adults. Participants received either a placebo, or 20, or 40 IU of insulin detemir twice daily for 21 days.

The effects of insulin detemir treatment on cognition were: (1) 20 IU insulin detemir treatment did not improve cognitive outcomes in any of the participants; (2) 40 IU insulin detemir treatment did not improve executive functioning or caregiver-rated daily functioning in any of the participants; (3) 40 IU insulin detemir treatment improved: verbal memory for ApoE- $\epsilon$ 4+ adults with MCI or AD; visuospatial and verbal working memory for ApoE- $\epsilon$ 4+ and APOE- $\epsilon$ 4– participants; and peripheral insulin resistance for ApoE- $\epsilon$ 4+ adults; and (4) 40 IU insulin detemir treatment increased peripheral insulin resistance in APOE- $\epsilon$ 4– participants [316].

## 5. Conclusions

Alzheimer's disease (AD) is the clinical manifestation of altered proteostasis, the process of accumulating misfolded, pathological proteins in the brain. The cause of altered proteostasis in most AD patients is not known. Known are some of the modulating factors that increase or decrease the risk of AD. The progression of AD is accelerated by several chronic disorders, among which the contribution of diabetes mellitus to AD is well understood at the cell biology level. At present, there is no effective cure for AD, only symptomatic interventions are available. To optimise symptomatic treatment, a personalised therapy approach that selectively targets different stages of AD, has been suggested. Intranasal insulin administration seems to have opened the possibility for a safe and effective symptomatic intervention that delays loss of cognition in AD patients, at least for the short term.

**Acknowledgments:** This work was supported by ARRS grant number P3-0171.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

1. Pahnke, J.; Walker, L.C.; Scheffler, K.; Krohn, M. Alzheimer's disease and blood-brain barrier function—Why have anti-beta-amyloid therapies failed to prevent dementia progression? *Neurosci. Biobehav. Rev.* **2009**, *33*, 1099–1108. [[CrossRef](#)] [[PubMed](#)]
2. Querfurth, H.W.; LaFerla, F.M. Alzheimer's disease. *N. Engl. J. Med.* **2010**, *362*, 329–344. [[CrossRef](#)] [[PubMed](#)]
3. Correia, S.C.; Santos, R.X.; Carvalho, C.; Cardoso, S.; Candeias, E.; Santos, M.S.; Oliveira, C.R.; Moreira, P.I. Insulin signaling, glucose metabolism and mitochondria: Major players in Alzheimer's disease and diabetes interrelation. *Brain Res.* **2012**, *1441*, 64–78. [[CrossRef](#)] [[PubMed](#)]

4. Freiherr, J.; Hallschmid, M.; Frey, W.H.; Brünner, Y.F.; Chapman, C.D.; Hölscher, C.; Craft, S.; De Felice, F.G.; Benedict, C. Intranasal insulin as a treatment for Alzheimer's disease: A review of basic research and clinical evidence. *CNS Drugs* **2013**, *27*, 505–514. [CrossRef] [PubMed]
5. Vandal, M.; Bourassa, P.; Calon, F. Can insulin signaling pathways be targeted to transport A $\beta$  out of the brain? *Front. Aging Neurosci.* **2015**, *7*. [CrossRef] [PubMed]
6. Yang, Y.; Song, W. Molecular links between Alzheimer's disease and diabetes mellitus. *Neuroscience* **2013**, *250*, 140–150. [CrossRef] [PubMed]
7. Chen, Z.; Zhong, C. Decoding Alzheimer's disease from perturbed cerebral glucose metabolism: Implications for diagnostic and therapeutic strategies. *Prog. Neurobiol.* **2013**, *108*, 21–43. [CrossRef] [PubMed]
8. Corder, E.H.; Saunders, A.M.; Strittmatter, W.J.; Schmeichel, D.E.; Gaskell, P.C.; Small, G.W.; Roses, A.D.; Haines, J.L.; Pericak-Vance, M.A. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* **1993**, *261*, 921–923. [CrossRef] [PubMed]
9. Hoyer, S. Causes and consequences of disturbances of cerebral glucose metabolism in sporadic Alzheimer disease: Therapeutic implications. *Adv. Exp. Med. Biol.* **2004**, *541*, 135–152. [PubMed]
10. Irie, F.; Fitzpatrick, A.L.; Lopez, O.L.; Kuller, L.H.; Peila, R.; Newman, A.B.; Launer, L.J. Enhanced risk for Alzheimer disease in persons with type 2 diabetes and APOE epsilon4: The Cardiovascular Health Study Cognition Study. *Arch. Neurol.* **2008**, *65*, 89–93. [CrossRef] [PubMed]
11. Kivipelto, M.; Helkala, E.L.; Laakso, M.P.; Hanninen, T.; Hallikainen, M.; Alhainen, K.; Iivonen, S.; Mannermaa, A.; Tuomilehto, J.; Nissinen, A.; et al. Apolipoprotein E epsilon4 allele, elevated midlife total cholesterol level, and high midlife systolic blood pressure are independent risk factors for late-life Alzheimer disease. *Ann. Intern. Med.* **2002**, *137*, 149–155. [CrossRef] [PubMed]
12. Luchsinger, J.A.; Reitz, C.; Patel, B.; Tang, M.X.; Manly, J.J.; Mayeux, R. Relation of diabetes to mild cognitive impairment. *Arch. Neurol.* **2007**, *64*, 570–575. [CrossRef] [PubMed]
13. Ott, A.; Stolk, R.P.; van Harskamp, F.; Pols, H.A.; Hofman, A.; Breteler, M.M. Diabetes mellitus and the risk of dementia: The Rotterdam Study. *Neurology* **1999**, *53*, 1937–1942. [CrossRef] [PubMed]
14. Rocchi, A.; Pellegrini, S.; Siciliano, G.; Murri, L. Causative and susceptibility genes for Alzheimer's disease: A review. *Brain Res. Bull.* **2003**, *61*, 1–24. [CrossRef]
15. Wild, S.; Roglic, G.; Green, A.; Sicree, R.; King, H. Global prevalence of diabetes: Estimates for the year 2000 and projections for 2030. *Diabetes Care* **2004**, *27*, 1047–1053. [CrossRef] [PubMed]
16. De la Monte, S.M.; Tong, M. Brain metabolic dysfunction at the core of Alzheimer's disease. *Biochem. Pharmacol.* **2014**, *88*, 548–559. [CrossRef] [PubMed]
17. Akisaki, T.; Sakurai, T.; Takata, T.; Umegaki, H.; Araki, A.; Mizuno, S.; Tanaka, S.; Ohashi, Y.; Iguchi, A.; Yokono, K.; Ito, H. Cognitive dysfunction associates with whitematter hyperintensities and subcortical atrophy onmagnetic resonance imaging of the elderly diabetes mellitus Japanese elderly diabetes intervention trial (J-EDIT). *Diabetes Metab. Res. Rev.* **2006**, *22*, 376–384. [CrossRef] [PubMed]
18. Den Heijer, T.; Vermeer, S.E.; van Dijk, E.J.; Prins, N.D.; Koudstaal, P.J.; Hofman, A.; Breteler, M.M. Type 2 diabetes and atrophy of medial temporal lobe structures on brain MRI. *Diabetologia* **2003**, *46*, 1604–1610. [CrossRef] [PubMed]
19. Manschot, S.M.; Brands, A.M.; van der Grond, J.; Kessels, R.P.; Algra, A.; Kappelle, L.J.; Biessels, G.J. Brain magnetic resonance imaging correlates of impaired cognition in patients with type 2 diabetes. *Diabetes* **2006**, *55*, 1106–1113. [CrossRef] [PubMed]
20. Schmidt, R.; Launer, L.J.; Nilsson, L.G.; Pajak, A.; Sans, S.; Berger, K.; Breteler, M.M.; de Ridder, M.; Dufouil, C.; Fuhrer, R.; et al. Magnetic resonance imaging of the brain in diabetes: The Cardiovascular Determinants of Dementia (CASCADE) Study. *Diabetes* **2004**, *53*, 687–692. [CrossRef] [PubMed]
21. Toth, C.; Martinez, J.; Zochodne, D.W. RAGE, diabetes, and the nervous system. *Curr. Mol. Med.* **2007**, *7*, 766–776. [CrossRef] [PubMed]
22. Kroner, Z. The relationship between Alzheimer's disease and diabetes: Type 3 diabetes? *Altern. Med. Rev.* **2009**, *14*, 373–379. [PubMed]
23. Janson, J.; Laedtke, T.; Parisi, J.E.; O'Brien, P.; Petersen, R.C.; Butler, P.C. Increased risk of type 2 diabetes in Alzheimer disease. *Diabetes* **2004**, *53*, 474–481. [CrossRef] [PubMed]
24. Pasquier, F.; Boulogne, A.; Leys, D.; Fontaine, P. Diabetes mellitus and dementia. *Diabetes Metab.* **2006**, *32*, 403–414. [CrossRef]

25. Nelson, P.T.; Smith, C.D.; Abner, E.A.; Schmitt, F.A.; Scheff, S.W.; Davis, G.J.; Keller, J.N.; Jicha, G.A.; Davis, D.; Wang-Xia, W.; *et al.* Human cerebral neuropathology of Type 2 diabetes mellitus. *Biochim. Biophys. Acta* **2009**, *1792*, 454–469. [[CrossRef](#)] [[PubMed](#)]
26. Lyn-Cook, L.E., Jr.; Lawton, M.; Tong, M.; Silbermann, E.; Longato, L.; Jiao, P.; Mark, P.; Wands, J.R.; Xu, H.; de la Monte, S.M. Hepatic ceramide may mediate brain insulin resistance and neurodegeneration in type 2 diabetes and non-alcoholic steatohepatitis. *J. Alzheimers Dis.* **2009**, *16*, 715–729. [[PubMed](#)]
27. Winocur, G.; Greenwood, C.E. Studies of the effects of high fat diets on cognitive function in a rat model. *Neurobiol. Aging* **2005**, *26*, 46–49. [[CrossRef](#)] [[PubMed](#)]
28. Tong, M.; Longato, L.; de la Monte, S.M. Early limited nitrosamine exposures exacerbate high fat diet-mediated type2 diabetes and neurodegeneration. *BMC Endocr. Dis.* **2010**, *10*. [[CrossRef](#)]
29. Chiti, F.; Dobson, C.M. Protein misfolding, functional amyloid, and human disease. *Annu. Rev. Biochem.* **2006**, *75*, 333–366. [[CrossRef](#)] [[PubMed](#)]
30. Kirkitadze, M.D.; Kowalska, A. Molecular mechanisms initiating amyloid beta-fibril formation in Alzheimer's disease. *Acta Biochim. Pol.* **2005**, *52*, 417–423. [[PubMed](#)]
31. Hardy, J.; Selkoe, D.J. The amyloid hypothesis of Alzheimer's disease: Progress and problems on the road to therapeutics. *Science* **2002**, *297*, 353–356. [[CrossRef](#)] [[PubMed](#)]
32. Selkoe, D.J. Translating cell biology into therapeutic advances in Alzheimer's disease. *Nature* **1999**, *399*, A23–A31. [[CrossRef](#)] [[PubMed](#)]
33. Shoji, M.; Golde, T.E.; Ghiso, J.; Cheung, T.T.; Estus, S.; Shaffer, L.M.; Cai, X.D.; McKay, D.M.; Tintner, R.; Frangione, B.; *et al.* Production of the Alzheimer amyloid beta protein by normal proteolytic processing. *Science* **1992**, *258*, 126–129. [[CrossRef](#)] [[PubMed](#)]
34. Tanzi, R.E.; Bertram, L. Twenty years of the Alzheimer's disease amyloid hypothesis: A genetic perspective. *Cell* **2005**, *120*, 545–555. [[CrossRef](#)] [[PubMed](#)]
35. Yankner, B.A.; Duffy, L.K.; Kirschner, D.A. Neurotropic and neurotoxic effects of amyloid b protein: Reversal by tachykinin neuropeptides. *Science* **1990**, *250*, 279–282. [[CrossRef](#)] [[PubMed](#)]
36. Williams, T.L.; Choi, J.K.; Surewicz, K.; Surewicz, W.K. Soluble Prion Protein Binds Isolated Low Molecular Weight Amyloid- $\beta$  Oligomers Causing Cytotoxicity Inhibition. *ACS Chem. Neurosci.* **2015**, *6*, 1972–1980. [[CrossRef](#)] [[PubMed](#)]
37. Lue, L.F.; Kuo, Y.M.; Roher, A.E.; Brachova, L.; Shen, Y.; Sue, L.; Beach, T.; Kurth, J.H.; Rydel, R.E.; Rogers, J. Soluble amyloid beta peptide concentration as a predictor of synaptic change in Alzheimer's disease. *Am. J. Pathol.* **1999**, *155*, 853–862. [[CrossRef](#)]
38. McLean, C.A.; Cherny, R.A.; Fraser, F.W.; Fuller, S.J.; Smith, M.J.; Beyreuther, K.; Bush, A.I.; Masters, C.L. Soluble pool of Abeta amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease. *Ann. Neurol.* **1999**, *46*, 860–866. [[CrossRef](#)]
39. Lambert, M.P.; Barlow, A.K.; Chromy, B.A.; Edwards, C.; Freed, R.; Liosatos, M.; Morgan, T.E.; Rozovsky, I.; Trommer, B.; Viola, K.L.; *et al.* Diffusible, nonfibrillar ligands derived from Abeta1–42 are potent central nervous system neurotoxins. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 6448–6453. [[CrossRef](#)] [[PubMed](#)]
40. Shankar, G.M.; Li, S.; Mehta, T.H.; Garcia-Munoz, A.; Shepardson, N.E.; Smith, I.; Brett, F.M.; Farrell, M.A.; Rowan, M.J.; Lemere, C.A.; *et al.* Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat. Med.* **2008**, *14*, 837–842. [[CrossRef](#)] [[PubMed](#)]
41. Townsend, M.; Shankar, G.M.; Mehta, T.; Walsh, D.M.; Selkoe, D.J. Effects of secreted oligomers of amyloid beta-protein on hippocampal synaptic plasticity: A potent role for trimers. *J. Physiol.* **2006**, *572*, 477–492. [[CrossRef](#)] [[PubMed](#)]
42. Wang, H.W.; Pasternak, J.F.; Kuo, H.; Ristic, H.; Lambert, M.P.; Chromy, B.; Viola, K.L.; Klein, W.L.; Stine, W.B.; Kraft, G.A.; *et al.* Soluble oligomers of beta amyloid (1–42) inhibit long-term potentiation but not long-term depression in rat dentate gyrus. *Brain Res.* **2002**, *924*, 133–140. [[CrossRef](#)]
43. Cleary, J.P.; Walsh, D.M.; Hofmeister, J.J.; Shankar, G.M.; Kuskowski, M.A.; Selkoe, D.J.; Ashe, K.H. Natural oligomers of the amyloid-beta protein specifically disrupt cognitive function. *Nat. Neurosci.* **2005**, *8*, 79–84. [[CrossRef](#)] [[PubMed](#)]
44. Lesne, S.; Koh, M.T.; Kotilinek, L.; Kayed, R.; Glabe, C.G.; Yang, A.; Gallagher, M.; Ashe, K.H. A specific amyloid-beta protein assembly in the brain impairs memory. *Nature* **2006**, *440*, 352–357. [[CrossRef](#)] [[PubMed](#)]

45. Poling, A.; Morgan-Paisley, K.; Panos, J.J.; Kim, E.M.; O'Hare, E.; Cleary, J.P.; Lesne, S.; Ashe, K.H.; Porritt, M.; Baker, L.E. Oligomers of the amyloid-beta protein disrupt working memory: Confirmation with two behavioral procedures. *Behav. Brain Res.* **2008**, *193*, 230–234. [CrossRef] [PubMed]
46. Shankar, G.M.; Bloodgood, B.L.; Townsend, M.; Walsh, D.M.; Selkoe, D.J.; Sabatini, B.L. Natural oligomers of the Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J. Neurosci.* **2007**, *27*, 2866–2875. [CrossRef] [PubMed]
47. Kawahara, M.; Kuroda, Y.; Arispe, N.; Rojas, E. Alzheimer's beta-amyloid, human islet amylin, and prion protein fragment evoke intracellular free calcium elevations by a common mechanism in a hypothalamic GnRH neuronal cell line. *J. Biol. Chem.* **2000**, *275*, 14077–14083. [CrossRef] [PubMed]
48. Anguiano, M.; Nowak, R.J.; Lansbury, P.T., Jr. Protofibrillar islet amyloid polypeptide permeabilizes synthetic vesicles by a pore-like mechanism that may be relevant to type II diabetes. *Biochemistry* **2002**, *41*, 11338–11343. [CrossRef] [PubMed]
49. Lim, Y.A.; Ittner, L.M.; Lim, Y.L.; Gotz, J. Human but not rat amylin shares neurotoxic properties with Abeta42 in long-term hippocampal and cortical cultures. *FEBS Lett.* **2008**, *582*, 2188–2194. [CrossRef] [PubMed]
50. Clark, A.; Nilsson, M.R. Islet amyloid: A complication of islet dysfunction or an aetiological factor in Type 2 diabetes? *Diabetologia* **2004**, *47*, 157–169. [CrossRef] [PubMed]
51. Haataja, L.; Gurlo, T.; Huang, C.J.; Butler, P.C. Islet amyloid in type 2 diabetes, and the toxic oligomer hypothesis. *Endocr. Rev.* **2008**, *29*, 303–316. [CrossRef] [PubMed]
52. Cooper, G.J. Amylin compared with calcitonin gene-related peptide: Structure, biology, and relevance to metabolic disease. *Endocr. Rev.* **1994**, *15*, 163–201. [CrossRef] [PubMed]
53. Green, J.D.; Goldsbury, C.; Kistler, J.; Cooper, G.J.; Aeby, U. Human amylin oligomer growth and fibril elongation define two distinct phases in amyloid formation. *J. Biol. Chem.* **2004**, *279*, 12206–12212. [CrossRef] [PubMed]
54. Porat, Y.; Kolusheva, S.; Jelinek, R.; Gazit, E. The human islet amyloid polypeptide forms transient membrane-active prefibrillar assemblies. *Biochemistry* **2003**, *42*, 10971–10977. [CrossRef] [PubMed]
55. Kayed, R.; Head, E.; Thompson, J.L.; McIntire, T.M.; Milton, S.C.; Cotman, C.W.; Glabe, C.G. Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science* **2003**, *300*, 486–489. [CrossRef] [PubMed]
56. Luca, S.; Yau, W.M.; Leapman, R.; Tycko, R. Peptide conformation and supramolecular organization in amylin fibrils: Constraints from solid-state NMR. *Biochemistry* **2007**, *46*, 13505–13522. [CrossRef] [PubMed]
57. Janson, J.; Ashley, R.H.; Harrison, D.; McIntyre, S.; Butler, P.C. The mechanism of islet amyloid polypeptide toxicity is membrane disruption by intermediate-sized toxic amyloid particles. *Diabetes* **1999**, *48*, 491–498. [CrossRef] [PubMed]
58. Konarkowska, B.; Aitken, J.F.; Kistler, J.; Zhang, S.; Cooper, G.J. The aggregation potential of human amylin determines its cytotoxicity towards islet beta-cells. *FEBS J.* **2006**, *273*, 3614–3624. [CrossRef] [PubMed]
59. Meier, J.J.; Kayed, R.; Lin, C.Y.; Gurlo, T.; Haataja, L.; Jayasinghe, S.; Langen, R.; Glabe, C.G.; Butler, P.C. Inhibition of human IAPP fibril formation does not prevent beta-cell death: Evidence for distinct actions of oligomers and fibrils of human IAPP. *Am. J. Physiol. Endocrinol. Metab.* **2006**, *291*, E1317–E1324. [CrossRef] [PubMed]
60. Huang, C.J.; Lin, C.Y.; Haataja, L.; Gurlo, T.; Butler, A.E.; Rizza, R.A.; Butler, P.C. High expression rates of human islet amyloid polypeptide induce endoplasmic reticulum stress mediated beta-cell apoptosis, a characteristic of humans with type 2 but not type 1 diabetes. *Diabetes* **2007**, *56*, 2016–2027. [CrossRef] [PubMed]
61. Kayed, R.; Sokolov, Y.; Edmonds, B.; McIntire, T.M.; Milton, S.C.; Hall, J.E.; Glabe, C.G. Permeabilization of lipid bilayers is a common conformation-dependent activity of soluble amyloid oligomers in protein misfolding diseases. *J. Biol. Chem.* **2004**, *279*, 46363–46366. [CrossRef] [PubMed]
62. Mirzabekov, T.A.; Lin, M.C.; Kagan, B.L. Pore formation by the cytotoxic islet amyloid peptide amylin. *J. Biol. Chem.* **1996**, *271*, 1988–1992. [PubMed]
63. Clark, A.; Wells, C.A.; Buley, I.D.; Cruickshank, J.K.; Vanhegan, R.I.; Matthews, D.R.; Cooper, G.J.; Holman, R.R.; Turner, R.C. Islet amyloid, increased A-cells, reduced B-cells and exocrine fibrosis: Quantitative changes in the pancreas in type 2 diabetes. *Diabetes Res.* **1988**, *9*, 151–159.

64. Kahn, S.E.; Andrikopoulos, S.; Verchere, C.B. Islet amyloid: A long-recognized but underappreciated pathological feature of type 2 diabetes. *Diabetes* **1999**, *48*, 241–253. [CrossRef] [PubMed]
65. Westerman, P.; Engstrom, U.; Johnson, K.H.; Westerman, G.T.; Betsholtz, C. Islet amyloid polypeptide: Pinpointing amino acid residues linked to amyloid fibril formation. *Proc. Natl. Acad. Sci. USA* **1990**, *87*, 5036–5040. [CrossRef] [PubMed]
66. Marzban, L.; Trigo-Gonzalez, G.; Verchere, C.B. Processing of pro-islet amyloid polypeptide in the constitutive and regulated secretory pathways of beta cells. *Mol. Endocrinol.* **2005**, *19*, 2154–2163. [CrossRef] [PubMed]
67. Hou, X.; Ling, Z.; Quartier, E.; Foriers, A.; Schuit, F.; Pipeleers, D.; Van Schravendijk, C. Prolonged exposure of pancreatic beta cells to raised glucose concentrations results in increased cellular content of islet amyloid polypeptide precursors. *Diabetologia* **1999**, *42*, 188–194. [CrossRef] [PubMed]
68. Marzban, L.; Rhodes, C.J.; Steiner, D.F.; Haataja, L.; Halban, P.A.; Verchere, C.B. Impaired NH<sub>2</sub>-terminal processing of human proislet amyloid polypeptide by the prohormone convertase PC2 leads to amyloid formation and cell death. *Diabetes* **2006**, *55*, 2192–2201. [CrossRef] [PubMed]
69. Paulsson, J.F.; Westerman, G.T. Aberrant processing of human proislet amyloid polypeptide results in increased amyloid formation. *Diabetes* **2005**, *54*, 2117–2125. [CrossRef] [PubMed]
70. Janson, J.; Soeller, W.C.; Roche, P.C.; Nelson, R.T.; Torchia, A.J.; Kreutter, D.K.; Butler, P.C. Spontaneous diabetes mellitus in transgenic mice expressing human islet amyloid polypeptide. *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 7283–7288. [CrossRef] [PubMed]
71. Verchere, C.B.; D'Alessio, D.A.; Palmiter, R.D.; Weir, G.C.; Bonner-Weir, S.; Baskin, D.G.; Kahn, S.E. Islet amyloid formation associated with hyperglycemia in transgenic mice with pancreatic beta cell expression of human islet amyloid polypeptide. *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 3492–3496. [CrossRef] [PubMed]
72. Zraika, S.; Hull, R.L.; Udayasankar, J.; Utzschneider, K.M.; Tong, J.; Gerchman, F.; Kahn, S.E. Glucose- and time-dependence of islet amyloid formation *in vitro*. *Biochem. Biophys. Res. Commun.* **2007**, *354*, 234–239. [CrossRef] [PubMed]
73. Zraika, S.; Hull, R.L.; Udayasankar, J.; Aston-Mourney, K.; Subramanian, S.L.; Kisilevsky, R.; Szarek, W.A.; Kahn, S.E. Oxidative stress is induced by islet amyloid formation and time-dependently mediates amyloid-induced beta cell apoptosis. *Diabetologia* **2009**, *52*, 626–635. [CrossRef] [PubMed]
74. Jackson, K.; Barisone, G.A.; Diaz, E.; Jin, L.W.; DeCarli, C.; Despa, F. Amylin deposition in the brain: A second amyloid in Alzheimer disease? *Ann. Neurol.* **2013**, *74*, 517–526. [CrossRef] [PubMed]
75. Srodninski, S.; Sharma, S.; Bachstetter, A.B.; Brelsfoard, J.M.; Pascual, C.; Xie, X.S.; Saatman, K.E.; Van Eldik, L.J.; Despa, F. Neuroinflammation and neurologic deficits in diabetes linked to brain accumulation of amylin. *Mol. Neurodegener.* **2014**, *22*, 9. [CrossRef] [PubMed]
76. de la Monte, S.M. Contributions of brain insulin resistance and deficiency in amyloid-related neurodegeneration in Alzheimer's disease. *Drugs* **2012**, *72*, 49–66. [CrossRef] [PubMed]
77. Farrall, A.J.; Wardlaw, J.M. Blood-brain barrier: Ageing and microvascular disease—systematic review and meta-analysis. *Neurobiol. Aging* **2009**, *30*, 337–352. [CrossRef] [PubMed]
78. Erickson, M.A.; Banks, W.A. Blood-brain barrier dysfunction as a cause and consequence of Alzheimer's disease. *J. Cereb. Blood Flow Metab.* **2013**, *33*, 1500–1513. [CrossRef] [PubMed]
79. Nugent, S.; Castellano, C.A.; Goffaux, P.; Whittingstall, K.; Lepage, M.; Paquet, N.; Bocti, C.; Fulop, T.; Cunnane, S.C. Glucose hypometabolism is highly localized, but lower cortical thickness and brain atrophy are widespread in cognitively normal older adults. *Am. J. Physiol. Endocrinol. Metab.* **2014**, *306*, E1315–E1321. [CrossRef] [PubMed]
80. Montagne, A.; Barnes, S.R.; Sweeney, M.D.; Halliday, M.R.; Sagare, A.P.; Zhao, Z.; Toga, A.W.; Jacobs, R.E.; Liu, C.Y.; Amezcua, L.; et al. Blood-brain barrier breakdown in the aging human hippocampus. *Neuron* **2015**, *85*, 296–302. [CrossRef] [PubMed]
81. Wang, H.; Golob, E.J.; Su, M.Y. Vascular volume and blood-brain barrier permeability measured by dynamic contrast enhanced MRI in hippocampus and cerebellum of patients with MCI and normal controls. *J. Magn. Reson. Imaging* **2006**, *24*, 695–700. [CrossRef] [PubMed]
82. Taheri, S.; Gasparovic, C.; Shah, N.J.; Rosenberg, G.A. Quantitative measurement of blood-brain barrier permeability in human using dynamic contrast-enhanced MRI with fast T1 mapping. *Magn. Reson. Med.* **2011**, *65*, 1036–1042. [CrossRef] [PubMed]

83. Sagare, A.P.; Bell, R.D.; Zlokovic, B.V. Neurovascular dysfunction and faulty amyloid beta-peptide clearance in Alzheimer disease. *Cold Spring Harb. Perspect. Med.* **2012**, *2*, a011452. [CrossRef] [PubMed]
84. Mosconi, L.; de Santi, S.; Li, J.; Tsui, W.H.; Li, Y.; Boppana, M.; Laska, E.; Rusinek, H.; de Leon, M.J. Hippocampal hypometabolism predicts cognitive decline from normal aging. *Neurobiol. Aging* **2008**, *29*, 676–692. [CrossRef] [PubMed]
85. Zhao, W.Q.; Townsend, M. Insulin resistance and amyloidogenesis as common molecular foundation for type 2 diabetes and Alzheimer's disease. *Biochim. Biophys. Acta* **2009**, *1792*, 482–496. [CrossRef] [PubMed]
86. Hoyer, S.; Nitsch, R. Cerebral excess release of neurotransmitter amino acids subsequent to reduced cerebral glucose metabolism in early-onset dementia of Alzheimer type. *J. Neural. Transm.* **1989**, *75*, 227–232. [CrossRef] [PubMed]
87. Reiman, E.M.; Chen, K.; Alexander, G.E.; Caselli, R.J.; Bandy, D.; Osborne, D.; Saunders, A.M.; Hardy, J. Functional brain abnormalities in young adults at genetic risk for late-onset Alzheimer's dementia. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 284–289. [CrossRef] [PubMed]
88. Hooijmans, C.R.; Graven, C.; Dederen, P.J.; Tanila, H.; van Groen, T.; Kiliaan, A.J. Amyloid beta deposition is related to decreased glucose transporter-1 levels and hippocampal atrophy in brains of aged APP/PS1 mice. *Brain Res.* **2007**, *1181*, 93–103. [CrossRef] [PubMed]
89. Liu, Y.; Liu, F.; Iqbal, K.; Grundke-Iqbali, I.; Gong, C.X. Decreased glucose transporters correlate to abnormal hyperphosphorylation of tau in Alzheimer disease. *FEBS Lett.* **2008**, *582*, 359–364. [CrossRef] [PubMed]
90. Li, H.; Guo, Q.; Inoue, T.; Polito, V.A.; Tabuchi, K.; Hammer, R.E.; Pautler, R.G.; Taffet, G.E.; Zheng, H. Vascular and parenchymal amyloid pathology in an Alzheimer disease knock-in mouse model: Interplay with cerebral blood flow. *Mol. Neurodegener.* **2014**, *9*. [CrossRef] [PubMed]
91. Deane, R.; du Yan, S.; Subramanyan, R.K.; LaRue, B.; Jovanovic, S.; Hogg, E.; Welch, D.; Manness, L.; Lin, C.; Yu, J.; et al. RAGE mediates amyloid-beta peptide transport across the blood-brain barrier and accumulation in brain. *Nat. Med.* **2003**, *9*, 907–913. [CrossRef] [PubMed]
92. Baker, L.D.; Cross, D.J.; Minoshima, S.; Belongia, D.; Watson, G.S.; Craft, S. Insulin resistance and Alzheimer-like reductions in regional cerebral glucose metabolism for cognitively normal adults with prediabetes or early type 2 diabetes. *Arch. Neurol.* **2011**, *68*, 51–57. [CrossRef] [PubMed]
93. Ouwendijk, D.M.; van Duinkerken, E.; Schoonenboom, S.N.; Herzfeld de Wiza, D.; Klein, M.; van Golen, L.; Pouwels, P.J.; Barkhof, F.; Moll, A.C.; Snoek, F.J.; et al. Cerebrospinal fluid levels of Alzheimer's disease biomarkers in middle-aged patients with type 1 diabetes. *Diabetologia* **2014**, *57*, 2208–2214. [CrossRef] [PubMed]
94. Hong, H.; Liu, L.P.; Liao, J.M.; Wang, T.S.; Ye, F.Y.; Wu, J.; Wang, Y.Y.; Wang, Y.; Li, Y.Q.; Long, Y.; Xia, Y.Z. Downregulation of LRP1 [correction of LPR1] at the blood-brain barrier in streptozotocin-induced diabetic mice. *Neuropharmacology* **2009**, *56*, 1054–1059. [CrossRef] [PubMed]
95. Sun, Y.N.; Liu, L.B.; Xue, Y.X.; Wang, P. Effects of insulin combined with idebenone on blood-brain barrier permeability in diabetic rats. *J. Neurosci. Res.* **2015**, *93*, 666–677. [CrossRef] [PubMed]
96. Jonas, E.A.; Knox, R.J.; Smith, T.C.; Wayne, N.L.; Connor, J.A.; Kaczmarek, L.K. Regulation by insulin of a unique neuronal Ca<sup>2+</sup> pool and of neuropeptide secretion. *Nature* **1997**, *385*, 343–346. [CrossRef] [PubMed]
97. Wan, Q.; Xiong, Z.G.; Man, H.Y.; Ackerley, C.A.; Brauton, J.; Lu, W.Y.; Becker, L.E.; MacDonald, J.F.; Wang, Y.T. Recruitment of functional GABA(A) receptors to postsynaptic domains by insulin. *Nature* **1997**, *388*, 686–690. [PubMed]
98. Wang, Y.T.; Linden, D.J. Expression of cerebellar long-term depression requires postsynaptic clathrin-mediated endocytosis. *Neuron* **2000**, *25*, 635–647. [CrossRef]
99. Skeberdis, V.A.; Lan, J.; Zheng, X.; Zukin, R.S.; Bennett, M.V. Insulin promotes rapid delivery of N-methyl-D-aspartate receptors to the cell surface by exocytosis. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 3561–3566. [CrossRef] [PubMed]
100. Ahmadian, G.; Ju, W.; Liu, L.; Wyszynski, M.; Lee, S.H.; Dunah, A.W.; Taghibiglou, C.; Wang, Y.; Lu, J.; Wong, T.P.; et al. Tyrosine phosphorylation of GluR2 is required for insulin-stimulated AMPA receptor endocytosis and LTD. *EMBO J.* **2004**, *23*, 1040–1050. [CrossRef] [PubMed]
101. Zhao, W.Q.; Alkon, D.L. Role of insulin and insulin receptor in learning and memory. *Mol. Cell Endocrinol.* **2001**, *177*, 125–134. [CrossRef]
102. Kaidanovich, O.; Eldar-Finkelman, H. The role of glycogen synthase kinase-3 in insulin resistance and type 2 diabetes. *Expert Opin. Ther. Targets* **2002**, *6*, 555–561. [PubMed]

103. Lee, J.; Kim, M.S. The role of GSK3 in glucose homeostasis and the development of insulin resistance. *Diabetes Res. Clin. Pract.* **2007**, *77*, S49–S57. [CrossRef] [PubMed]
104. Balaraman, Y.; Limaye, A.R.; Levey, A.I.; Srinivasan, S. Glycogen synthase kinase 3beta and Alzheimer's disease: Pathophysiological and therapeutic significance. *Cell Mol. Life Sci.* **2006**, *63*, 1226–1235. [CrossRef] [PubMed]
105. Shiiki, T.; Ohtsuki, S.; Kurihara, A.; Naganuma, H.; Nishimura, K.; Tachikawa, M.; Hosoya, K.; Terasaki, T. Brain insulin impairs amyloid-beta(1–40) clearance from the brain. *J. Neurosci.* **2004**, *24*, 9632–9637. [CrossRef] [PubMed]
106. Authier, F.; Posner, B.I.; Bergeron, J.J. Insulin-degrading enzyme. *Clin. Investig. Med.* **1996**, *19*, 149–160.
107. Vekrellis, K.; Ye, Z.; Qiu, W.Q.; Walsh, D.; Hartley, D.; Chesneau, V.; Rosner, M.R.; Selkoe, D.J. Neurons regulate extracellular levels of amyloid beta-protein via proteolysis by insulin-degrading enzyme. *J. Neurosci.* **2000**, *20*, 1657–1665. [PubMed]
108. Farris, W.; Mansourian, S.; Chang, Y.; Lindsley, L.; Eckman, E.A.; Frosch, M.P.; Eckman, C.B.; Tanzi, R.E.; Selkoe, D.J.; Guenette, S. Insulin-degrading enzyme regulates the levels of insulin, amyloid beta-protein, and the beta-amyloid precursor protein intracellular domain *in vivo*. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 4162–4167. [CrossRef] [PubMed]
109. Zhao, L.; Teter, B.; Morihara, T.; Lim, G.P.; Ambegaokar, S.S.; Ubeda, O.J.; Frautschy, S.A.; Cole, G.M. Insulin-degrading enzyme as a downstream target of insulin receptor signaling cascade: Implications for Alzheimer's disease intervention. *J. Neurosci.* **2004**, *24*, 11120–11126. [CrossRef] [PubMed]
110. Zhao, W.Q.; Lacor, P.N.; Chen, H.; Lambert, M.P.; Quon, M.J.; Krafft, G.A.; Klein, W.L. Insulin receptor dysfunction impairs cellular clearance of neurotoxic oligomeric AB. *J. Biol. Chem.* **2009**, *284*, 18742–18753. [CrossRef] [PubMed]
111. De Felice, F.G.; Vieira, M.N.; Bomfim, T.R.; Decker, H.; Velasco, P.T.; Lambert, M.P.; Viola, K.L.; Zhao, W.Q.; Ferreira, S.T.; Klein, W.L. Protection of synapses against Alzheimer's-linked toxins: Insulin signaling prevents the pathogenic binding of Abeta oligomers. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 1971–1976. [CrossRef] [PubMed]
112. De Felice, F.G.; Lourenco, M.V.; Ferreira, S.T. How does brain insulin resistance develop in Alzheimer's disease? *Alzheimers Dement.* **2014**, *10*, S26–S32. [CrossRef] [PubMed]
113. De Felice, F.G.; Ferreira, S.T. Inflammation, defective insulin signaling, and mitochondrial dysfunction as common molecular denominators connecting type 2 diabetes to Alzheimer disease. *Diabetes* **2014**, *63*, 2262–2272. [CrossRef] [PubMed]
114. De Felice, F.G. Alzheimer's disease and insulin resistance: Translating basic science into clinical applications. *J. Clin. Invest.* **2013**, *123*, 531–539. [CrossRef] [PubMed]
115. Ferreira, S.T.; Clarke, J.R.; Bomfim, T.R.; de Felice, F.G. Inflammation, defective insulin signaling, and neuronal dysfunction in Alzheimer's disease. *Alzheimers Dement.* **2014**, *10*, S76–S83. [CrossRef] [PubMed]
116. Hooper, C.; Killick, R.; Lovestone, S. The GSK3 hypothesis of Alzheimer's disease. *J. Neurochem.* **2008**, *104*, 1433–1439. [CrossRef] [PubMed]
117. Reger, M.A.; Watson, G.S.; Green, P.S.; Baker, L.D.; Cholerton, B.; Fishel, M.A.; Plymate, S.R.; Cherrier, M.M.; Schellenberg, G.D.; Frey, W.H., II; et al. Intranasal insulin administration dose-dependently modulates verbal memory and plasma amyloid-beta in memory-impaired older adults. *J. Alzheimers Dis.* **2008**, *13*, 323–331. [PubMed]
118. Craft, S.; Peskind, E.; Schwartz, M.W.; Schellenberg, G.D.; Raskind, M.; Porte, D., Jr. Cerebrospinal fluid and plasma insulin levels in Alzheimer's disease: Relationship to severity of dementia and apolipoprotein E genotype. *Neurology* **1999**, *50*, 164–168. [CrossRef]
119. Steen, E.; Terry, B.M.; Rivera, E.J.; Cannon, J.L.; Neely, T.R.; Tavares, R.; Xu, X.J.; Wands, J.R.; de la Monte, S.M. Impaired insulin and insulin-like growth factor expression and signaling mechanisms in Alzheimer's disease—is this type 3 diabetes? *J. Alzheimers Dis.* **2005**, *7*, 63–80. [PubMed]
120. Rivera, E.J.; Goldin, A.; Fulmer, N.; Tavares, R.; Wands, J.R.; de la Monte, S.M. Insulin and insulin-like growth factor expression and function deteriorate with progression of Alzheimer's disease: Link to brain reductions in acetylcholine. *J. Alzheimers Dis.* **2005**, *8*, 247–268. [PubMed]
121. Craft, S.; Newcomer, J.; Kanne, S.; Dagogo-Jack, S.; Cryer, P.; Sheline, Y.; Luby, J.; Dagogo-Jack, A.; Alderson, A. Memory improvement following induced hyperinsulinemia in Alzheimer's disease. *Neurobiol. Aging* **1996**, *17*, 123–130. [CrossRef]

122. Reger, M.A.; Watson, G.S.; Green, P.S.; Wilkinson, C.W.; Baker, L.D.; Cholerton, B.; Fishel, M.A.; Plymate, S.R.; Breitner, J.C.; DeGroodt, W.; *et al.* Intranasal insulin improves cognition and modulates beta-amyloid in early AD. *Neurology* **2008**, *70*, 440–448. [CrossRef] [PubMed]
123. Schubert, M.; Brazil, D.P.; Burks, D.J.; Kushner, J.A.; Ye, J.; Flint, C.L.; Farhang-Fallah, J.; Dikkes, P.; Warot, X.M.; Rio, C.; *et al.* Insulin receptor substrate-2 deficiency impairs brain growth and promotes tau phosphorylation. *J. Neurosci.* **2003**, *23*, 7084–7092. [PubMed]
124. Schubert, M.; Gautam, D.; Surjo, D.; Ueki, K.; Baudler, S.; Schubert, D.; Kondo, T.; Alber, J.; Galldiks, N.; Kustermann, E.; *et al.* Role for neuronal insulin resistance in neurodegenerative diseases. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 3100–3105. [CrossRef] [PubMed]
125. Grunblatt, E.; Salkovic-Petrisic, M.; Osmanovic, J.; Riederer, P.; Hoyer, S. Brain insulin system dysfunction in streptozotocin intracerebroventricularly treated rats generates hyperphosphorylated tau protein. *J. Neurochem.* **2007**, *101*, 757–770. [CrossRef] [PubMed]
126. Lucas, J.J.; Hernandez, F.; Gomez-Ramos, P.; Moran, M.A.; Hen, R.; Avila, J. Decreased nuclear beta-catenin, tau hyperphosphorylation and neurodegeneration in GSK-3beta conditional transgenic mice. *EMBOJ* **2001**, *20*, 27–39. [CrossRef] [PubMed]
127. Phiel, C.J.; Wilson, C.A.; Lee, V.M.; Klein, P.S. GSK-3alpha regulates production of Alzheimer's disease amyloid-beta peptides. *Nature* **2003**, *423*, 435–439. [CrossRef] [PubMed]
128. Noble, W.; Planell, E.; Zehr, C.; Olm, V.; Meyerson, J.; Suleman, F.; Gaynor, K.; Wang, L.; LaFrancois, J.; Feinstein, B.; *et al.* Inhibition of glycogen synthase kinase-3 by lithium correlates with reduced tauopathy and degeneration *in vivo*. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 6990–6995. [CrossRef] [PubMed]
129. Qing, H.; He, G.; Ly, P.T.; Fox, C.J.; Staufenbiel, M.; Cai, F.; Zhang, Z.; Wei, S.; Sun, X.; Chen, C.H.; *et al.* Valproic acid inhibits Abeta production, neuritic plaque formation, and behavioral deficits in Alzheimer's disease mouse models. *J. Exp. Med.* **2008**, *205*, 2781–2789. [CrossRef] [PubMed]
130. Ly, P.T.; Wu, Y.; Zou, H.; Wang, R.; Zhou, W.; Kinoshita, A.; Zhang, M.; Yang, Y.; Cai, F.; Woodgett, J.; *et al.* Inhibition of GSK3beta-mediated BACE1 expression reduces Alzheimer-associated phenotypes. *J. Clin. Investig.* **2013**, *123*, 224–235. [CrossRef] [PubMed]
131. Ling, X.; Martins, R.N.; Racchi, M.; Craft, S.; Helmerhorst, E. Amyloid beta antagonizes insulin promoted secretion of the amyloid beta protein precursor. *J. Alzheimers. Dis.* **2002**, *4*, 369–374. [PubMed]
132. Xie, L.; Helmerhorst, E.; Taddei, K.; Plewright, B.; van Bronswijk, W.; Martins, R. Alzheimer's beta-amyloid peptides compete for insulin binding to the insulin receptor. *J. Neurosci.* **2002**, *22*, (RC221), 1–5. [PubMed]
133. Biessels, G.J.; Kamal, A.; Urban, I.J.; Spruijt, B.M.; Erkelens, D.W.; Gispen, W.H. Water maze learning and hippocampal synaptic plasticity in streptozotocin-diabetic rats: Effects of insulin treatment. *Brain Res.* **1998**, *800*, 125–135. [CrossRef]
134. Li, Z.G.; Zhang, W.; Grunberger, G.; Sima, A.A. Hippocampal neuronal apoptosis in type 1 diabetes. *Brain Res.* **2002**, *946*, 221–231. [CrossRef]
135. Sima, A.A.; Li, Z.G. The effect of C-peptide on cognitive dysfunction and hippocampal apoptosis in type 1 diabetic rats. *Diabetes* **2005**, *54*, 1497–1505. [CrossRef] [PubMed]
136. Lannert, H.; Hoyer, S. Intracerebroventricular administration of streptozotocin causes long-term diminutions in learning and memory abilities and in cerebral energy metabolism in adult rats. *Behav. Neurosci.* **1998**, *112*, 1199–1208. [CrossRef] [PubMed]
137. Kim, B.; Backus, C.; Oh, S.; Hayes, J.M.; Feldman, E.L. Increased tau phosphorylation and cleavage in mouse models of type 1 and type 2 diabetes. *Endocrinology* **2009**, *150*, 5294–5301. [CrossRef] [PubMed]
138. Li, Z.G.; Zhang, W.; Sima, A.A. Alzheimer-like changes in rat models of spontaneous diabetes. *Diabetes* **2007**, *56*, 1817–1824. [CrossRef] [PubMed]
139. Moreira, P.I.; Santos, M.S.; Seica, R.; Oliveira, C.R. Brain mitochondrial dysfunction as a link between Alzheimer's disease and diabetes. *J. Neurol. Sci.* **2007**, *257*, 206–214. [CrossRef] [PubMed]
140. Cardoso, S.; Santos, M.S.; Seica, R.; Moreira, P.I. Cortical and hippocampal mitochondria bioenergetics and oxidative status during hyperglycemia and/or insulin-induced hypoglycemia. *Biochim. Biophys. Acta.* **2010**, *1802*, 942–951. [CrossRef] [PubMed]
141. Casley, C.S.; Canevari, L.; Land, J.M.; Clark, J.B.; Sharpe, M.A. Beta-amyloid inhibits integrated mitochondrial respiration and key enzyme activities. *J. Neurochem.* **2002**, *80*, 91–100. [CrossRef] [PubMed]

142. Crouch, P.J.; Blake, R.; Duce, J.A.; Ciccosto, G.D.; Li, Q.X.; Barnham, K.J.; Curtain, C.C.; Cherny, R.A.; Cappai, R.; Dyrks, T.; et al. Copper-dependent inhibition of human cytochrome c oxidase by a dimeric conformer of amyloid-beta1-42. *J. Neurosci.* **2005**, *5*, 672–679. [CrossRef] [PubMed]
143. Manczak, M.; Anekonda, T.S.; Henson, E.; Park, B.S.; Quinn, J.; Reddy, P.H. Mitochondria are a direct site of A beta accumulation in Alzheimer's disease neurons: Implications for free radical generation and oxidative damage in disease progression. *Hum. Mol. Genet.* **2006**, *15*, 1437–1449. [CrossRef] [PubMed]
144. Caspersen, C.; Wang, N.; Yao, J.; Sosunov, A.; Chen, X.; Lustbader, J.W.; Xu, H.W.; Stern, D.; McKhann, G.; Yan, S.D. Mitochondrial Abeta: A potential focal point for neuronal metabolic dysfunction in Alzheimer's disease. *FASEB J.* **2005**, *19*, 2040–2041. [PubMed]
145. Devi, L.; Prabhu, B.M.; Galati, D.F.; Avadhani, N.G.; Anandatheerthavarada, H.K. Accumulation of amyloid precursor protein in the mitochondrial import channels of human Alzheimer's disease brain is associated with mitochondrial dysfunction. *J. Neurosci.* **2006**, *26*, 9057–9068. [CrossRef] [PubMed]
146. Lustbader, J.W.; Cirilli, M.; Lin, C.; Xu, H.W.; Takuma, K.; Wang, N.; Caspersen, C.; Chen, X.; Pollak, S.; Chaney, M.; et al. ABAD directly links Abeta to mitochondrial toxicity in Alzheimer's disease. *Science* **2004**, *304*, 448–452. [CrossRef] [PubMed]
147. Yao, J.; Du, H.; Yan, S.; Fang, F.; Wang, C.; Lue, L.F.; Guo, L.; Chen, D.; Stern, D.M.; Gunn Moore, F.J.; et al. Inhibition of amyloid-beta (Abeta) peptide-binding alcohol dehydrogenase-Abeta interaction reduces Abeta accumulation and improves mitochondrial function in a mouse model of Alzheimer's disease. *J. Neurosci.* **2011**, *31*, 2313–2320. [CrossRef] [PubMed]
148. Kish, S.J.; Bergeron, C.; Rajput, A.; Dozic, S.; Mastrogiacomo, F.; Chang, L.J.; Wilson, J.M.; DiStefano, L.M.; Nobrega, J.N. Brain cytochrome oxidase in Alzheimer's disease. *J. Neurochem.* **1992**, *59*, 776–779. [CrossRef] [PubMed]
149. Parker, W.D., Jr.; Mahr, N.J.; Filley, C.M.; Parks, J.K.; Hughes, D.; Young, D.A.; Cullum, C.M. Reduced platelet cytochrome c oxidase activity in Alzheimer's disease. *Neurology* **1994**, *44*, 1086–1090. [CrossRef] [PubMed]
150. Gibson, G.E.; Sheu, K.F.; Blass, J.P. Abnormalities of mitochondrial enzymes in Alzheimer disease. *J. Neural. Transm.* **1998**, *105*, 855–870. [CrossRef] [PubMed]
151. Aksenov, M.Y.; Tucker, H.M.; Nair, P.; Aksenova, M.V.; Butterfield, D.A.; Estus, S.; Markesberry, W.R. The expression of several mitochondrial and nuclear genes encoding the subunits of electron transport chain enzyme complexes, cytochrome c oxidase, and NADH dehydrogenase, in different brain regions in Alzheimer's disease. *Neurochem. Res.* **1999**, *24*, 767–774. [CrossRef] [PubMed]
152. Hirai, K.; Aliev, G.; Nunomura, A.; Fujioka, H.; Russell, R.L.; Atwood, C.S.; Johnson, A.B.; Kress, Y.; Vinters, H.V.; Tabaton, M.; et al. Mitochondrial abnormalities in Alzheimer's disease. *J. Neurosci.* **2001**, *21*, 3017–3023. [PubMed]
153. Bosetti, F.; Brizzi, F.; Barogi, S.; Mancuso, M.; Siciliano, G.; Tendi, E.A.; Murri, L.; Rapoport, S.I.; Solaini, G. Cytochrome c oxidase and mitochondrial F1F0-ATPase (ATP synthase) activities in platelets and brain from patients with Alzheimer's disease. *Neurobiol. Aging* **2002**, *23*, 371–376. [CrossRef]
154. Valla, J.; Schneider, L.; Niedzielko, T.; Coon, K.D.; Caselli, R.; Sabbagh, M.N.; Ahern, G.L.; Baxter, L.; Alexander, G.; Walker, G.; et al. Impaired platelet mitochondrial activity in Alzheimer's disease and mild cognitive impairment. *Mitochondrion* **2006**, *6*, 323–330. [CrossRef] [PubMed]
155. Bubber, P.; Haroutunian, V.; Fisch, G.; Blass, J.P.; Gibson, G.E. Mitochondrial abnormalities in Alzheimer brain: Mechanistic implications. *Ann. Neurol.* **2005**, *57*, 695–703. [CrossRef] [PubMed]
156. Huang, H.M.; Ou, H.C.; Xu, H.; Chen, H.L.; Fowler, C.; Gibson, G.E. Inhibition of alpha-ketoglutarate dehydrogenase complex promotes cytochrome c release from mitochondria, caspase-3 activation, and necrotic cell death. *J. Neurosci. Res.* **2003**, *74*, 309–317. [CrossRef] [PubMed]
157. Tretter, L.; Adam-Vizi, V. Inhibition of Krebs cycle enzymes by hydrogen peroxide: A key role of [alpha]-ketoglutarate dehydrogenase in limiting NADH production under oxidative stress. *J. Neurosci.* **2000**, *20*, 8972–8979. [PubMed]
158. Busciglio, J.; Pelsman, A.; Wong, C.; Pigino, G.; Yuan, M.; Mori, H.; Yankner, B.A. Altered metabolism of the amyloid beta precursor protein is associated with mitochondrial dysfunction in Down's syndrome. *Neuron* **2002**, *33*, 677–688. [CrossRef]
159. Swerdlow, R.H.; Parks, J.K.; Cassarino, D.S.; Maguire, D.J.; Maguire, R.S.; Bennett, J.P., Jr.; Davis, R.E.; Parker, W.D., Jr. Cybrids in Alzheimer's disease: A cellular model of the disease? *Neurology* **1997**, *49*, 918–925. [CrossRef] [PubMed]

160. Chou, J.L.; Shenoy, D.V.; Thomas, N.; Choudhary, P.K.; Laferla, F.M.; Goodman, S.R.; Breen, G.A. Early dysregulation of the mitochondrial proteome in a mouse model of Alzheimer's disease. *J. Proteomics* **2011**, *74*, 466–479. [CrossRef] [PubMed]
161. Li, F.; Calingasan, N.Y.; Yu, F.; Mauck, W.M.; Toidze, M.; Almeida, C.G.; Takahashi, R.H.; Carlson, G.A.; Flint Beal, M.; Lin, M.T.; et al. Increased plaque burden in brains of APP mutant MnSOD heterozygous knockout mice. *J. Neurochem.* **2004**, *89*, 1308–1312. [CrossRef] [PubMed]
162. Anandatheerthavarada, H.K.; Biswas, G.; Robin, M.A.; Avadhani, N.G. Mitochondrial targeting and a novel transmembrane arrest of Alzheimer's amyloid precursor protein impairs mitochondrial function in neuronal cells. *J. Cell Biol.* **2003**, *161*, 41–54. [CrossRef] [PubMed]
163. Kelley, D.E.; He, J.; Menshikova, E.V.; Ritov, V.B. Dysfunction of mitochondria in human skeletal muscle in type 2 diabetes. *Diabetes* **2002**, *51*, 2944–2950. [CrossRef] [PubMed]
164. Anello, M.; Lupi, R.; Spampinato, D.; Piro, S.; Masini, M.; Boggi, U.; del Prato, S.; Rabuazzo, A.M.; Purrello, F.; Marchetti, P. Functional and morphological alterations of mitochondria in pancreatic beta cells from type 2 diabetic patients. *Diabetologia* **2005**, *48*, 282–289. [CrossRef] [PubMed]
165. Moreira, P.I.; Santos, M.S.; Moreno, A.M.; Seica, R.; Oliveira, C.R. Increased vulnerability of brain mitochondria in diabetic (Goto-Kakizaki) rats with aging and amyloid-beta exposure. *Diabetes* **2003**, *52*, 1449–1456. [CrossRef] [PubMed]
166. Mastrocola, R.; Restivo, F.; Vercellinatto, I.; Danni, O.; Brignardello, E.; Aragno, M.; Bocuzzi, G. Oxidative and nitrosative stress in brain mitochondria of diabetic rats. *J. Endocrinol.* **2005**, *187*, 37–44. [CrossRef] [PubMed]
167. Moreira, P.I.; Santos, M.S.; Sena, C.; Seica, R.; Oliveira, C.R. Insulin protects against amyloid beta-peptide toxicity in brain mitochondria of diabetic rats. *Neurobiol. Dis.* **2005**, *18*, 628–637. [CrossRef] [PubMed]
168. Edwards, J.L.; Quattrini, A.; Lentz, S.I.; Figueroa-Romero, C.; Cerri, F.; Backus, C.; Hong, Y.; Feldman, E.L. Diabetes regulates mitochondrial biogenesis and fission in mouse neurons. *Diabetologia* **2010**, *53*, 160–169. [CrossRef] [PubMed]
169. Fernyhough, P.; Roy Chowdhury, S.K.; Schmidt, R.E. Mitochondrial stress and the pathogenesis of diabetic neuropathy. *Expert. Rev. Endocrinol. Metab.* **2010**, *5*, 39–49. [CrossRef] [PubMed]
170. Cardoso, S.; Carvalho, C.; Santos, R.; Correia, S.; Santos, M.S.; Seica, R.; Oliveira, C.R.; Moreira, P.I. Impact of STZ-induced hyperglycemia and insulin-induced hypoglycemia in plasma amino acids and cortical synaptosomal neurotransmitters. *Synapse* **2010**, *65*, 457–466. [CrossRef] [PubMed]
171. Tong, Y.; Zhou, W.; Fung, V.; Christensen, M.A.; Qing, H.; Sun, X.; Song, W. Oxidative stress potentiates BACE1 gene expression and Abeta generation. *J. Neural. Transm.* **2005**, *112*, 455–469. [CrossRef] [PubMed]
172. Shen, C.; Chen, Y.; Liu, H.; Zhang, K.; Zhang, T.; Lin, A.; Jing, N. Hydrogen peroxide promotes Abeta production through JNK-dependent activation of gamma-secretase. *J. Biol. Chem.* **2008**, *283*, 17721–17730. [CrossRef] [PubMed]
173. Tamagno, E.; Guglielotto, M.; Aragno, M.; Borghi, R.; Autelli, R.; Giliberto, L.; Muraca, G.; Danni, O.; Zhu, X.; Smith, M.A.; et al. Oxidative stress activates a positive feedback between the gamma- and beta-secretase cleavages of the beta-amyloid precursor protein. *J. Neurochem.* **2008**, *104*, 683–695. [PubMed]
174. Quiroz-Baez, R.; Rojas, E.; Arias, C. Oxidative stress promotes JNK-dependent amyloidogenic processing of normally expressed human APP by differential modification of alpha-, beta- and gamma-secretase expression. *Neurochem. Int.* **2009**, *55*, 662–670. [CrossRef] [PubMed]
175. Jo, D.G.; Arumugam, T.V.; Woo, H.N.; Park, J.S.; Tang, S.C.; Mughal, M.; Hyun, D.H.; Park, J.H.; Choi, Y.H.; Gwon, A.R.; et al. Evidence that gamma-secretase mediates oxidative stress-induced beta-secretase expression in Alzheimer's disease. *Neurobiol. Aging* **2010**, *31*, 917–925. [CrossRef] [PubMed]
176. Oda, A.; Tamaoka, A.; Araki, W. Oxidative stress up-regulates presenilin 1 in lipid rafts in neuronal cells. *J. Neurosci. Res.* **2010**, *88*, 1137–1145. [CrossRef] [PubMed]
177. Reddy, P.H.; Manczak, M.; Mao, P.; Calkins, M.J.; Reddy, A.P.; Shirendeb, U. Amyloid-beta and mitochondria in aging and Alzheimer's disease: Implications for synaptic damage and cognitive decline. *J. Alzheimers Dis.* **2010**, *20*, S499–S512. [PubMed]
178. Sultana, R.; Boyd-Kimball, D.; Poon, H.F.; Cai, J.; Pierce, W.M.; Klein, J.B.; Markesberry, W.R.; Zhou, X.Z.; Lu, K.P.; Butterfield, D.A. Oxidative modification and down-regulation of Pin1 in Alzheimer's disease hippocampus: A redox proteomics analysis. *Neurobiol. Aging* **2006**, *27*, 918–925. [CrossRef] [PubMed]

179. Sayre, L.M.; Zelasko, D.A.; Harris, P.L.; Perry, G.; Salomon, R.G.; Smith, M.A. 4-Hydroxynonenal-derived advanced lipid peroxidation end products are increased in Alzheimer's disease. *J. Neurochem.* **1997**, *68*, 2092–2097. [CrossRef] [PubMed]
180. Smith, M.A.; Richey Harris, P.L.; Sayre, L.M.; Beckman, J.S.; Perry, G. Widespread peroxynitrite-mediated damage in Alzheimer's disease. *J. Neurosci.* **1997**, *17*, 2653–2657. [PubMed]
181. Gabbita, S.P.; Lovell, M.A.; Markesberry, W.R. Increased nuclear DNA oxidation in the brain in Alzheimer's disease. *J. Neurochem.* **1998**, *71*, 2034–2040. [CrossRef] [PubMed]
182. Montine, T.J.; Markesberry, W.R.; Morrow, J.D.; Roberts, L.J., II. Cerebrospinal fluid F2-isoprostane levels are increased in Alzheimer's disease. *Ann. Neurol.* **1998**, *44*, 410–413. [CrossRef] [PubMed]
183. Nourooz-Zadeh, J.; Liu, E.H.; Yhlen, B.; Anggard, E.E.; Halliwell, B. F4-isoprostanes as specific marker of docosahexaenoic acid peroxidation in Alzheimer's disease. *J. Neurochem.* **1999**, *72*, 734–740. [CrossRef] [PubMed]
184. Nunomura, A.; Perry, G.; Pappolla, M.A.; Wade, R.; Hirai, K.; Chiba, S.; Smith, M.A. RNA oxidation is a prominent feature of vulnerable neurons in Alzheimer's disease. *J. Neurosci.* **1999**, *19*, 1959–1964. [PubMed]
185. Lovell, M.A.; Markesberry, W.R. Ratio of 8-hydroxyguanine in intact DNA to free 8-hydroxyguanine is increased in Alzheimer disease ventricular cerebrospinal fluid. *Arch. Neurol.* **2001**, *58*, 392–396. [CrossRef] [PubMed]
186. Nunomura, A.; Chiba, S.; Lippa, C.F.; Cras, P.; Kalaria, R.N.; Takeda, A.; Honda, K.; Smith, M.A.; Perry, G. Neuronal RNA oxidation is a prominent feature of familial Alzheimer's disease. *Neurobiol. Dis.* **2004**, *17*, 108–113. [CrossRef] [PubMed]
187. Nunomura, A.; Perry, G.; Aliev, G.; Hirai, K.; Takeda, A.; Balraj, E.K.; Jones, P.K.; Ghanbari, H.; Wataya, T.; Shimohama, S.; et al. Oxidative damage is the earliest event in Alzheimer disease. *J. Neuropathol. Exp. Neurol.* **2001**, *60*, 759–767. [CrossRef] [PubMed]
188. Pratico, D.; Uryu, K.; Leight, S.; Trojanowski, J.Q.; Lee, V.M. Increased lipid peroxidation precedes amyloid plaque formation in an animal model of Alzheimer amyloidosis. *J. Neurosci.* **2001**, *21*, 4183–4187. [PubMed]
189. Resende, R.; Moreira, P.I.; Proenca, T.; Deshpande, A.; Busciglio, J.; Pereira, C.; Oliveira, C.R. Brain oxidative stress in a triple-transgenic mouse model of Alzheimer disease. *Free Radic. Biol. Med.* **2008**, *44*, 2051–2057. [CrossRef] [PubMed]
190. Apelt, J.; Bigl, M.; Wunderlich, P.; Schliebs, R. Aging-related increase in oxidative stress correlates with developmental pattern of beta-secretase activity and beta-amyloid plaque formation in transgenic Tg2576 mice with Alzheimer-like pathology. *Int. J. Dev. Neurosci.* **2004**, *22*, 475–484. [CrossRef] [PubMed]
191. Vincent, A.M.; Russell, J.W.; Low, P.; Feldman, E.L. Oxidative stress in the pathogenesis of diabetic neuropathy. *Endocr. Rev.* **2004**, *25*, 612–628. [CrossRef] [PubMed]
192. Rösen, P.; Nawrotzky, P.P.; King, G.; Möller, W.; Tritschler, H.J.; Packer, L. The role of oxidative stress in the onset and progression of diabetes and its complications: A summary of a Congress Series sponsored by UNESCO-MCBN, the American Diabetes Association and the German Diabetes Society. *Diabetes Metab. Res. Rev.* **2001**, *17*, 189–212. [CrossRef] [PubMed]
193. Russell, J.W.; Berent-Spillson, A.; Vincent, A.M.; Freimann, C.L.; Sullivan, K.A.; Feldman, E.L. Oxidative injury and neuropathy in diabetes and impaired glucose tolerance. *Neurobiol. Dis.* **2008**, *30*, 420–429. [CrossRef] [PubMed]
194. Yan, S.D.; Chen, X.; Fu, J.; Chen, M.; Zhu, H.; Roher, A.; Slattery, T.; Zhao, L.; Nagashima, M.; Morser, J.; et al. RAGE and amyloid-beta peptide neurotoxicity in Alzheimer's disease. *Nature* **1996**, *382*, 685–691. [CrossRef] [PubMed]
195. Srikanth, V.; Maczurek, A.; Phan, T.; Steele, M.; Westcott, B.; Juskiw, D.; Munch, G. Advanced glycation endproducts and their receptor RAGE in Alzheimer's disease. *Neurobiol. Aging* **2011**, *32*, 763–777. [CrossRef] [PubMed]
196. Bucala, R.; Cerami, A. Advanced glycosylation: Chemistry, biology, and implications for diabetes and aging. *Adv. Pharmacol.* **1992**, *23*, 1–34. [PubMed]
197. Hunt, J.V.; Dean, R.T.; Wolff, S.P. Hydroxyl radical production and autoxidative glycosylation. Glucose autoxidation as the cause of protein damage in the experimental glycation model of diabetes mellitus and ageing. *Biochem. J.* **1988**, *256*, 205–212. [CrossRef] [PubMed]
198. Mullarkey, C.J.; Edelstein, D.; Brownlee, M. Free radical generation by early glycation products: A mechanism for accelerated atherogenesis in diabetes. *Biochem. Biophys. Res. Commun.* **1990**, *173*, 932–939. [CrossRef]

199. Smith, M.A.; Taneda, S.; Richey, P.L.; Miyata, S.; Yan, S.D.; Stern, D.; Sayre, L.M.; Monnier, V.M.; Perry, G. Advanced Maillard reaction end products are associated with Alzheimer disease pathology. *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 5710–5714. [CrossRef] [PubMed]
200. Vitek, M.P.; Bhattacharya, K.; Glendening, J.M.; Stopa, E.; Vlassara, H.; Bucala, R.; Manogue, K.; Cerami, A. Advanced glycation end products contribute to amyloidosis in Alzheimer disease. *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 4766–4770. [CrossRef] [PubMed]
201. Ledesma, M.D.; Bonay, P.; Colaco, C.; Avila, J. Analysis of microtubule-associated protein tau glycation in paired helical filaments. *J. Biol. Chem.* **1994**, *269*, 21614–21619. [PubMed]
202. Yan, S.D.; Yan, S.F.; Chen, X.; Fu, J.; Chen, M.; Kuppusamy, P.; Smith, M.A.; Perry, G.; Godman, C.G.; Nawroth, P.; et al. Non-enzymatically glycated tau in Alzheimer's disease induces neuronal oxidant stress resulting in cytokine gene expression and release of amyloid beta-peptide. *Nat. Med.* **1995**, *1*, 693–699. [CrossRef] [PubMed]
203. McCance, D.R.; Dyer, D.G.; Dunn, J.A.; Bailie, K.E.; Thorpe, S.R.; Baynes, J.W.; Lyons, T.J. Maillard reaction products and their relation to complications in insulin-dependent diabetes mellitus. *J. Clin. Investig.* **1993**, *91*, 2470–2478. [CrossRef] [PubMed]
204. Vlassara, H. Advanced glycation end-products and atherosclerosis. *Ann. Med.* **1996**, *28*, 419–426. [CrossRef] [PubMed]
205. Wells-Knecht, K.J.; Brinkmann, E.; Wells-Knecht, M.C.; Litchfield, J.E.; Ahmed, M.U.; Reddy, S.; Zyzak, D.V.; Thorpe, S.R.; Baynes, J.W. New biomarkers of Maillard reaction damage to proteins. *Nephrol. Dial. Transplant.* **1996**, *11*, 41–47. [CrossRef] [PubMed]
206. Hammes, H.P.; Alt, A.; Niwa, T.; Clausen, J.T.; Bretzel, R.G.; Brownlee, M.; Schleicher, E.D. Differential accumulation of advanced glycation end products in the course of diabetic retinopathy. *Diabetologia* **1999**, *42*, 728–736. [CrossRef] [PubMed]
207. Goh, S.Y.; Cooper, M.E. Clinical review: The role of advanced glycation end products in progression and complications of diabetes. *J. Clin. Endocrinol. Metab.* **2008**, *93*, 1143–1152. [CrossRef] [PubMed]
208. Smith, M.A.; Sayre, L.M.; Perry, G. Diabetes mellitus and Alzheimer's disease: Glycation as a biochemical link. *Diabetologia* **1996**, *39*, 247. [CrossRef] [PubMed]
209. Smith, M.A.; Tabaton, M.; Perry, G. Early contribution of oxidative glycation in Alzheimer disease. *Neurosci. Lett.* **1996**, *217*, 210–211. [CrossRef]
210. Wahrle, S.; Das, P.; Nyborg, A.C.; McLendon, C.; Shoji, M.; Kawarabayashi, T.; Younkin, L.H.; Younkin, S.G.; Golde, T.E. Cholesterol-dependent gamma-secretase activity in buoyant cholesterol-rich membrane microdomains. *Neurobiol. Dis.* **2002**, *9*, 11–23. [CrossRef] [PubMed]
211. Kalvodova, L.; Kahya, N.; Schwille, P.; Ehehalt, R.; Verkade, P.; Drechsel, D.; Simons, K. Lipids as modulators of proteolytic activity of BACE: Involvement of cholesterol, glycosphingolipids, and anionic phospholipids *in vitro*. *J. Biol. Chem.* **2005**, *280*, 36815–36823. [CrossRef] [PubMed]
212. Osenkowski, P.; Ye, W.; Wang, R.; Wolfe, M.S.; Selkoe, D.J. Direct and potent regulation of gamma-secretase by its lipid microenvironment. *J. Biol. Chem.* **2008**, *283*, 22529–22540. [CrossRef] [PubMed]
213. Canevari, L.; Clark, J.B. Alzheimer's disease and cholesterol: The fat connection. *Neurochem. Res.* **2007**, *32*, 739–750. [CrossRef] [PubMed]
214. Brecht, W.J.; Harris, F.M.; Chang, S.; Tesseur, I.; Yu, G.Q.; Xu, Q.; Dee Fish, J.; Wyss-Coray, T.; Buttini, M.; Mucke, L.; et al. Neuron-specific apolipoprotein e4 proteolysis is associated with increased tau phosphorylation in brains of transgenic mice. *J. Neurosci.* **2004**, *24*, 2527–2534. [CrossRef] [PubMed]
215. Mahley, R.W.; Rall, S.C., Jr. Apolipoprotein E: Far more than a lipid transport protein. *Annu. Rev. Genomics Hum. Genet.* **2000**, *1*, 507–537. [CrossRef] [PubMed]
216. Lynch, J.R.; Tang, W.; Wang, H.; Vitek, M.P.; Bennett, E.R.; Sullivan, P.M.; Warner, D.S.; Laskowitz, D.T. APOE genotype and an ApoE-mimetic peptide modify the systemic and central nervous system inflammatory response. *J. Biol. Chem.* **2003**, *278*, 48529–48533. [CrossRef] [PubMed]
217. Van den Elzen, P.; Garg, S.; Leon, L.; Brigl, M.; Leadbetter, E.A.; Gumperz, J.E.; Dascher, C.C.; Cheng, T.Y.; Sacks, F.M.; Illarionov, P.A.; et al. Apolipoprotein-mediated pathways of lipid antigen presentation. *Nature* **2005**, *437*, 906–910. [CrossRef] [PubMed]
218. Roses, A.D. Apolipoprotein E alleles as risk factors in Alzheimer's disease. *Annu. Rev. Med.* **1996**, *47*, 387–400. [CrossRef] [PubMed]

219. Bertram, L.; Tanzi, R.E. Thirty years of Alzheimer's disease genetics: The implications of systematic meta-analyses. *Nat. Rev. Neurosci.* **2008**, *9*, 768–778. [CrossRef] [PubMed]
220. Gibson, G.E.; Haroutunian, V.; Zhang, H.; Park, L.C.; Shi, Q.; Lesser, M.; Mohs, R.C.; Sheu, R.K.; Blass, J.P. Mitochondrial damage in Alzheimer's disease varies with apolipoprotein E genotype. *Ann. Neurol.* **2000**, *48*, 297–303. [CrossRef]
221. Puglielli, L.; Konopka, G.; Pack-Chung, E.; Ingano, L.A.; Berezovska, O.; Hyman, B.T.; Chang, T.Y.; Tanzi, R.E.; Kovacs, D.M. Acyl-coenzyme A: Cholesterol acyltransferase modulates the generation of the amyloid beta-peptide. *Nat. Cell Biol.* **2001**, *3*, 905–912. [CrossRef] [PubMed]
222. Shibuya, Y.; Niu, Z.; Bryleva, E.Y.; Harris, B.T.; Murphy, S.R.; Kheirullah, A.; Bowen, Z.D.; Chang, C.C.; Chang, T.Y. Acyl-coenzyme A: Cholesterol acyltransferase 1 blockage enhances autophagy in the neurons of triple transgenic Alzheimer's disease mouse and reduces human P301L-tau content at the presymptomatic stage. *Neurobiol. Aging* **2015**, *36*, 2248–2259. [CrossRef] [PubMed]
223. Hutter-Paier, B.; Huttunen, H.J.; Puglielli, L.; Eckman, C.B.; Kim, D.Y.; Hofmeister, A.; Moir, R.D.; Domnitz, S.B.; Frosch, M.P.; Windisch, M.; et al. The ACAT inhibitor CP-113,818 markedly reduces amyloid pathology in a mouse model of Alzheimer's disease. *Neuron* **2004**, *44*, 227–238. [CrossRef] [PubMed]
224. Bhattacharyya, R.; Kovacs, D.M. ACAT inhibition and amyloid beta reduction. *Biochim. Biophys. Acta* **2010**, *1801*, 960–965. [CrossRef] [PubMed]
225. Bryleva, E.Y.; Rogers, M.A.; Chang, C.C.; Buen, F.; Harris, B.T.; Rousselet, E.; Seidah, N.G.; Oddo, S.; LaFerla, F.M.; Spencer, T.A.; et al. ACAT1 gene ablation increases 24(S)-hydroxycholesterol content in the brain and ameliorates amyloid pathology in mice with AD. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 3081–3086. [CrossRef] [PubMed]
226. Simons, M.; Keller, P.; de Strooper, B.; Beyreuther, K.; Dotti, C.G.; Simons, K. Cholesterol depletion inhibits the generation of beta-amyloid in hippocampal neurons. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 6460–6464. [CrossRef] [PubMed]
227. Bien-Ly, N.; Andrews-Zwilling, Y.; Xu, Q.; Bernardo, A.; Wang, C.; Huang, Y. C-terminal-truncated apolipoprotein (apo) E4 inefficiently clears amyloid-beta (Abeta) and acts in concert with Abeta to elicit neuronal and behavioral deficits in mice. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 4236–4241. [CrossRef] [PubMed]
228. Harris, F.M.; Brecht, W.J.; Xu, Q.; Tesseur, I.; Kekonius, L.; Wyss-Coray, T.; Fish, J.D.; Maslia, E.; Hopkins, P.C.; Scearce-Levie, K.; et al. Carboxyl-terminal-truncated apolipoprotein E4 causes Alzheimer's disease-like neurodegeneration and behavioral deficits in transgenic mice. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 10966–10971. [CrossRef] [PubMed]
229. Castellano, J.M.; Kim, J.; Stewart, F.R.; Jiang, H.; DeMattos, R.B.; Patterson, B.W.; Fagan, A.M.; Morris, J.C.; Mawuenyega, K.G.; Cruchaga, C.; et al. Human apoE isoforms differentially regulate brain amyloid-beta peptide clearance. *Sci. Transl. Med.* **2011**, *3*. [CrossRef] [PubMed]
230. Raber, J.; Wong, D.; Buttini, M.; Orth, M.; Bellobusta, S.; Pitas, R.E.; Mahley, R.W.; Mucke, L. Isoform-specific effects of human apolipoprotein E on brain function revealed in ApoE knockout mice: increased susceptibility of females. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 10914–10919. [CrossRef] [PubMed]
231. Raber, J.; Wong, D.; Yu, G.Q.; Buttini, M.; Mahley, R.W.; Pitas, R.E.; Mucke, L. Apolipoprotein E and cognitive performance. *Nature* **2000**, *404*, 352–354. [CrossRef] [PubMed]
232. Peila, R.; Rodriguez, B.L.; Launer, L.J. Type 2 diabetes, APOE gene, and the risk for dementia and related pathologies: The Honolulu-Asia aging study. *Diabetes* **2002**, *51*, 1256–1262. [CrossRef] [PubMed]
233. Pickup, J.C.; Mattock, M.B.; Chusney, G.D.; Burt, D. NIDDM as a disease of the innate immune system: Association of acute-phase reactants and interleukin-6 with metabolic syndrome X. *Diabetologia* **1997**, *40*, 1286–1292. [CrossRef] [PubMed]
234. Spranger, J.; Kroke, A.; Mohlig, M.; Hoffmann, K.; Bergmann, M.M.; Ristow, M.; Boeing, H.; Pfeiffer, A.F. Inflammatory cytokines and the risk to develop type 2 diabetes: Results of the prospective population-based European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam study. *Diabetes* **2003**, *52*, 812–817. [CrossRef] [PubMed]
235. Herder, C.; Illig, T.; Rathmann, W.; Martin, S.; Haastert, B.; Muller-Scholze, S.; Holle, R.; Thorand, B.; Koenig, W.; Wichmann, H.E.; et al. Inflammation and type 2 diabetes: Results from KORA Augsburg. *Gesundheitswesen* **2005**, *67*, S115–S121. [CrossRef] [PubMed]

236. Herder, C.; Brunner, E.J.; Rathmann, W.; Strassburger, K.; Tabak, A.G.; Schloot, N.C.; Witte, D.R. Elevated levels of the anti-inflammatory interleukin-1 receptor antagonist precede the onset of type 2 diabetes: The Whitehall II study. *Diabetes Care* **2009**, *32*, 421–423. [CrossRef] [PubMed]
237. Maedler, K.; Sergeev, P.; Ris, F.; Oberholzer, J.; Joller-Jemelka, H.I.; Spinas, G.A.; Kaiser, N.; Halban, P.A.; Donath, M.Y. Glucose-induced beta cell production of IL-1beta contributes to glucotoxicity in human pancreatic islets. *J. Clin. Investig.* **2002**, *110*, 851–860. [CrossRef] [PubMed]
238. Maedler, K.; Sergeev, P.; Ehses, J.A.; Mathe, Z.; Bosco, D.; Berney, T.; Dayer, J.M.; Reinecke, M.; Halban, P.A.; Donath, M.Y. Leptin modulates beta cell expression of IL-1 receptor antagonist and release of IL-1beta in human islets. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 8138–8143. [CrossRef] [PubMed]
239. Ehses, J.A.; Perren, A.; Eppler, E.; Ribaux, P.; Pospisilik, J.A.; Maor-Cahn, R.; Gueripel, X.; Ellingsgaard, X.; Schneider, M.K.; Biollaz, G.; et al. Increased number of islet-associated macrophages in type 2 diabetes. *Diabetes* **2007**, *56*, 2356–2370. [CrossRef] [PubMed]
240. Akiyama, H.; Barger, S.; Barnum, S.; Bradt, B.; Bauer, J.; Cole, G.M.; Cooper, N.R.; Eikelenboom, P.; Emmerling, M.; Fiebich, B.L.; et al. Inflammation and Alzheimer's disease. *Neurobiol. Aging* **2000**, *21*, 383–421. [CrossRef]
241. Lee, S.; Tong, M.; Hang, S.; Deochand, C.; de la Monte, S.M. CSF and brain indices of insulin resistance, oxidative stress and neuro-inflammation in early versus late Alzheimer's disease. *J. Alzheimers Dis. Parkinsonism* **2013**, *3*. [CrossRef]
242. Blalock, E.M.; Chen, K.C.; Stromberg, A.J.; Norris, C.M.; Kadish, I.; Kraner, S.D.; Porter, N.M.; Landfield, P.W. Harnessing the power of gene microarrays for the study of brain aging and Alzheimer's disease: Statistical reliability and functional correlation. *Ageing Res. Rev.* **2005**, *4*, 481–512. [CrossRef] [PubMed]
243. Katsel, P.L.; Davis, K.L.; Haroutunian, V. Large-scale microarray studies of gene expression in multiple regions of the brain in schizophrenia and Alzheimer's disease. *Int. Rev. Neurobiol.* **2005**, *63*, 41–82. [PubMed]
244. Morgan, D.; Gordon, M.N.; Tan, J.; Wilcock, D.; Rojiani, A.M. Dynamic complexity of the microglial activation response in transgenic models of amyloid deposition: Implications for Alzheimer therapeutics. *J. Neuropathol. Exp. Neurol.* **2005**, *64*, 743–753. [CrossRef] [PubMed]
245. Heneka, M.T.; Sastre, M.; Dumitrescu-Ozimek, L.; Dewachter, I.; Walter, J.; Klockgether, T.; Van Leuven, F. Focal glial activation coincides with increased BACE1 activation and precedes amyloid plaque deposition in APP[V717I] transgenic mice. *J. Neuroinflamm.* **2005**, *2*. [CrossRef] [PubMed]
246. Kitazawa, M.; Oddo, S.; Yamasaki, T.R.; Green, K.N.; LaFerla, F.M. Lipopolysaccharide-induced inflammation exacerbates tau pathology by a cyclin-dependent kinase 5-mediated pathway in a transgenic model of Alzheimer's disease. *J. Neurosci.* **2005**, *25*, 8843–8853. [CrossRef] [PubMed]
247. Szekely, C.A.; Thorne, J.E.; Zandi, P.P.; Ek, M.; Messias, E.; Breitner, J.C.; Goodman, S.N. Nonsteroidal anti-inflammatory drugs for the prevention of Alzheimer's disease: A systematic review. *Neuroepidemiology* **2004**, *23*, 159–169. [CrossRef] [PubMed]
248. Aisen, P.S.; Davis, K.L.; Berg, J.D.; Schafer, K.; Campbell, K.; Thomas, R.G.; Weiner, M.F.; Farlow, M.R.; Sano, M.; Grundman, M.; et al. A randomized controlled trial of prednisone in Alzheimer's disease. Alzheimer's Disease Cooperative Study. *Neurology* **2000**, *54*, 588–593. [CrossRef] [PubMed]
249. Aisen, P.S.; Schafer, K.A.; Grundman, M.; Pfeiffer, E.; Sano, M.; Davis, K.L.; Farlow, M.R.; Jin, S.; Thomas, R.G.; Thal, L.J. Effects of rofecoxib or naproxen vs. placebo on Alzheimer disease progression: A randomized controlled trial. *JAMA* **2003**, *289*, 2819–2826. [CrossRef] [PubMed]
250. Van Gool, W.A.; Weinstein, H.C.; Scheltens, P.; Walstra, G.J. Effect of hydroxychloroquine on progression of dementia in early Alzheimer's disease: An 18-month randomised, double-blind, placebo-controlled study. *Lancet* **2001**, *358*, 455–460. [CrossRef]
251. Reines, S.A.; Block, G.A.; Morris, J.C.; Liu, G.; Nessly, M.L.; Lines, C.R.; Norman, B.A.; Baranak, C.C. Rofecoxib: No effect on Alzheimer's disease in a 1-year, randomized, blinded, controlled study. *Neurology* **2004**, *62*, 66–71. [CrossRef] [PubMed]
252. Zhao, W.Q.; De Felice, F.G.; Fernandez, S.; Chen, H.; Lambert, M.P.; Quon, M.J.; Krafft, G.A.; Klein, W.L. Amyloid beta oligomers induce impairment of neuronal insulin receptors. *FASEB J.* **2008**, *22*, 246–260. [CrossRef] [PubMed]
253. Zhao, W.Q.; Chen, H.; Quon, M.J.; Alkon, D.L. Insulin and the insulin receptor in experimental models of learning and memory. *Eur. J. Pharmacol.* **2004**, *490*, 71–81. [CrossRef] [PubMed]

254. Haj-ali, V.; Mohaddes, G.; Babri, S.H. Intracerebroventricular insulin improves spatial learning and memory in male Wistar rats. *Behav. Neurosci.* **2009**, *123*, 1309–1314. [CrossRef] [PubMed]
255. Francis, G.J.; Martinez, J.A.; Liu, W.Q.; Xu, K.; Ayer, A.; Fine, J.; Tour, U.I.; Glazner, G.; Hanson, L.R.; Frey, W.H., II; et al. Intranasal insulin prevents cognitive decline, cerebral atrophy and white matter changes in murine type I diabetic encephalopathy. *Brain* **2008**, *131*, 3311–3334. [PubMed]
256. Biessels, G.J.; Deary, I.J.; Ryan, C.M. Cognition and diabetes: A lifespan perspective. *Lancet Neurol.* **2008**, *7*, 184–190. [CrossRef]
257. Brands, A.M.; Biessels, G.J.; de Haan, E.H.; Kappelle, L.J.; Kessels, R.P. The effects of type 1 diabetes on cognitive performance: A meta-analysis. *Diabetes Care* **2005**, *28*, 726–735. [CrossRef] [PubMed]
258. Kodl, C.T.; Seaquist, E.R. Cognitive dysfunction and diabetes mellitus. *Endocr. Rev.* **2008**, *29*, 494–511. [CrossRef] [PubMed]
259. Benedict, C.; Hallschmid, M.; Hatke, A.; Schultes, B.; Fehm, H.L.; Born, J.; Kern, W. Intranasal insulin improves memory in humans. *Psychoneuroendocrinology* **2004**, *29*, 1326–1334. [CrossRef] [PubMed]
260. Benedict, C.; Kern, W.; Schultes, B.; Born, J.; Hallschmid, M. Differential sensitivity of men and women to anorexigenic and memory-improving effects of intranasal insulin. *J. Clin. Endocrinol. Metab.* **2008**, *93*, 1339–1344. [CrossRef] [PubMed]
261. Benedict, C.; Hallschmid, M.; Schmitz, K.; Schultes, B.; Ratter, F.; Fehm, H.L.; Born, J.; Kern, W. Intranasal insulin improves memory in humans: Superiority of insulin aspart. *Neuropsychopharmacology* **2007**, *32*, 239–243. [CrossRef] [PubMed]
262. Krug, R.; Benedict, C.; Born, J.; Hallschmid, M. Comparable sensitivity of postmenopausal and young women to the effects of intranasal insulin on food intake and working memory. *J. Clin. Endocrinol. Metab.* **2010**, *95*, E468–E472. [CrossRef] [PubMed]
263. Reger, M.A.; Watson, G.S.; Frey, W.H., II; Baker, L.D.; Cholerton, B.; Keeling, M.L.; Belongia, D.A.; Fishel, M.A.; Plymate, S.R.; Schellenberg, G.D.; et al. Effects of intranasal insulin on cognition in memory-impaired older adults: Modulation by APOE genotype. *Neurobiol. Aging* **2006**, *27*, 451–458. [CrossRef] [PubMed]
264. DeJong, R.N. CNS manifestations of diabetes mellitus. *Postgrad. Med.* **1977**, *61*, 101–107. [PubMed]
265. Reske-Nielsen, E.; Lundbaek, K. Diabetic encephalopathy. Diffuse and focal lesions of the brain in long-term diabetes. *Acta Neurol. Scand. Suppl.* **1963**, *39*, 273–290. [CrossRef] [PubMed]
266. Allen, K.V.; Frier, B.M.; Strachan, M.W. The relationship between type 2 diabetes and cognitive dysfunction: Longitudinal studies and their methodological limitations. *Eur. J. Pharmacol.* **2004**, *490*, 169–175. [CrossRef] [PubMed]
267. Roriz-Filho, J.S.; Sa-Roriz, T.M.; Rosset, I.; Camozzato, A.L.; Santos, A.C.; Chaves, M.L.; Moriguti, J.C.; Roriz-Cruz, M. (Pre)diabetes, brain aging, and cognition. *Biochim. Biophys. Acta* **2009**, *1792*, 432–443. [CrossRef] [PubMed]
268. Fernandez, A.M.; Torres-Alemán, I. The many faces of insulin-like peptide signalling in the brain. *Nat. Rev. Neurosci.* **2012**, *13*, 225–239. [CrossRef] [PubMed]
269. Zheng, W.H.; Kar, S.; Dore, S.; Quirion, R. Insulin-like growth factor-1 (IGF-1): A neuroprotective trophic factor acting via the Akt kinase pathway. *J. Neural. Transm. Suppl.* **2000**, *60*, 261–272. [PubMed]
270. Dore, S.; Bastianetto, S.; Kar, S.; Quirion, R. Protective and rescuing abilities of IGF-I and some putative free radical scavengers against beta-amyloid-inducing toxicity in neurons. *Ann. N. Y. Acad. Sci.* **1999**, *890*, 356–364. [CrossRef] [PubMed]
271. Dore, S.; Kar, S.; Quirion, R. Insulin-like growth factor I protects and rescues hippocampal neurons against beta-amyloid- and human amylin-induced toxicity. *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 4772–4777. [CrossRef] [PubMed]
272. Evin, G.; Weidemann, A. Biogenesis and metabolism of Alzheimer's disease Abeta amyloid peptides. *Peptides* **2002**, *23*, 1285–1297. [CrossRef]
273. Tsukamoto, E.; Hashimoto, Y.; Kanekura, K.; Niikura, T.; Also, S.; Nishimoto, I. Characterization of the toxic mechanism triggered by Alzheimer's amyloid-beta peptides via p75 neurotrophin receptor in neuronal hybrid cells. *J. Neurosci. Res.* **2003**, *73*, 627–636. [CrossRef] [PubMed]
274. Pandini, G.; Pace, V.; Copani, A.; Squatrito, S.; Milardi, D.; Vigneri, R. Insulin has multiple antiamyloidogenic effects on human neuronal cells. *Endocrinology* **2012**, *154*, 375–387. [CrossRef] [PubMed]

275. Watson, G.S.; Peskind, E.R.; Asthana, S.; Purganan, K.; Wait, C.; Chapman, D.; Schwartz, M.W.; Plymate, S.; Craft, S. Insulin increases CSF Abeta42 levels in normal older adults. *Neurology* **2003**, *60*, 1899–1903. [CrossRef] [PubMed]
276. Benedict, C.; Brooks, S.J.; Kullberg, J.; Burgos, J.; Kempton, M.J.; Nordenskjold, R.; Nylander, R.; Kilander, L.; Craft, S.; Larsson, E.M.; *et al.* Impaired insulin sensitivity as indexed by the HOMA score is associated with deficits in verbal fluency and temporal lobe gray matter volume in the elderly. *Diabetes Care* **2012**, *35*, 488–494. [CrossRef] [PubMed]
277. Talbot, K.; Wang, H.Y.; Kazi, H.; Han, L.Y.; Bakshi, K.P.; Stucky, A.; Fuino, R.L.; Kawaguchi, K.R.; Samoyedny, A.J.; Wilson, R.S.; *et al.* Demonstrated brain insulin resistance in Alzheimer’s disease patients is associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive decline. *J. Clin. Investig.* **2012**, *122*, 1316–1338. [CrossRef] [PubMed]
278. Bomfim, T.R.; Forny-Germano, L.; Sathler, L.B.; Brito-Moreira, J.; Houzel, J.C.; Decker, H.; Silverman, M.A.; Kazi, H.; Melo, H.M.; McClean, P.L.; *et al.* An anti-diabetes agent protects the mouse brain from defective insulin signaling caused by Alzheimer’s disease—Associated Abeta oligomers. *J. Clin. Investig.* **2012**, *22*, 1339–1353. [CrossRef] [PubMed]
279. Kenna, H.; Hoeft, F.; Kelley, R.; Wroolie, T.; Demuth, B.; Reiss, A.; Rasgon, N. Fasting plasma insulin and the default mode network in women at risk for Alzheimer’s disease. *Neurobiol. Aging* **2013**, *34*, 641–649. [CrossRef] [PubMed]
280. Asai, M.; Hattori, C.; Iwata, N.; Saido, T.C.; Sasagawa, N.; Szabo, B.; Hashimoto, Y.; Maruyama, K.; Tanuma, S.; Kiso, Y.; Ishiura, S. The novel beta-secretase inhibitor KMI-429 reduces amyloid beta peptide production in amyloid precursor protein transgenic and wild-type mice. *J. Neurochem.* **2006**, *96*, 533–540. [CrossRef] [PubMed]
281. Comery, T.A.; Martone, R.L.; Aschmies, S.; Atchison, K.P.; Diamantidis, G.; Gong, X.; Zhou, H.; Kreft, A.F.; Pangalos, M.N.; Sonnenberg-Reines, J.; *et al.* Acute gamma-secretase inhibition improves contextual fear conditioning in the Tg2576 mouse model of Alzheimer’s disease. *J. Neurosci.* **2005**, *25*, 8898–8902. [CrossRef] [PubMed]
282. Schenk, D.; Barbour, R.; Dunn, W.; Gordon, G.; Grajeda, H.; Guido, T.; Hu, K.; Huang, J.; Johnson-Wood, K.; Khan, K.; *et al.* Immunization with amyloid-beta attenuates Alzheimer-disease-like pathology in the PDAPP mouse. *Nature* **1999**, *400*, 173–177. [CrossRef] [PubMed]
283. Gravitz, L. Drugs: A tangled web of targets. *Nature* **2011**, *475*, S9–S11. [CrossRef] [PubMed]
284. Holmes, C.; Boche, D.; Wilkinson, D.; Yadegarfar, G.; Hopkins, V.; Bayer, A.; Jones, R.W.; Bullock, R.; Love, S.; Neal, J.W.; *et al.* Long-term effects of Abeta42 immunisation in Alzheimer’s disease: Follow-up of a randomised, placebo-controlled phase I trial. *Lancet* **2008**, *372*, 216–223. [CrossRef]
285. Gilman, S.; Koller, M.; Black, R.S.; Jenkins, L.; Griffith, S.G.; Fox, N.C.; Eisner, L.; Kirby, L.; Rovira, M.B.; Forette, F.; *et al.* Clinical effects of Abeta immunization (AN1792) in patients with AD in an interrupted trial. *Neurology* **2005**, *64*, 1553–1562. [CrossRef] [PubMed]
286. Orgogozo, J.M.; Gilman, S.; Dartigues, J.F.; Laurent, B.; Puel, M.; Kirby, L.C.; Jouanny, P.; Dubois, B.; Eisner, L.; Flitman, S.; *et al.* Subacute meningoencephalitis in a subset of patients with AD after Abeta42 immunization. *Neurology* **2003**, *61*, 46–54. [CrossRef] [PubMed]
287. Zandi, P.P.; Anthony, J.C.; Khachaturian, A.S.; Stone, S.V.; Gustafson, D.; Tschanz, J.T.; Norton, M.C.; Welsh-Bohmer, K.A.; Breitner, J.C. Reduced risk of Alzheimer disease in users of antioxidant vitamin supplements: The Cache County Study. *Arch. Neurol.* **2004**, *61*, 82–88. [CrossRef] [PubMed]
288. Aisen, P.S.; Saumier, D.; Briand, R.; Laurin, J.; Gervais, F.; Tremblay, P.; Garceau, D. A Phase II study targeting amyloid-beta with 3APS in mild-to-moderate Alzheimer disease. *Neurology* **2006**, *67*, 1757–1763. [CrossRef] [PubMed]
289. Scharf, S.; Mander, A.; Ugoni, A.; Vajda, F.; Christophidis, N. A double-blind, placebo-controlled trial of diclofenac/misoprostol in Alzheimer’s disease. *Neurology* **1999**, *53*, 197–201. [CrossRef] [PubMed]
290. Porsteinsson, A.P.; Grossberg, G.T.; Mintzer, J.; Olin, J.T. Memantine treatment in patients with mild to moderate Alzheimer’s disease already receiving a cholinesterase inhibitor: A randomized, double-blind, placebo-controlled trial. *Curr. Alzheimer Res.* **2008**, *5*, 83–89. [CrossRef] [PubMed]
291. Yarchoan, M.; Arnold, S.E. Repurposing diabetes drugs for brain insulin resistance in Alzheimer disease. *Diabetes* **2014**, *63*, 2253–2261. [CrossRef] [PubMed]
292. American Diabetes Association. Standards of medical care in diabetes—2013. *Diabetes Care* **2013**, *36*, S11–S66.

293. Nicolakakis, N.; Aboulkassim, T.; Ongali, B.; Lecrux, C.; Fernandes, P.; Rosa-Neto, P.; Tong, X.K.; Hamel, E. Complete rescue of cerebrovascular function in aged Alzheimer's disease transgenic mice by antioxidants and pioglitazone, a peroxisome proliferator-activated receptor gamma agonist. *J. Neurosci.* **2008**, *28*, 9287–9296. [CrossRef] [PubMed]
294. Watson, G.S.; Cholerton, B.A.; Reger, M.A.; Baker, L.D.; Plymate, S.R.; Asthana, S.; Fishel, M.A.; Kulstad, J.J.; Green, P.S.; Cook, D.G.; et al. Preserved cognition in patients with early Alzheimer disease and amnestic mild cognitive impairment during treatment with rosiglitazone: A preliminary study. *Am. J. Geriatr. Psychiatry* **2005**, *13*, 950–958. [CrossRef] [PubMed]
295. Harrington, C.; Sawchak, S.; Chiang, C.; Davies, J.; Saunders, A.; Irizarry, M.; Zvartau-Hind, M.; van Dyck, C.; Gold, M. Effects of rosiglitazone-extended release as adjunctive therapy to acetylcholinesterase inhibitors over 48 weeks on cognition in Apoe4-stratified subjects with mild-to-moderate Alzheimer's disease. *Alzheimer Dement.* **2009**, *5*, e17–e18. [CrossRef]
296. Festuccia, W.T.; Oztezcan, S.; Laplante, M.; Berthiaume, M.; Michel, C.; Dohgu, S.; Denis, R.G.; Brito, M.N.; Brito, N.A.; Miller, D.S.; et al. Peroxisome proliferatoractivated receptor-gamma-mediated positive energy balance in the rat is associated with reduced sympathetic drive to adipose tissues and thyroid status. *Endocrinology* **2008**, *149*, 2121–2130. [CrossRef] [PubMed]
297. Kim, B.; Sullivan, K.A.; Backus, C.; Feldman, E.L. Cortical neurons develop insulin resistance and blunted Akt signaling: A potential mechanism contributing to enhanced ischemic injury in diabetes. *Antioxid. Redox Signal.* **2011**, *14*, 1829–1839. [CrossRef] [PubMed]
298. Łabuzek, K.; Suchy, D.; Gabryel, B.; Bielecka, A.; Liber, S.; Okopień, B. Quantification of metformin by the HPLC method in brain regions, cerebrospinal fluid and plasma of rats treated with lipopolysaccharide. *Pharmacol. Rep.* **2010**, *62*, 956–965.
299. Nath, N.; Khan, M.; Paintlia, M.K.; Singh, I.; Hoda, M.N.; Giri, S. Metformin attenuated the autoimmune disease of the central nervous system in animal models of multiple sclerosis. *J. Immunol.* **2009**, *182*, 8005–8014. [CrossRef] [PubMed]
300. Gupta, A.; Bisht, B.; Dey, C.S. Peripheral insulin-sensitizer drug metformin ameliorates neuronal insulin resistance and Alzheimer's-like changes. *Neuropharmacology* **2011**, *60*, 910–920. [CrossRef] [PubMed]
301. Chen, Y.; Zhou, K.; Wang, R.; Liu, Y.; Kwak, Y.D.; Ma, T.; Thompson, R.C.; Zhao, Y.; Smith, L.; Gasparini, L.; et al. Antidiabetic drug metformin (GlucophageR) increases biogenesis of Alzheimer's amyloid peptides via up-regulating BACE1 transcription. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 3907–3912. [CrossRef] [PubMed]
302. Perry, T.A.; Greig, N.H. A new Alzheimer's disease interventional strategy: GLP-1. *Curr. Drug Targets* **2004**, *5*, 565–571. [CrossRef] [PubMed]
303. McClean, P.L.; Parthsarathy, V.; Faivre, E.; Hölscher, C. The diabetes drug liraglutide prevents degenerative processes in a mouse model of Alzheimer's disease. *J. Neurosci.* **2011**, *31*, 6587–6594. [CrossRef] [PubMed]
304. Li, X.-L.; Aou, S.; Oomura, Y.; Hori, N.; Fukunaga, K.; Hori, T. Impairment of longterm potentiation and spatial memory in leptin receptor-deficient rodents. *Neuroscience* **2002**, *113*, 607–615.
305. Paz-Filho, G.J.; Babikian, T.; Asarnow, R.; Delibasi, T.; Esposito, K.; Erol, H.K.; Wong, M.L.; Licinio, J. Leptin replacement improves cognitive development. *PLoS ONE* **2008**, *3*, e3098. [CrossRef]
306. Greco, S.J.; Bryan, K.J.; Sarkar, S.; Zhu, X.; Smith, M.A.; Ashford, J.W.; Johnston, J.M.; Tezapsidis, N.; Casadesus, G. Leptin reduces pathology and improves memory in a transgenic mouse model of Alzheimer's disease. *J. Alzheimers Dis.* **2010**, *19*, 1155–1167. [PubMed]
307. Lieb, W.; Beiser, A.S.; Vasan, R.S.; Tan, Z.S.; Au, R.; Harris, T.B.; Roubenoff, R.; Auerbach, S.; DeCarli, C.; Wolf, P.A.; et al. Association of plasma leptin levels with incident Alzheimer disease and MRI measures of brain aging. *JAMA* **2009**, *302*, 2565–2572. [CrossRef] [PubMed]
308. Adler, B.L.; Yarchan, M.; Hwang, H.M.; Louneva, N.; Blair, J.A.; Palm, R.; Smith, M.A.; Lee, H.G.; Arnold, S.E.; Casadesus, G. Neuroprotective effects of the amylin analogue pramlintide on Alzheimer's disease pathogenesis and cognition. *Neurobiol. Aging* **2014**, *35*, 793–801. [CrossRef] [PubMed]
309. Ravussin, E.; Smith, S.R.; Mitchell, J.A.; Shringarpure, R.; Shan, K.; Maier, H.; Koda, J.E.; Weyer, C. Enhanced weight loss with pramlintide/metreleptin: An integrated neurohormonal approach to obesity pharmacotherapy. *Obesity* **2009**, *17*, 1736–1743. [CrossRef] [PubMed]
310. Unger, J.W.; Livingston, J.N.; Moss, A.M. Insulin receptors in the central nervous system: Localization, signalling mechanisms and functional aspects. *Prog. Neurobiol.* **1991**, *36*, 343–362. [CrossRef]

311. Baura, G.D.; Foster, D.M.; Porte, D., Jr.; Kahn, S.E.; Bergman, R.N.; Cobelli, C.; Schwartz, M.W. Saturable transport of insulin from plasma into the central nervous system of dogs *in vivo*. A mechanism for regulated insulin delivery to the brain. *J. Clin. Invest.* **1993**, *92*, 1824–1830. [CrossRef] [PubMed]
312. Hallschmid, M.; Schultes, B.; Marshall, L.; Molle, M.; Kern, W.; Bredthauer, J.; Fehm, H.L.; Born, J. Transcortical direct current potential shift reflects immediate signaling of systemic insulin to the human brain. *Diabetes* **2004**, *53*, 2202–2208. [CrossRef] [PubMed]
313. Marks, D.R.; Tucker, K.; Cavallin, M.A.; Mast, T.G.; Fadool, D.A. Awake intranasal insulin delivery modifies protein complexes and alters memory, anxiety, and olfactory behaviors. *J. Neurosci.* **2009**, *29*, 6734–6751. [CrossRef] [PubMed]
314. Khafagy, el-S.; Morishita, M.; Onuki, Y.; Takayama, K. Current challenges in non-invasive insulin delivery systems: A comparative review. *Adv. Drug. Deliv. Rev.* **2007**, *59*, 1521–1546.
315. Kupila, A.; Sipila, J.; Keskinen, P.; Simell, T.; Knip, M.; Pulkki, K.; Simell, O. Intranasally administered insulin intended for prevention of type 1 diabetes—a safety study in healthy adults. *Diabetes Metab. Res. Rev.* **2003**, *19*, 415–420. [CrossRef] [PubMed]
316. Claxton, A.; Baker, L.D.; Hanson, A.; Tritschuh, E.H.; Cholerton, B.; Morgan, A.; Callaghan, M.; Arbuckle, M.; Behl, C.; Craft, S. Long-acting intranasal insulin detemir improves cognition for adults with mild cognitive impairment or early-stage Alzheimer’s disease dementia. *J. Alzheimers Dis.* **2015**, *44*, 897–906. [PubMed]
317. Watson, G.S.; Bernhardt, T.; Reger, M.A.; Cholerton, B.A.; Baker, L.D.; Peskind, E.R.; Asthana, S.; Plymate, S.R.; Frölich, L.; Craft, S. Insulin effects on CSF norepinephrine and cognition in Alzheimer’s disease. *Neurobiol. Aging* **2006**, *27*, 38–41. [CrossRef] [PubMed]
318. Messier, C.; Teutenberg, K. The role of insulin, insulin growth factor, and insulin-degrading enzyme in brain aging and Alzheimer’s disease. *Neural. Plast.* **2005**, *12*, 311–328. [CrossRef] [PubMed]
319. Galasko, D. Insulin and Alzheimer’s disease: An amyloid connection. *Neurology* **2003**, *60*, 1886–1887. [CrossRef] [PubMed]
320. Benedict, C.; Dodd, C.; Hallschmid, M.; Lepiorz, M.; Fehm, H.L.; Born, J.; Kern, W. Immediate but not long-term intranasal administration of insulin raises blood pressure in human beings. *Metabolism* **2005**, *54*, 1356–1361. [CrossRef] [PubMed]
321. Craft, S.; Baker, L.D.; Montine, T.J.; Minoshima, S.; Watson, G.S.; Claxton, A.; Arbuckle, M.; Callaghan, M.; Tsai, E.; Plymate, S.R.; et al. Intranasal insulin therapy for Alzheimer disease and amnestic mild cognitive impairment: A pilot clinical trial. *Arch. Neurol.* **2012**, *69*, 29–38. [CrossRef] [PubMed]
322. Dhamoon, M.S.; Noble, J.M.; Craft, S. Intranasal insulin improves cognition and modulates beta-amyloid in early AD. *Neurology* **2009**, *72*, 292–293; author reply 293–294. [CrossRef] [PubMed]
323. Reger, M.A.; Watson, G.S.; Green, P.S.; Wilkinson, C.W.; Baker, L.D.; Cholerton, B.; Fishel, M.A.; Plymate, S.R.; Breitner, J.C.S.; DeGroodt, W.; et al. Intranasal insulin improves cognition and modulates beta-amyloid in early AD. *Neurology* **2008**, *70*, 440–448. [CrossRef] [PubMed]
324. Schiöth, H.B.; Craft, S.; Brooks, S.J.; Frey, W.H., II; Benedict, C. Brain insulin signaling and Alzheimer’s disease: Current evidence and future directions. *Mol. Neurobiol.* **2012**, *46*, 4–10. [CrossRef] [PubMed]
325. Shemesh, E.; Rudich, A.; Harman-Boehm, I.; Cukierman-Yaffe, T. Effect of intranasal insulin on cognitive function: A systematic review. *J. Clin. Endocrinol. Metab.* **2012**, *97*, 366–376. [CrossRef] [PubMed]
326. Dash, S.; Xiao, C.; Morgantini, C.; Koulajian, K.; Lewis, G.F. Intranasal insulin suppresses endogenous glucose production in humans compared with placebo in the presence of similar venous insulin concentrations. *Diabetes* **2005**, *64*, 766–774. [CrossRef] [PubMed]
327. Gancheva, S.; Koliaki, C.; Bierwagen, A.; Nowotny, P.; Heni, M.; Fritzsche, A.; Häring, H.U.; Szendroedi, J.; Roden, M. Effects of intranasal insulin on hepatic fat accumulation and energy metabolism in humans. *Diabetes* **2015**, *64*, 1966–1975. [CrossRef] [PubMed]
328. Born, J.; Lange, T.; Kern, W.; McGregor, G.P.; Bickel, U.; Fehm, H.L. Sniffing neuropeptides: A transnasal approach to the human brain. *Nat. Neurosci.* **2002**, *5*, 514–516. [CrossRef] [PubMed]
329. Hallschmid, M.; Benedict, C.; Schultes, B.; Perras, B.; Fehm, H.L.; Kern, W.; Born, J. Towards the therapeutic use of intranasal neuropeptide administration in metabolic and cognitive disorders. *Regul. Pept.* **2008**, *149*, 79–83. [CrossRef] [PubMed]
330. Zhang, Y.; Lee, D.H. Sink hypothesis and therapeutic strategies for attenuating Abeta levels. *Neuroscientist* **2011**, *17*, 163–173. [CrossRef] [PubMed]

331. Ketterer, C.; Tschritter, O.; Preissl, H.; Heni, M.; Häring, H.U.; Fritzsche, A. Insulin sensitivity of the human brain. *Diabetes Res. Clin. Pract.* **2011**, *93*, S47–S51. [[CrossRef](#)]
332. Kurtzhals, P. Engineering predictability and protraction in a basal insulin analogue: The pharmacology of insulin detemir. *Int. J. Obes. Relat. Metab. Disord.* **2004**, *28*, S23–S28. [[CrossRef](#)] [[PubMed](#)]
333. De Leeuw, I.; Vague, P.; Selam, J.L.; Skeie, S.; Lang, H.; Draeger, E.; Elte, J.W. Insulin detemir used in basal-bolus therapy in people with type 1 diabetes is associated with a lower risk of nocturnal hypoglycaemia and less weight gain over 12 months in comparison to NPH insulin. *Diabetes Obes. Metab.* **2005**, *7*, 73–82. [[CrossRef](#)] [[PubMed](#)]
334. Shen, D.D.; Artru, A.A.; Adkison, K.K. Principles and applicability of CSF sampling for the assessment of CNS drug delivery and pharmacodynamics. *Adv. Drug Deliv. Rev.* **2004**, *56*, 1825–1857. [[CrossRef](#)] [[PubMed](#)]
335. Monami, M.; Marchionni, N.; Mannucci, E. Long-acting insulin analogues vs. NPH human insulin in type 1 diabetes. A meta-analysis. *Diabetes. Obes. Metab.* **2009**, *11*, 372–378. [[CrossRef](#)] [[PubMed](#)]
336. Hennige, A.M.; Sartorius, T.; Tschritter, O.; Preissl, H.; Fritzsche, A.; Ruth, P.; Haring, H.U. Tissue selectivity of insulin detemir action *in vivo*. *Diabetologia* **2006**, *49*, 1274–1282. [[CrossRef](#)] [[PubMed](#)]
337. Hallschmid, M.; Jauch-Chara, K.; Korn, O.; Molle, M.; Rasch, B.; Born, J.; Schultes, B.; Kern, W. Euglycemic infusion of insulin detemir compared with human insulin appears to increase direct current brain potential response and reduces food intake while inducing similar systemic effects. *Diabetes* **2010**, *59*, 1101–1107. [[CrossRef](#)] [[PubMed](#)]



© 2016 by the author; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).