

Synthesis and Evaluation of New Pyrazoline Derivatives as Potential Anticancer Agents in HepG-2 Cell Line

Wei jie Xu ^{1,†}, Ying Pan ^{1,†}, Hong Wang ^{1,†}, Haiyan Li ², Qing Peng ³, Duncan Wei ¹, Cheng Chen ¹ and Jinhong Zheng ^{1,*}

¹ Department of Chemistry, Shantou University Medical College, Shantou 515041, Guangdong, China; 13536920534@163.com (W.X.); ypan@stu.edu.cn (Y.P.); 13592851552@163.com (H.W.); weiduncan2012@163.com (D.W.); 13411963192@163.com (C.C.); jhzheng@stu.edu.cn (J.Z)


² Department of Pharmacology, Shantou University Medical College, Shantou 515041, Guangdong, China; 13066360950@163.com (H.L.)

³ Department of Hepatobiliary Surgery II, Zhujiang Hospital of Southern Medical University, Guangzhou 510280, Guangdong, China; zhujiangzhuanhua@163.com (Q.P.)

* Correspondence: jhzheng@stu.edu.cn; Tel.: +86-754-8890-0499; Fax: +86-754-8855-7562

† These authors contributed equally to this work.

Structure of compound **b17**:



Analysis: b17

Acquisition Date: 2/28/2017

Name: b17

Comment: ESI Negative

Operator: TIPC

2017022805 #12 RT: 0.12 AV: 1 NL: 5.11E7

T: FTMS - p ESI Full ms [50.00-750.00]

