

Supplementary Materials

Two New Anisic Acid Derivatives from Endophytic fungus *Rhizopycnis vagum* Nitaf22 and Their Antibacterial activity


Ali Wang, Peng Li, Xuping Zhang, Peipei Han, Daowan Lai and Ligang Zhou *

Department of Plant Pathology, College of Plant Protection, China Agricultural University, Beijing 100193, China; wangali526@163.com (A.W.); ytmuyue1989@163.com (P.L.); zhangxuping5@163.com (X.Z.); mhanpeipei@126.com (P.H.); dwlai@cau.edu.cn (D.L.)

* Correspondence: lgzhou@cau.edu.cn; Tel.: +86-10-6273-1199

Contents

Figure S1. HRESIMS spectrum of 1	2
Figure S2. IR spectrum of 1	2
Figure S3. ^1H NMR spectrum of 1 (CD_3OD , 400 MHz).....	3
Figure S4. ^{13}C NMR spectrum of 1 (CD_3OD , 100 MHz).....	3
Figure S5. HMBC spectrum of 1 (400 MHz).....	4
Figure S6. NOESY spectrum of 1 (400 MHz)	4
Figure S7. HRESIMS spectrum of 2	5
Figure S8. IR spectrum of 2	5
Figure S9. ^1H NMR spectrum of 2 (CD_3OD , 400 MHz).....	6
Figure S10. ^{13}C NMR spectrum of 2 (CD_3OD , 100 MHz).....	6
Figure S11. HMBC spectrum of 2 (400 MHz).....	7



<i>m/z</i>	<i>Calc m/z</i>	Diff (ppm)	<i>z</i>	Abund	Formula	Ion
249.0771	249.0768	1.14	-1	329658.7	C ₁₃ H ₁₃ O ₅	(M-H) ⁻

Figure S1. HRESIMS spectrum of **1**.




Figure S2. IR spectrum of **1**.




Figure S3. ^1H NMR spectrum of **1** (CD_3OD , 400 MHz).




Figure S4. ^{13}C NMR spectrum of **1** (CD_3OD , 100 MHz).




Figure S5. HMBC spectrum of **1** (400 MHz).





Figure S6. NOESY spectrum of **1** (400 MHz).



<i>m/z</i>	Calc <i>m/z</i>	Diff (ppm)	<i>z</i>	Abund	Formula	Ion
251.093	251.0925	2.01	-1	991886.9	C ₁₃ H ₁₅ O ₅	(M-H) ⁻

Figure S7. HRESIMS spectrum of 2.




Figure S8. IR spectrum of 2.




Figure S9. ^1H NMR spectrum of **2** (CD_3OD , 400 MHz).




Figure S10. ^{13}C NMR spectrum of **2** (CD_3OD , 100 MHz).




Figure S11. HMBC spectrum of **2** (400 MHz).