

Supplementary Materials

Phenolic compositions and antioxidant activities differ significantly among sorghum grains with different applications

Shuyu Shen ¹, Rui Huang ¹, Charlie Li ², WenYan Wu ¹, Honglin Chen ¹, John Shi ^{3,*}, Shiguo Chen ^{1,*} and Xingqian Ye ^{1,*}

¹ Zhejiang University, College of Biosystem Engineering and Food Science, Zhejiang Key Laboratory for Agro-Food Processing, Fuli Institute of Food Science, Hangzhou 310058, China;

² University of California-Davis, Department of Environmental Toxicology, Davis, CA 94564, USA;

³ Guelph Research and Development Center, Agriculture and Agri-Food Canada, Guelph, Ontario, Canada;

* Correspondence: johnshi2006@yahoo.ca; chenshiguo210@163.com, Tel.: 86-571-88982151.; psu@zju.edu.cn, Tel.: 86-571-88982155

Figure S1. MS/MS spectra of Protocatechuic acid in negative electrospray ionization

Figure S2. MS/MS spectra of Caffeic acid in negative electrospray ionization

Figure S3. MS/MS spectra of Luteolinidin in negative electrospray ionization

Figure S4. MS/MS spectra of Apigeninidin in negative electrospray ionization

Figure S5. MS/MS spectra of *p*-coumaric acid in negative electrospray ionization

Figure S6. MS/MS spectra of Ferulic acid in negative electrospray ionization

Figure S7. MS/MS spectra of Taxifolin in negative electrospray ionization

Figure S8. MS/MS spectra of Luteolin in negative electrospray ionization

Figure S9. MS/MS spectra of Apigenin in negative electrospray ionization

Figure S10. MS/MS spectra of Narigenin in negative electrospray ionization




Figure S1. MS/MS spectra of Protocatechuic acid in negative electrospray ionization




Figure S2. MS/MS spectra of Caffeic acid in negative electrospray ionization




Figure S3. MS/MS spectra of Luteolinidin in negative electrospray ionization




Figure S4. MS/MS spectra of Apigeninidin in negative electrospray ionization




Figure S5. MS/MS spectra of *p*-coumaric acid in negative electrospray ionization




Figure S6. MS/MS spectra of Ferulic acid in negative electrospray ionization




Figure S7. MS/MS spectra of Taxifolin in negative electrospray ionization




Figure S8. MS/MS spectra of Luteolin in negative electrospray ionization




Figure S9. MS/MS spectra of Apigenin in negative electrospray ionization




Figure S10. MS/MS spectra of Narigenin in negative electrospray ionization