

Supporting Information

Synthesis and antioxidant activity of caffeic acid derivatives

Katarzyna Sidoryk¹, Anna Jaromin², Nina Filipczak², Piotr Cmoch¹, Marcin Cybulski¹

¹Pharmaceutical Research Institute, Rydygiera 8, 01-793 Warsaw, Poland

²Department of Lipids and Liposomes, Faculty of Biotechnology, University of Wrocław, Joliot-Curie 14A, 50-383 Wrocław, Poland

To whom correspondence should be addressed: Phone: +48-22-456-3928 Fax: +48-22-456-3838 E-mail: k.sidoryk@ifarm.eu

Supporting material contains ¹H and ¹³C NMR spectra of compounds:

1. 3-(2'-Hydroxyphenyl)-(E)-propenoic acid methyl ester (**9**)
2. 3-(3'-Hydroxyphenyl)-(E)-propenoic acid methyl ester (**10**)
3. 3-(4'-Hydroxyphenyl)-(E)-propenoic acid methyl ester (**8**)
4. 3-(2',4'-dihydroxyphenyl)-(E)-propenoic acid methyl ester (**11**)
5. 3-(2',3',4'-trihydroxyphenyl)-(E)-propenoic acid methyl ester (**12**)
6. 3-(2',4',5'-trihydroxyphenyl)-(E)-propenoic acid methyl ester (**13**)
7. 3-(4'-hydroxy-3-methoxyphenyl)-(E)-propenoic acid methyl ester (**14**)
8. 4-(2'-hydroxyphenyl)-3(E)-buten-2-one (**6**)
9. 4-(3'-hydroxyphenyl)-3(E)-buten-2-one (**15**)
10. 4-(4'-hydroxyphenyl)-3(E)-buten-2-one (**16**)
11. 4-(2',4'-dihydroxyphenyl)-3(E)-buten-2-one (**17**)
12. 4-(2',3',4'-trihydroxyphenyl)-3(E)-buten-2-one (**18**)
13. 4-(3'-methoxy-4'-hydroxyphenyl)-3(E)-buten-2-one (**20**)



















































