

Article

N-methyl-D-glucamine–Calix[4]resorcinarene Conjugates: Self-Assembly and Biological Properties

Ruslan R. Kashapov ^{1,2,*}, Yuliya S. Razuvayeva ^{1,2}, Albina Y. Ziganshina ¹, Rezeda K. Mukhitova ¹, Anastasiia S. Sapunova ¹, Alexandra D. Voloshina ¹, Victor V. Syakaev ¹, Shamil K. Latypov ¹, Irek R. Nizameev ^{1,2}, Marsil K. Kadirov ^{1,2} and Lucia Y. Zakharova ^{1,2}

¹ Arbuzov Institute of Organic and Physical Chemistry, FRC Kazan Scientific Center of RAS, 8 Arbuzov str., Kazan 420088, Russia; kashapov@iopc.ru

² Kazan National Research Technological University, 68 K. Marks str., Kazan 420015, Russia

* Correspondence: kashapov@iopc.ru; Tel.: +7843-273-22-93; Fax: +7843-273-22-53 (F.L.)

Supplementary materials




Figure S1. ¹H and ¹³C NMR spectra of GCR-1.




Figure S2. ^1H and ^{13}C NMR spectra of GCR-2.




Figure S3. IR spectra of GCR-1 and GCR-2.




Figure S4. Concentration dependence of pH of water solutions of CR-1 and GCR-1, 25°C.




Figure S5. Concentration dependence of specific conductivity (χ) of GCR-1 and GCR-2 in 50% DMSO – 50% water, 25°C.




Figure S6. Concentration dependence of absorption at 500 nm for GCR-1 and GCR-2 in 50% DMSO – 50% water, 25°C.