

Supplementary Materials

Anti-Amyloidogenic Effects of Asarone Derivatives From *Perilla frutescens* Leaves Against Beta-Amyloid Aggregation and Nitric Oxide Production


Jae Eun Lee ^{1,†}, Nayeon Kim ^{1,†}, Ji Yun Yeo ¹, Dae-Gun Seo ¹, Sunggun Kim ¹, Jae-Sun Lee ¹, Kwang Woo Hwang ² and So-Young Park ^{1,*}

¹ College of Pharmacy, Dankook University, 119 Dandae-ro, Dongnam-gu, Cheonan-si, Chungnam 31116, Republic of Korea; jnlee88@korea.kr (J.E.L.); nayeon02200@gmail.com (N.K.); dankook_jjiyun@naver.com (J.Y.Y.); gun3691@naver.com (D.-G.S.); jeju@hanmail.net (S.K.); sailious34@naver.com (J.-S.L.)


² College of Pharmacy, Chung-Ang University, 84 Heukseok-ro, Dongjak-gu, Seoul 06974, Republic of Korea; khwang@cau.ac.kr

† These authors contributed equally to this work.

* Correspondence: soypark23@dankook.ac.kr, Tel.: +82-41-550-1434, Fax: +82-41-559-7899



Supplementary Figure 1. Direct effect of asarone derivatives on Th T fluorescence. In order to determine quenching effect of asarone derivatives asarone derivatives were incubated with Th T without A β and the fluorescence values were determined. The Th T fluorescence values were compared to Th T only.



Supplementary Figure 2. The effect of asarone derivatives on Aβ aggregation determined by western blot analysis. The western blot analysis with native gels using anti-Aβ(1–20) antibody and the band intensity was compared to Aβ only samples (Due to the lack of amount, compound 3 was excluded from this experiment). * $p < 0.05$ compared to Aβ only.