

Supplementary Material

Ionic Liquids and Ohmic Heating in Combination for Pd-catalyzed Cross-coupling Reactions: Sustainable Synthesis of Flavonoids

Vera L. M. Silva ¹, Raquel G. Soengas ² and Artur M. S. Silva ^{1,*}

¹ LAQV-REQUIMTE, Department of Chemistry, University of Aveiro, 3810-193 Aveiro, Portugal; verasilva@ua.pt

² Department of Organic and Inorganic Chemistry, University of Oviedo, c/Julián Clavería 6, 33006 Oviedo, Spain; rsoengas@uniovi.es

* Correspondence: artur.silva@ua.pt

Academic Editor: Vito Capriati

Received: 24 February 2020; Accepted: 26 March 2020; Published: date.

Figure S1. 3-Phenyl-4*H*-chromen-4-one (3a)





Figure S2. 3-(3,4-Dimethoxyphenyl)-4*H*-chromen-4-one (3b)

Figure S3. 3-(4-Chlorophenyl)-4H-chromen-4-one (3c)




Figure S4. 3-Styryl-4H-chromen-4-one (3d)




Figure S5. 3-(3,4-Dimethoxystyryl)-4H-chromen-4-one (3e)




Figure S6. 7-(Benzyloxy)-3-(3,4-dimethoxystyryl)-4H-chromen-4-one (3f)




Figure S7. 3-(2,2-Bis(3,4-dimethoxyphenyl)vinyl)-4H-chromen-4-one (3g)




Figure S8. 3-(2,2-Bis(4-chlorophenyl)vinyl)-6-chloro-4H-chromen-4-one (3h)




Figure S9. 7-(Benzyloxy)-3-(2,2-bis(4-chlorophenyl)vinyl)-4H-chromen-4-one (3i)




Figure S10. 3-(2,2-Bis(3,4-dimethoxyphenyl)vinyl)-6-methyl-4H-chromen-4-one (3j)

