

PEEK-WC-based Mixed Matrix Membranes Containing Polyimine Cages for Gas Separation

Marcello Monteleone ¹, Riccardo Mobili ², Chiara Milanese ², Elisa Esposito ¹, Alessio Fuoco ^{1,*}, Sonia La Cognata^{2,*}, Valeria Amendola ² and Johannes C. Jansen ¹

¹ Institute on Membrane Technology, National Research Council of Italy (CNR-ITM), via P. Bucci 17/C, Rende (CS), 87036, Italy; m.monteleone@itm.cnr.it (M.M.) e.esposito@itm.cnr.it (E.E.), jc.jansen@itm.cnr.it (J.C.J.)

² Dipartimento di Chimica, Università di Pavia, via Taramelli 12, I-27100 Pavia, Italy; riccardo.mobili01@universitadipavia.it (R.M.), chiara.milanese@unipv.it (C.M.), valeria.amendola@unipv.it (V.A.)

* Correspondence: a.fuoco@itm.cnr.it (A.F.), sonia.lacognata01@universitadipavia.it (S.L.C.)

Index

1. Characterization of the Cages.....	2
1.1 ¹ H-NMR spectra	2
1.2 FTIR-ATR spectra.....	4
2. Differential scanning calorimetric (DSC) measurements	6

1. Characterization of the Cages

1.1 ^1H -NMR spectra




Figure S1. ^1H -NMR spectrum of Fura in CDCl₃.




Figure S2. ^1H -NMR spectrum of m-xy in CDCl₃.




Figure S3. ¹H-NMR spectrum of p-xy in CDCl₃.




Figure S4. ¹H-NMR spectrum of the diphenyl cage in CDCl₃.

1.2 FTIR-ATR spectra




Figure S5. FTIR-ATR spectrum of Fura.




Figure S6 FTIR-ATR spectrum of m-xy.




Figure S7 FTIR-ATR spectrum of p-xy




Figure S8 FTIR-ATR spectrum of the diphenyl cage.

2. Differential scanning calorimetric (DSC) measurements




Figure S9 DSC curve of Fura.




Figure S10 DSC curve of m-xy




Figure S11 DSC curve of p-xy.




Figure S12 DSC curve of the diphenyl cage.