

Impact of Static-Oriented Electric Fields on the Kinetics of Some Representative Suzuki–Miyaura and Metal-Cluster Mediated Reactions

Navya Areppalli ^{1,†}, Sukanta Mondal ^{2,†}, Debdutta Chakraborty ^{1,*} and Pratim Kumar Chattaraj ^{1,*}

¹ Department of Chemistry, Birla Institute of Technology, Mesra, Ranchi 835215, Jharkhand, India

² Department of Education, A. M. School of Educational Sciences, Assam University, Silchar 788011, Assam, India

* Correspondence: debdutta.chakraborty@bitmesra.ac.in (D.C.);
pkc@chem.iitkgp.ac.in (P.K.C.)

† These authors contributed equally to this work.

Supporting Information




Figure S1: Results of EDA for reaction 1a.




Figure S2: Results of EDA for reaction 1b.




Figure S3: Results of EDA for reaction 1c.




Figure S4: Results of EDA for reaction 2a.




Figure S5: Results of EDA for reaction 2b.




Figure S6: Results of EDA for reaction 3a.




Figure S7: Results of EDA for reaction 3b.




Figure S8: Results of EDA for reaction 4a.




Figure S9: Results of EDA for reaction 4b.




Figure S10: Results of EDA for reaction 4c.

Reaction Path: 5




Figure S11: Results of EDA for reaction 5.