

Supporting Information

Systematic Study of Resveratrol Nanoliposomes Transdermal Delivery System for Enhancing Anti-Aging and Skin-Brightening Efficacy

Xinchao Zhang ¹, Siyuan Chen ², Dan Luo ³, Dan Chen ³, Hong Zhou ¹, Shuting Zhang ¹, Xuan Chen ¹, Wangwang Lu ^{4,*} and Wei Liu ^{1,3,*}

¹ College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan 430074, China

² College of Materials Science and Engineering, Suqian Advanced Materials Industry Technology Innovation Center, Nanjing Tech University, Nanjing 211816, China

³ National Engineering Research Center for Nanomedicine, Huazhong University of Science and Technology, Wuhan 430075, China

⁴ Guangzhou Jiyan Cosmetics Technology Co., Ltd., Guangzhou 510275, China

* Correspondence: leo.lu@yatsenglobal.com (W.L.); wliu@hust.edu.cn (W.L.)

Figure S1




Figure S1. In vitro cytotoxicity of the free Res, Res-NLPs against (a) HaCaT, (b) HSF and (c) B16-F10 cells detected by CCK-8. The cells were treated with the free Res or Res-NLPs at the same Res concentrations over a wide range (5–200 μM). Mean ± SD (n = 3). **p* < 0.05, ***p* < 0.01, compared with the control group; #*p* < 0.05, compared with the free Res group.

Figure S2




Figure S2. Informed Consent Statement. (a) Informed consent statement from all volunteers (Chinese version), (b) especially, informed consent statement from two volunteers whose facial information was included in the manuscript (English version).