

Phytoconstituents and Ergosterol Biosynthesis-Targeting Antimicrobial Activity of Nutmeg (*Myristica fragrans* Houtt.) against Phytopathogens

Adriana Cruz ^{1,2}, Eva Sánchez-Hernández ³, Ana Teixeira ^{1,4}, Rui Oliveira ^{1,2}, Ana Cunha ^{1,2}, and Pablo Martín-Ramos ^{3,*}

¹ Department of Biology, School of Sciences, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal; cruzadriana73@gmail.com (A.C.).

² Centre of Molecular and Environmental Biology (CBMA), University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal; accunha@bio.uminho.pt (A.C.); ruipso@bio.uminho.pt (R.O.).

³ Department of Agricultural and Forestry Engineering, ETSIIAA, Universidad de Valladolid, Avenida de Madrid 44, 34004 Palencia, Spain; eva.sanchez.hernandez@uva.es (E.S.-H.), pmr@uva.es (P.M.-R.).

⁴ Centre for Research and Technology of Agro-Environmental and Biological Sciences (CITAB), Inov4Agro, University of Trás-os-Montes and Alto Douro (UTAD), Quinta de Prados, 5000-801 Vila Real, Portugal; anaspereirateixeira@gmail.com (A.T.).

* Correspondence: pmr@uva.es

SUPPLEMENTARY MATERIAL




Figure S1. ATR–FTIR spectrum of the freeze-dried *M. fragrans* hydroethanolic extract




Figure S2. GC–MS chromatogram of *M. fragrans* hydroethanolic extract.