


Molecules **1998**, *3*, M81**(1R,2S)-N-Benzyl-N-[(2,2-dimethyl-1,3-dioxolan-4-yl)-(2-benzothiényl)methyl] Hydroxylamine****Pedro Merino*, Santiago Franco, Francisco L. Merchan and Tomas Tejero**

Department of Organic Chemistry. Faculty of Sciences-ICMA. University of Zaragoza. E-50009 Zaragoza. Spain

Tel. +34 976 762075, Fax +34 976 761194, Email: pmerino@posta.unizar.es

Received: 15 May 1998 / Published: 19 May 1998

A cooled solution (-90°C) of benzothiophene (0.402 g, 3 mmol) in THF (10 mL) was treated with butyllithium (2 mL of a 1.6 M solution in hexane, 3.2 mmol) under an inert atmosphere. The resulting solution was stirred at -60 °C for 1 h during which time the reaction mixture was cooled to -90 °C and treated via a syringe with a solution of nitron **1** (0.235 g, 1 mmol) in Et₂O (10 mL), previously treated with Et₂AlCl (1 mL of a 1.0 M solution in hexane, 1 mmol) at ambient temperature for 5 min. The rate of addition was adjusted so as to keep the internal temperature below -80°C. The reaction mixture was stirred for 1 h at -80 °C and then quenched with a 1N aqueous solution of sodium hydroxide (10 mL). The mixture was stirred at ambient temperature for 10 min and diluted with ethyl acetate (15 mL). The organic layer was separated, and the aqueous layer was extracted with ethyl acetate (3 x 10 mL). The combined organic extracts were washed with brine and dried over magnesium sulfate, and the solvent was evaporated under reduced pressure to give the crude mixture of diastereomeric hydroxylamines. The NMR analysis of this mixture revealed a diastereoselectivity of 70%. The absolute configuration of compound **3** was assumed according to previous results [1]. Purification by column chromatography (hexane/diethyl ether 60:40) on silica gel afforded **3** as a white solid (0.218 g, 59%).

Mp 127-129°C

TLC (hexane/diethyl ether 60:40) R_f 0.39[α]_D²⁰ = -22.3 (c 0.60, CHCl₃)

¹H NMR (CDCl₃) δ 1.33 (s, 3H), 1.39 (s, 3H), 3.70 (d, 1H, J = 13.2 Hz), 3.86 (d, 1H, J = 13.2 Hz), 4.09 (dd, 1H, J = 5.9, 8.6 Hz), 4.13 (d, 1H, J = 7.6 Hz), 4.19 (dd, 1H, J = 6.2, 8.6 Hz), 4.72 (ddd, 1H, J = 5.9, 6.2, 7.6 Hz), 4.85 (bs, 1H, ex. D₂O), 7.18 (s, 1H), 7.21-7.46 (m, 9H).

¹³C NMR (CDCl₃) δ 25.3, 26.7, 62.0, 67.4, 68.0, 76.6, 109.6, 122.2, 124.5, 124.1, 124.2, 125.1, 127.5, 128.4, 129.4, 137.2, 137.3, 138.6, 139.0.

Anal. Calcd. for C₂₁H₂₃NO₃S: C, 68.27; H, 6.27; N, 3.79. Found: C, 68.33; H, 6.18; N, 3.95.

References and Notes

1. Merino, P.; Franco, S.; Merchan, F. L.; Tejero, T. *Molecules*, 1998, 3, M80.

Sample Availability: Available from the authors and MDPI: [MDPI Reg. No. 15815](#).

©1998 MDPI. All rights reserved. *Molecules* website <http://www.mdpi.org/molecules/>