Molecules 2000, 5, M164

6-Amino-2-phenylimidazo[4,5-e][1,3]diazepine-4,8(1H, 5H)-dione

Huan-Ming Chen and Ramachandra S. Hosmane*

Laboratory for Drug Design and Synthesis, Department of Chemistry & Biochemistry, University of Maryland, Baltimore County, 1000 Hilltop Circle, Baltimore, MD 21250, USA. Tel. 410-455-2520, Fax 410-455-1148 (E-mail: hosmane@umbc.edu)

Received: 17 May 2000 / Accepted: 2 June 2000 / Published: 10 July 2000

The starting imidazole diester (1) was prepared by esterification of the corresponding dicarboxylic acid according to the literature procedure [1,2]. A solution of dimethyl 2-phenyl-1*H*-imidazole-4,5-dicarboxylate (1) (0.26 g, 1 mmol) and guanidine hydrochloride (0.48 g, 5 mmol) in 50 mL of absolute methanol was stirred for 30 min at room temperature, then 2 mL of 25% sodium methoxide solution in methanol was added. The resulting mixture was stirred for 48 h at room temperature. The precipitate formed was filtered and washed with methanol to give sodium salt of the product as a white solid. The sodium salt was dissolved in water and neutralized with 2 M hydrochloric acid to yield the title compound (2) as a white solid after filtration and washing with water, yield 0.21 g (78%).

M.p. > 260 $^{o}C.$

TLC (CHCl₃: MeOH: 30% NH₄OH, 2:2:1): R_f 0.61.

 1 H NMR (DMSO- d_6): 13.74 (br s, 1H, NH, exchangeable with D₂O), 10.68 (br s, 1H, NH, exchangeable with D₂O), 8.20 (m, 2H, Ph), 7.60 (br s, 1H, NH, exchangeable with D₂O), 7.48 (m, 3H, Ph), 6.50 (br s, 1H, NH, exchangeable with D₂O).

FABMS: 256 [MH⁺].

Anal. Calcd. for $C_{12}H_9N_5O_2^{-3}/4H_2O$ (268.75): C, 53.63; H, 3.94; N, 26.06. Found: C, 53.62, H, 4.02; N, 26.03.

References

- 1. Anderson, W. K.; Bhattacharjee, D.; Houston, D. M. J. Med. Chem. 1989, 32, 119-127.
- 2. Manecke, G.; Schlegel, R. Chem. Ber. 1974, 107, 892-897.

Sample availability: available from the authors and from MDPI (MDPI ID 18548).

© 2000 MDPI. All rights reserved. *Molecules* website www.mdpi.org/molecules/

1 von 1 04.05.2009 14:59