

## Supplementary Information

**Table S1.** Univariate analysis using the log-rank test according to the characteristics.

	<b><i>p</i>-value</b>
<b>Tumor size</b>	0.758
<i>T</i> ≤ 2 cm vs. <i>T</i> > 2 cm	
<b>Nodal status</b>	0.825
Negative vs. Positive	
<b>Stage</b>	0.371
I vs. II vs. III	
<b>Histologic grade</b>	0.442
I-II vs. III	
<b>Progesterone receptor</b>	0.754
Positive vs. Negative	
<b>HER-2 *</b>	0.756
Negative vs. Positive	
<b>70GS</b>	0.013
Good vs. Poor	
<b>Adjuvant chemotherapy</b>	0.024
Yes or No	
<b>Adjuvant endocrine therapy</b>	0.811
Yes or No	
<b>Adjuvant radiotherapy</b>	0.085
Yes or No	

HER-2, human epidermal growth factor receptor-2; \* HER2 positivity was defined by three positive findings in an immunohistochemical examination or amplification in fluorescence *in situ* hybridization.

© 2013 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).