

Supplementary Materials

Article

Time-Dependent DNA Origami Denaturation by Guanidinium Chloride, Guanidinium Sulfate, and Guanidinium Thiocyanate

Marcel Hanke [†], Niklas Hansen ^{†‡}, Emilia Tomm, Guido Grundmeier and Adrian Keller ^{*}

Technical and Macromolecular Chemistry, Paderborn University, Warburger Str. 100, 33098 Paderborn, Germany; marcelha@mail.uni-paderborn.de (M.H.); niklashansen94@aol.de (N.H.);

emiliat@mail.uni-paderborn.de (E.T.); g.grundmeier@tc.uni-paderborn.de (G.G.)

^{*} Correspondence: adrian.keller@uni-paderborn.de; Tel.: +49-5251-60-5722

[†] These authors contributed equally to this work.

[‡] Present address: Department of Biophysical Chemistry, J. Heyrovsky Institute of Physical Chemistry, Czech Academy of Sciences, Dolejškova 3, 182 23 Prague 8, Czech Republic.




Figure S1: Additional AFM images of DNA origami triangles exposed to GdmCl at 25 °C.




Figure S2: Additional AFM images of DNA origami triangles exposed to GdmCl at 40 °C.




Figure S3: Additional AFM images of DNA origami triangles exposed to GdmCl at 50 °C.




Figure S4: Additional AFM images of DNA origami triangles exposed to Gdm_2SO_4 at 25 °C.




Figure S5: Additional AFM images of DNA origami triangles exposed to Gdm_2SO_4 at 40 °C.




Figure S6: Additional AFM images of DNA origami triangles exposed to Gdm₂SO₄ at 50 °C.




Figure S7: Additional AFM images of DNA origami triangles exposed to GdmSCN at 25 °C.




Figure S8: Additional AFM images of DNA origami triangles exposed to GdmSCN at 40 °C.




Figure S9: Additional AFM images of DNA origami triangles exposed to GdmSCN at 50 °C.