

*Article*

# Specific Amino Acid Substitutions in OXA-51-Type $\beta$ -Lactamase Enhance Catalytic Activity to a Level Comparable to Carbapenemase OXA-23 and OXA-24/40

Kwan-Wai Chan <sup>1,2,†</sup>, Chen-Yu Liu <sup>2,†</sup>, Ho-Yin Wong <sup>1</sup>, Wai-Chi Chan <sup>1</sup>, Kwok-Yin Wong <sup>1</sup> and Sheng Chen <sup>2,\*</sup>

<sup>1</sup> State Key Laboratory of Chemical Biology and Drug Discovery, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong; bill.kw.chan@connect.polyu.hk (K.-W.C.); marcus-ho-yin.wong@polyu.edu.hk (H.-Y.W.); edwardwchan@yahoo.com.hk (W.-C.C.); kwok-yin.wong@polyu.edu.hk (K.-Y.W.)

<sup>2</sup> Department of Infectious Diseases and Public Health, City University of Hong Kong, Kowloon, Hong Kong; chenyliu4-c@my.cityu.edu.hk

\* Correspondence: shechen@cityu.edu.hk; Tel.: +852-3442-5782

† These authors contributed equally to this work.

**Keywords:** *A. baumannii*; carbapenem resistance; carbapenem-hydrolyzing class D  $\beta$ -lactamases (CHDLs); OXA-51 variants; structure/function relationship

**Table S1.** Characteristics of 14 *A. baumannii* clinical strains included in this study.

Strain	Year and Place of Isolation	ST Type (Oxford)	International Clone	Presence of Resistance Determinants						
				<i>carO</i>	<i>adeB</i>	<i>adeRS</i>	<i>bla</i> <sub>OXA-51</sub>	<i>ISAbal</i> - <i>bla</i> <sub>OXA-51</sub>	<i>bla</i> <sub>OXA-</sub>	<i>ISAbal</i> - <i>bla</i> <sub>OXA23</sub>
							Variants	24/40		
AB1	2004, Hong Kong	208	IC II	+	+	-	OXA-83	+	-	-
AB2	2008, Hong Kong	940	Non-IC II	+	+	+	OXA-99	-	OXA-23	+
AB3	2011, Hong Kong	208	IC II	+	-	-	OXA-66	+	OXA-23 <sup>#</sup>	+
AB4	2011, Hong Kong	253	IC II	+	+	+	OXA-66	-	OXA-23	+
AB5	2004, Hong Kong	940	Non-IC II	+	+	+	OXA-99	-	OXA-23	+
AB7	2011, Hong Kong	208	IC II	+	+	+	OXA-82	+	-	-
AB8	2003, Hong Kong	208	IC II	+	+	+	OXA-82	+	-	-
AB10	2000, Hong Kong	208	IC II	+	+	+	OXA-66	-	-	-
MH1	2000, Hong Kong	684	IC II	+	+	+	OXA-66	-	-	-
MH2	2010, Hong Kong	218	IC II	+	+	+	OXA-66	-	OXA-72	-
MH3	2011, Hong Kong	208	IC II	+	+	+	OXA-79	+	-	-
MH5	2006, Hong Kong	940	Non-IC II	+	-	-	OXA-99	-	OXA-23	+
MH6	2013, Henan, China	195	IC II	+	+	+	OXA-66	-	OXA-23	+
MH7	2013, Henan, China	369	IC II	+	+	+	OXA-66	-	OXA-23 <sup>#</sup>	+

Presence of endogenous genes and other resistance determinants were obtained from analysis of whole genome sequences. <sup>#</sup> Chromosomal located.