

Article

A Highly Sensitive and Selective Competition Assay for the Detection of Cysteine Using Mercury-Specific DNA, Hg^{2+} and Sybr Green I

Hui Xu ^{1,*}, Shuli Gao ¹, Quanwen Liu ¹, Dun Pan ², Lihua Wang ², Shuzhen Ren ³, Min Ding ³, Jingwen Chen ³ and Gang Liu ^{3,*}

¹ School of Chemistry and Material Sciences, Ludong University, Yantai 264025, China; E-Mails: shuli0049@126.com (S.G.); ldu_lqw@163.com (Q.L.)

² Laboratory of Physical Biology, Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China; E-Mails: pandun@sinap.ac.cn (D.P.); Wanglihua@sinap.ac.cn (L.W.)

³ Shanghai Institute of Measurement and Testing Technology, Shanghai 201203, China; E-Mails: renshuzhen@simt.com.cn (S.R.); dingmin@simt.com.cn (M.D.); chendingwen@simt.com.cn (J.C.)

* Authors to whom correspondence should be addressed; E-Mail: xuhui@ldu.edu.cn (H.X.); Tel.: 86-535-667-2176 (H.X.); Fax: 86-535-669-7667 (H.X.); E-Mail: liug@simt.com.cn (G.L.); Tel.: 86-21-3883-9800-35232 (G.L.); Fax: 86-21-5079-8552 (G.L.).

Received: 3 September 2011; in revised form: 18 October 2011 / Accepted: 23 October 2011 /

Published: 26 October 2011

Abstract: We here report a rapid, sensitive, selective and label-free fluorescence detection method for cysteine (Cys). The conformation of mercury-specific DNA (MSD) changes from a random coil form to a hairpin structure in the presence of Hg^{2+} due to the formation of a thymine- Hg^{2+} -thymine (T- Hg^{2+} -T) complex. Cys can selectively coordinate with Hg^{2+} and extract it from the thymine- Hg^{2+} -thymine complex. The hairpin structure dehybridizes and the fluorescence intensity of Sybr Green I (SG) decreases upon addition of Cys because SG efficiently discriminates mercury-specific DNA and mercury-specific DNA/ Hg^{2+} complex. The detection can be finished within 5 min with high sensitivity and selectivity. In addition, we can obtain variable dynamic ranges for Cys by changing the concentration of MSD/ Hg^{2+} .

Keywords: cysteine; competition assay; mercury-specific DNA; Hg²⁺; Sybr Green I

1. Introduction

Widely distributed in living cells, thiols such as cysteine (Cys) are involved in many biological functions. Their levels in biological fluids such as human plasma and urine are of great importance for clinical diagnostics of a variety of diseases [1-3]. Determination of these species can be achieved by effective separation/detection techniques, e.g., HPLC and capillary electrophoresis [4-6], spectrophotometry [7-9], electrochemical voltammetry [10-13], colorimetric methods [14-17], flow injection [18-20] and fluorescence analysis method [21-25]. However, some of them suffer from low sensitivity and/or selectivity or require cumbersome laboratory procedures. There is thus an intense demand for more sensitive, selective, convenient and low-cost methods to detect Cys.

Growing research interest has been focused on the development of sensitive, selective, and cost-effective biosensors based on target-responsive DNA structural switching [26-29]. The core technology of these sensors is a kind of special DNA molecules which change their conformations upon binding with the targets. For example, aptamers are *in vitro* selected functional oligonucleotides that can bind specifically to target molecules. In this work, we use a mercury-specific DNA (MSD) which presents a random coil form in the absence of Hg²⁺, and forms a hairpin structure in the presence of Hg²⁺ due to the formation of a thymine-Hg²⁺-thymine (T-Hg²⁺-T) complex. A fluorescent dye, Sybr Green I (SG), was applied to recognize the structural change due to the different interaction of SG with MSD and MSD/Hg²⁺ complex [30].


It is well-known that Cys can form a very stable complex with Hg²⁺ [31]. By using this property, Mirkin *et al.* [32] developed a highly sensitive and selective colorimetric detection method for Cys. However, their method needs an elevated temperature, which requires a long time. We recently developed a fluorescent turn-on “molecular beacon” probe for the detection of Cys, which is also based on the competitive ligation of Hg²⁺ ions by Cys and thymine-thymine (T-T) mismatches [33]. The method shows high sensitivity and selectivity, but still needs to be improved in terms of cost and convenience due to its requirement of a labelled “molecular beacon” and a solution heating process. Here we develop a simple, rapid, sensitive and selective method for detection of Cys by using a target-responsive DNA structural change. The MSD/Hg²⁺ complex is a hairpin structure, which dehybridizes when Hg²⁺ is extracted from the thymine-Hg²⁺-thymine complex by Cys due to the high formation constant of Hg²⁺-Cys complexes. The fluorescence intensity of SG then decreases upon addition of Cys due to the dehybridization of the MSD/Hg²⁺ complex.

2. Experimental Section

2.1. Chemicals and Apparatus

All chemicals used for these investigations were of analytical grade purity. L-Cysteine (Cys, minimum 98.5%) was purchased from Sigma Aldrich Chemical Company and used as received. MSD (5'-TTCTTTCTTCCCCTTGTTTGTT-3') was synthesized and purified by HPLC (Takara Biotech. Co., Dalian). SG (10,000×) was purchased from Invitrogen Inc. A stock solution of 400× was prepared

Figure 1. Fluorescence spectra of SG-Hg²⁺-MSD in the absence (black line) and in the presence of Cys (red line). [MSD] = 10 nM, [Hg²⁺] = 70 nM, [SG] = 1.225 × 10⁻⁷ M, [Cys] = 140 nM.



3.2. Optimization of the Assay

In previous study, Liu *et al.* [30] found that MSD kinetically accomplishes the hairpin structure at $[\text{Hg}^{2+}]/7[\text{MSD}] = 1$. One equivalent of MSD thus requires seven equivalents of Hg²⁺ to completely form the hairpin structure due to the T-Hg²⁺-T chemistry. So we fixed the concentration of MSD to be 10 nM and Hg²⁺ 70 nM, and then the concentration of SG was optimized. Figure 2 shows the fluorescence spectra of MSD and MSD-Hg²⁺ upon addition of different concentration of SG.

Figure 2. Fluorescence spectra of MSD (a) and MSD-Hg²⁺ (b) upon addition of different concentration of SG. [MSD] = 10 nM, [Hg²⁺] = 70 nM, [SG](from bottom to up) = 2.45, 12.25, 24.5, 36.75, 61.25 × 10⁻⁸ M, respectively. (c) Relationship between fluorescence intensity and the concentration of SG for SG-Hg²⁺-MSD and SG-MSD. Squares and triangles represent the fluorescence intensity of SG-MSD and SG-Hg²⁺-MSD, respectively; circles are the ratios of the fluorescence intensity SG-Hg²⁺-MSD to that of SG-MSD. 10 mM MOPS, 0.1 M NaNO₃, pH 7.50 was used as the buffer.





Figure 2. Cont.




We can see from the results that the fluorescence intensity of SG increases both in the absence of Hg^{2+} [Figure 2(a)] and in the presence of Hg^{2+} [Figure 2(b)]. The highest ratio of the fluorescence intensity of SG- Hg^{2+} -MSD to that of SG-MSD is 21.8 [Figure 2(c)], at which the concentration of SG is 1.225×10^{-7} M, so we selected the concentration of SG to be 1.225×10^{-7} M when the concentration of MSD is 10 nM and Hg^{2+} 70 nM.

3.3. Cys Sensor Sensitivity

To evaluate the sensitivity of the assay, different concentrations of Cys were mixed with MSD/ Hg^{2+} /SG, and then the fluorescence intensity was immediately measured. As shown in Figure 3(a), the fluorescence intensity of SG decreases gradually upon adding increased concentrations of Cys.

Figure 3. (a) Fluorescence spectra of SG-MSD- Hg^{2+} in the absence and presence of Cys, the concentration of Cys (from top to bottom): 0, 7, 14, 21, 28, 35, 42, 49, 56, 70, 84, 98, 112, 126, 140, 160, 200 nM. **(b)** The fluorescence intensity of SG-MSD- Hg^{2+} vs. [Cys]. [MSD] = 10 nM, [Hg^{2+}] = 70 nM, [SG] = 1.225×10^{-7} M. (b inset: the linear relationship of the fluorescence intensity of SG vs. [Cys], The error bars represent the standard deviation of three measurements).



When the concentration of Cys is two-fold higher than that of Hg^{2+} , the fluorescence intensity decreases very slowly due to formation of a 2:1 Cys/ Hg^{2+} adduct [31]. This implies that almost all the Hg^{2+} has been extracted from T- Hg^{2+} -T complex when the concentration of Cys is two fold higher than that of Hg^{2+} . When we increased the concentration of Cys further, we found that the fluorescence intensity decreased very slowly. Perhaps some Cys forms $\text{Hg}(\text{Cys})_3$ complexes. By measuring the fluorescence intensity at the emission maximum of SG- Hg^{2+} -MSD-Cys, a linear response of fluorescence intensity vs. [Cys] was observed in the range 7–84 nM [Figure 3(b) inset]. The detection limit may be estimated from Equation (1):

$$\text{LOD} = 3 \times \frac{S_0}{S} \quad (1)$$

where S_0 is the standard deviation of the blank and S is the sensitivity. 3.39 nM was the experimentally estimated detection limit for Cys, which was lower than most reported Cys sensors. Table 1 lists a comparison of methods for the determination of Cys and confirms the results.

Table 1. Comparison of methods for the determination of Cys.


Detection method	Linear range ($\mu\text{mol L}^{-1}$)	Detection limit ($\mu\text{mol L}^{-1}$)	Reference
Spectrophotometry	0.0082–0.12	0.0049	[9]
Spectrophotometry	0.17–50	Not given	[7]
Fluorimetry	100–5,000	Not given	[21]
Fluorimetry	0.05–4	0.025	[22]
Fluorimetry	0.025–6	0.02	[23]
Fluorimetry	0.01–0.8	0.0038	[24]
Fluorimetry	0.05–0.3	0.0096	[25]
Flow injection	0.001–0.5	0.0005	[18]
Flow injection	1–90	0.2	[19]
Flow injection	0.4–40	0.1	[20]
Capillary zone electrophoresis	0.1–100	0.058	[5]
Voltammetry	0.5–100	0.2	[12]
Voltammetry	2–10,000	1.0	[13]
Voltammetry	0.78–200	0.26 ± 0.01	[10]
Voltammetry	0.5–630	0.07	[11]
Fluorimetry	0.007–0.084 or 0.05–0.25 or 0.05–1	0.0034	This method

What is more important, our method is really fast. In the first step of this assay, SG staining was finished in 2 min, which was proved to be fully adequate for the SG binding by a previous study [30]. In the second step, after the addition of Cys, the fluorescence spectra was measured immediately, and as anticipated, the fluorescence intensity remained stable for a long time (data not shown), because Cys can bind to Hg^{2+} very strongly [31].

We further find that we can obtain different dynamic ranges by changing the concentration of MSD/ Hg^{2+} . For example, when we fixed the concentration of MSD to be 50 nM and that of Hg^{2+} to

be 350 nM (MSD kinetically forms the hairpin structure at $[\text{Hg}^{2+}]/7[\text{MSD}] = 1$ [30]), a linear dynamic range for Cys from 50 to 250 nM was obtained [Figure 4(a) inset]. When we changed the concentration of MSD and Hg^{2+} to be 100 nM and 700 nM, a very wide linear dynamic range for Cys from 50 to 1,000 nM was obtained [Figure 4(b) inset]. The concentration of SG was 6.125×10^{-7} M [Figure 4(a)] and 1.225×10^{-6} M [Figure 4(b)], respectively, which was determined by the same method mentioned above (Section 3.2). When the concentration of MSD/ Hg^{2+} changed, different concentration of Cys was needed to extract Hg^{2+} from T- Hg^{2+} -T structure due to the 2:1 Cys/ Hg^{2+} adduct, which leads to the different dynamic ranges for Cys.


Figure 4. The fluorescence intensity of SG-MSD- Hg^{2+} vs. [Cys] (a) [MSD] = 50 nM, $[\text{Hg}^{2+}] = 350$ nM, $[\text{SG}] = 6.125 \times 10^{-7}$ M, (b) [Cys] = 100 nM, $[\text{Hg}^{2+}] = 700$ nM, $[\text{SG}] = 1.225 \times 10^{-6}$ M (a and b inset: the linear relationship of the fluorescence intensity of SG vs. [Cys]).



3.4. Cys Sensor Selectivity

To challenge the assay's selectivity, other amino acids and the Cys-containing tripeptide glutathione (GSH) at a concentration of 140 nM were analyzed. The results are shown in Figure 5. It is clear that only Cys and GSH show significant fluorescence intensity changes. In contrast to the enormous fluorescence decrease observed for Cys and GSH, there is very little fluorescence change observed in the presence of other amino acids. A Cys-containing tripeptide GSH (γ -Glu-Cys-Gly) showed almost the same fluorescence decrease as Cys, indicating that Hg^{2+} can be extracted from the T- Hg^{2+} -T structure upon adding GSH. This is because GSH can also form a very stable 2:1 GSH/ Hg^{2+} complex with Hg^{2+} [34-36]. Hg^{2+} is well known to have an affinity to certain N-type ligands such as amino acids [37], but the amino acids, except for Cys, cannot extract Hg^{2+} from the thymine- Hg^{2+} -thymine complex. The binding affinity of Hg^{2+} to T-T mismatch sites thus appears to be stronger enough than that of Hg^{2+} to all of the amino acids studied, except Cys. We also found from Figure 5 that a sulfur-containing amino acid like methionine did not result in a significant fluorescence decrease. The formation constant for Cys to Hg^{2+} (about 10^{42}) is much higher than that of methionine to Hg^{2+} (about $10^{17.6}$) [31]. The method presented here therefore shows very high specificity to Cys.

Figure 5. The fluorescence intensity of the blank (SG-MSD-Hg²⁺, no amino acids) and solutions containing different amino acids (SG-MSD-Hg²⁺-amino acid). [MSD] = 10 nM, [Hg²⁺] = 70 nM, [SG] = 1.225 × 10⁻⁷ M, [amino acids] = [GSH] = 140 nM.



4. Conclusions

This paper describes a rapid, highly selective and sensitive fluorescence assay for Cys using mercury-specific DNA and SG. This assay is based on the extraction of Hg²⁺ by Cys from a T-Hg²⁺-T complex. The whole procedure can be done within 5 min. In addition, the assay is label free, low-cost and can provide variable linear dynamic ranges.

Acknowledgments

The authors would like to thank the Promotional Foundation for the Excellent Middle-Aged and Young Scientists of Shandong Province of China (BS2009SW040), the National Natural Science Foundation (20902096, 21104030) and a Project of Shandong Province Higher Educational Science and Technology Program (J11LB03, J10LF02), Shanghai Postdoctoral Sustentation Fund, China (11R21420800), General Administration of Quality Supervision, Inspection and Quarantine of PRC (2009QK098, 2010QK294).

References

1. Droge, W.; Holm, E. Role of cysteine and glutathione in HIV infection and other diseases associated with muscle wasting and immunological dysfunction. *FASEB J.* **1997**, *11*, 1077-1089.
2. Wang, X.F.; Cynader, M.S. Pyruvate released by astrocytes protects neurons from copper-catalyzed cysteine neurotoxicity. *J. Neurosci.* **2001**, *21*, 3322-3331.
3. Liu, J.; Yeo, H.C.; Overvik-Douki, E.; Hagen, T.; Doniger, S.J.; Chu, D.W.; Brook, G.A.; Ames, B.N. Chronically and acutely exercised rats: Biomarkers of oxidative stress and endogenous antioxidants. *J. Appl. Physiol.* **2000**, *89*, 21-28.
4. Lu, C.; Zu, Y.; Yam, V.W.W. Nonionic surfactant-capped gold nanoparticles as postcolumn reagents for high-performance liquid chromatography assay of low-molecular-mass biothiols. *J. Chromatogr. A* **2007**, *1163*, 328-332.

5. Jin, W.R.; Wang, Y. Determination of cysteine by capillary zone electrophoresis with end-column amperometric detection at a gold/mercury amalgam microelectrode without deoxygenation. *J. Chromatogr. A* **1997**, *769*, 307-314.
6. Nozal, M.J.; Bernal, J.L.; Toribio, L.; Marinero, P.; Moral, O.; Manzanar, L.; Rodriguez, E. Determination of glutathione, cysteine and N-acetylcysteine in rabbit eye tissues using high-performance liquid chromatography and post-column derivatization with 5,5'-dithiobis(2-nitrobenzoic acid). *J. Chromatogr. A* **1997**, *778*, 347-353.
7. Abu Eid, M. Spectrophotometric determination of cysteine and N-acetylcysteine in pharmaceutical preparations. *Microchim. Acta* **1998**, *129*, 91-95.
8. Wang, W.; Rusin, O.; Xu, X.; Kim, K.K.; Escobedo, J.O.; Fakayode, S.O.; Fletcher, K.A.; Lowry, M.; Schowalter, C.M.; Lawrence, C.M.; Fronczek, F.R.; Warner I.M.; Strongin, R.M. Detection of homocysteine and cysteine. *J. Am. Chem. Soc.* **2005**, *127*, 15949-15958.
9. Lunar, M.L.; Rubio, S.; Pérez-Bendito, D.; Carreto, M.L.; McLeod, C.W. Hexadecylpyridinium chloride micelles for the simultaneous kinetic determination of cysteine and cystine by their induction of the iodine-azide reaction. *Anal. Chim. Acta* **1997**, *337*, 341-349.
10. Deng, C.; Chen, J.; Chen, X.; Wang, M.; Nie, Z.; Yao, S. Electrochemical detection of l-cysteine using a boron-doped carbonnanotube-modified electrode. *Electrochim. Acta* **2009**, *54*, 3298-3302.
11. Bai, Y.H.; Xu, J.J.; Chen, H.Y. Selective sensing of cysteine on manganese dioxide nanowires and chitosan modified glassy carbon electrodes. *Biosens. Bioelectron.* **2009**, *24*, 2985-2990.
12. Shahrokhian, S.; Karimi, M. Voltammetric studies of a Cobalt(II)-4-methylsalophen modified carbon-paste electrode and its application for the simultaneous determination of cysteine and ascorbic acid. *Electrochim. Acta* **2004**, *50*, 77-84.
13. Amini, M.K.; Khorasani, J.H.; Khaloo, S.S.; Tangestaninejad, S. Cobalt(II) salophen-modified carbon-paste electrode for potentiometric and voltammetric determination of cysteine. *Anal. Biochem.* **2003**, *320*, 32-38.
14. Zhang, F.X.; Han, L.; Israel, L.B.; Daras, J.G.; Maye, M.M.; Ly, N.K.; Zhong, C.J. Colorimetric detection of thiol-containing amino acids using gold nanoparticles. *Analyst* **2002**, *127*, 462-465.
15. Li, L.; Li, B. Sensitive and selective detection of cysteine using gold nanoparticles as colorimetric probes. *Analyst* **2009**, *134*, 1361-1365.
16. Sudeep, P.K.; Joseph, S.T.S.; Thomas, K.G. Selective detection of cysteine and glutathione using gold nanorods. *J. Am. Chem. Soc.* **2005**, *127*, 6516-6517.
17. Jia, S.M.; Liu, X.F.; Li, P.; Kong, D.M.; Shen, H.X. G-quadruplex DNAzyme-based Hg²⁺ and cysteine sensors utilizing Hg²⁺-mediated oligonucleotide switching. *Biosens. Bioelectron.* **2011**, *27*, 148-152.
18. Waseem, A.; Yaqoob, M.; Nabi, A. Flow injection determination of cysteine in pharmaceuticals based on luminol-persulphate chemiluminescence detection. *Luminescence* **2008**, *23*, 144-149.
19. Teshima, N.; Nobuta, T.; Sakai, T. Simultaneous flow injection determination of ascorbic acid and cysteine using double flow cell. *Anal. Chim. Acta* **2001**, *438*, 21-29.
20. Zhao, C.; Zhang, J.C.; Song, J.F. Determination of l-Cysteine in amino acid mixture and human urine by flow-injection analysis with a biamperometric detector. *Anal. Biochem.* **2001**, *297*, 170-176.
21. Tanaka, F.; Mase, N.; Barbas, C.F., III. Determination of cysteine concentration by fluorescence increase: Reaction of cysteine with a fluorogenic aldehyde. *Chem. Commun.* **2004**, 1762-1763.

22. Shang, L.; Qin, C.J.; Wang, T.; Wang, M.; Wang, L.X.; Dong, S.J. Fluorescent conjugated polymer-stabilized gold nanoparticles for sensitive and selective detection of cysteine. *J. Phys. Chem. C* **2007**, *111*, 13414-13417.
23. Shang, L.; Dong, S.J. Sensitive detection of cysteine based on fluorescent silver clusters. *Biosens. Bioelectron.* **2009**, *24*, 1569-1573.
24. Huang, S.; Xiao, Q.; Li, R.; Guan, H.L.; Liu, J.; Liu, X.R.; He, Z.K.; Liu, Y. A simple and sensitive method for l-cysteine detection based on the fluorescence intensity increment of quantum dots. *Anal. Chim. Acta* **2009**, *645*, 73-78.
25. Ruan, Y.B.; Li, A.F.; Zhao, J.S.; Shen, J.S.; Jiang, Y.B. Specific Hg²⁺-mediated perylene bisimide aggregation for highly sensitive detection of cysteine. *Chem. Commun.* **2010**, *46*, 4938-4940.
26. Li, D.; Song, S.P.; Fan, C.H. Target-responsive structural switching for nucleic acid-based sensors. *Acc. Chem. Res.* **2010**, *43*, 631-641.
27. Zhang, J.; Wang, L.H.; Zhang, H.; Boey, F.; Song, S.P.; Fan, C.H. Aptamer-based multicolor fluorescent gold nanoprobe for multiplex detection in homogeneous solution. *Small* **2010**, *6*, 201-204.
28. Song, S.P.; Wang, L.H.; Li, J.; Zhao, J.; Fan, C.H. Aptamer-based biosensors. *Trends Anal. Chem.* **2008**, *27*, 108-117.
29. Zuo, X.L.; Song, S.P.; Zhang, J.; Pan, D.; Wang, L.H.; Fan, C.H. A target-responsive electrochemical aptamer switch (TREAS) for reagentless detection of nanomolar ATP. *J. Am. Chem. Soc.* **2007**, *129*, 1042-1043.
30. Wang, J.; Liu, B. Highly sensitive and selective detection of Hg²⁺ in aqueous solution with mercury-specific DNA and Sybr Green I. *Chem. Commun.* **2008**, 4759-4761.
31. Berthon, G. The stability constants of metal complexes of amino acids with polar side chains. *Pure Appl. Chem.* **1995**, *67*, 1117-1240.
32. Lee, J.S.; Ulmann, P.A.; Han, M.S.; Mirkin, C.A. A DNA-gold nanoparticle-based colorimetric competition assay for the detection of cysteine. *Nano Lett.* **2008**, *8*, 529-533.
33. Xu, H.; Hepel, M. "Molecular beacon"-based fluorescent assay for selective detection of glutathione and cysteine. *Anal. Chem.* **2011**, *83*, 813-819.
34. Oram, P.D.; Fang, X.; Fernando, Q. The formation of constants of mercury (II)-glutathione complexes. *Chem. Res. Toxicol.* **1996**, *9*, 709-712.
35. Fuhr, B.J.; Rabenstein, D.L. Nuclear magnetic resonance studies of the solution chemistry of metal complexes. IX. The binding of cadmium, zinc, lead, and mercury by glutathione. *J. Am. Chem. Soc.* **1973**, *95*, 6944-6950.
36. Stricks, W.; Kolthoff, I.M. Reactions between mercuric mercury and cysteine and glutathione. Apparent dissociation constants, heats and entropies of formation of various forms of mercuric mercapto-cysteine and -glutathione. *J. Am. Chem. Soc.* **1953**, *75*, 5673-5681.
37. Corradi, A.B.; Cramarossa, M.R.; Vezzosi, I.M.; Giusti, J.G. *trans*-2-Styrylbenzothiazole complexes with mercury(II) halides. Synthesis, characterization and X-ray crystal structure of diiodobis(*trans*-2-styrylbenzothiazole)mercury(II). *Polyhedron* **1993**, *12*, 2235-2239.