Pharmaceuticals **2012**, *5*, 745-757; doi:10.3390/ph5070745

OPEN ACCESS

pharmaceuticals

ISSN 1424-8247

www.mdpi.com/journal/pharmaceuticals

Article

Synthesis, Reactions and Evaluation of the Antimicrobial Activity of Some 4-(*p*-Halophenyl)-4*H*-naphthopyran, Pyranopyrimidine and Pyranotriazolopyrimidine Derivatives

Ashraf H. F. Abd El-Wahab

Chemistry Department, Faculty of Science, Al-Azhar University, Nasr City, Cairo 11884, Egypt; E-Mails: ash_abdelwahab@hotmail.com; ash_abdelwahab@yahoo.com; Tel: +20-10-0819-9893; Fax: +20-22-262-958

Received: 14 May 2012; in revised form: 15 June 2012 / Accepted: 21 June 2012 / Published: 12 July 2012

Abstract: A series of naphthopyran derivatives **3a–f** were prepared. Reaction of *2*-amino-4-(*p*-chlorophenyl)-7-methoxy-4*H*-naphtho[2,1-*b*]pyran-3-carbonitrile (**3b**) with Ac_2O afforded two products, 2-acetylamino-7-methoxy-4-(*p*-chlorophenyl)-4*H*-naphtho-[2,1-*b*]pyran-3-carbonitrile (**4**) and 10,11-dihydro-3-methoxy-9-methyl-12-(*p*-chloro-phenyl)-12*H*-naphtho[2,1-*b*]pyran[2,3-*d*]pyrimidine-11-one (**5**) and treatment of **3b** with benzoyl chloride gave the pyranopyrimidin-11-one derivative **6**. While treatment of **3b** with formamide afforded 11-amino-3-methoxy-12-(*p*-chlorophenyl)-12*H*-naphtho[2,1-*b*] pyrano[2,3-*d*]pyrimidine (**7**). Reaction of **3b** with triethyl orthoformate gave the corresponding 2-ethoxymethyleneamino-7-methoxy-4-(*p*-chlorophenyl)-4*H*-naphtho-[2,1-*b*] pyran-3-carbonitrile (**8**). Hydrazinolysis of **8** in EtOH at room temperature yielded 10-amino-10,11-dihydro-11-imino-3-methoxy-12-(*p*-chlorophenyl)-12*H*-naphtho[2,1-b]pyrano-[2,3-d]pyrimidine (**9**), while aminolysis of **8** with methylamine or dimethylamine gave the corresponding pyranopyrimidine and *N*,*N*-dimethylaminomethylene derivatives **10** and **11**. Condensation of **9** with some carboxylic acid derivatives afforded triazolopyrimidine derivatives **12–16**, while reaction of **9** with benzaldehyde gave 10-benzalamino-10,11-dihydro-11-imino-3-methoxy-12-(*p*-chlorophenyl)12*H*-naphtho[2,1-*b*]pyrano[2,3-*d*]pyrimidine (**17**). The structures of the newly synthesized compounds were confirmed by spectral data. The synthesized compounds were also screened for their antimicrobial activity.

Keywords: antimicrobial activity; arylidienemalonitrile; 6-methoxy-2-naphthol; naphthopyranopyrimidine; naphthopyranotriazolopyrimidine; carboxylic acid derivatives

---

1. Introduction

Pyran and fused 4*H*-pyran derivatives have attracted a great deal of interest owing to their antimicrobial activity [1–5], inhibition of influenza, virus sialidases [6], mutagenic activity [7], activity as antiviral [8] and antiproliferation agents [9], sex-pheromones [10], antitumor [11] and anti-inflammatory agents [12]. Moreover, pyran derivatives are well known for their antihistaminic activity [13]. Also pyrimdines are an important class of compounds and have widespread applications from pharmaceuticals to materials [14], with activities such as Tie-2 kinase inhibitors [15], HIV-1 inhibitor [16], anti-malarial [17], adenosine A1 receptor antagonism [18], anticancer [19], analgesic [20], cardiovascular [21] and antiallergic activities [22]. In view of the important biological properties of the pyran and pyrimidine derivatives as medicinal agents, we report here the synthesis and antimicrobial activities of new naphthopyrano, naphthopyranopyrimidine and naphthopyranotriazolopyrimidine derivatives.

2. Results and Discussion

Condensation of 6-methoxy-2-naphthol (**1**) with substituted 4-halobenzylidenmalononitriles **2a–c** and/or ethyl 4-halobenzylidenmalonates **2d–f** afforded the corresponding 2-amino-4-(*p*-halophenyl)-7-methoxy-4*H*-naphtho[2,1-*b*]pyran-3-carbonitriles **3a–c** and ethyl-2-amino-4-(*p*-halophenyl)-7-methoxy-4*H*-naphtho[2,1-*b*]pyran-3-carboxylates **3d–f**, respectively [23,24] (Scheme 1).

Scheme 1. Synthesis of naphthopyran derivatives **3a–f**.



a; Ar = *p*-Br-C_6H_4 X= CN
b; Ar = *p*-Cl-C_6H_4 X= CN
c; Ar = *p* F-C_6H_4 X= CN
d; Ar = *p* Br-C_6H_4 X= COOEt
e; Ar = *p* Cl-C_6H_4 X= COOEt
f; Ar = *p* F-C_6H_4 X= COOEt

The structure of compounds **3a–f** were established by spectral data. The IR spectrum of compounds **3a–f** showed absorptions at 3,466–3,350, 3,346–3,314 cm^{-1} (NH_2), 3,192–3,000 cm^{-1} (CH-aromatic), 2,950–2,900 (CH-aliphatic), 2,200–2,192 cm^{-1} (C≡N), 1,682–1,670 cm^{-1} (C=O). The ^{1}H-NMR of compounds **3a–f** showed chemical shifts δ_H at 3.75–3.78 (s, 3H, OCH_3), 5.27–5.530 (s, 1H, pyran CH), 6.98–7.21 (br, 2H, NH_2, exchangeable by D_2O) while the ^{13}C-NMR of compound **3b** showed δ_C at 56.5 (OCH_3), 28.4 (C-4), 117.1 (C≡N) and compound **3e** showed chemical shifts δ_C at 13.1 (CH_3-ester), 28.4 (C-4), 56.5 (OCH_3), 62.5 (CH_2-ester) and 172.2 (CO).

Treatment of **3b** with Ac_2O gave two products, depending on the reaction time; one product was identified as 2-acetylamino-7-methoxy-4-(*p*-chlorophenyl)-4*H*-naphtho[2,1-*b*]pyran-3-carbonitrile (**4**, 1/2 hour), while the other was identified as 10,11-dihydro-3-methoxy-9-methyl-12-(*p*-chlorophenyl)-12*H*-naphtho[2,1-*b*]pyrano-[2,3-*d*]pyrimidine-11-one (**5**, 3 hours). Support for structure **5** was obtained

by its independent synthesis by the reaction of **3e** with CH_3CN in the presence of dry HCl gas [25]. Reaction of **3b** with benzoyl chloride gave the pyranopyrimidin-11-one derivative **6**, while with formamide afforded 11-amino-3-methoxy-12-(*p*-chlorophenyl)-12*H*-naphtho[2,1-*b*]pyrano[2,3-d] pyrimidine **7** (Scheme 2).

Scheme 2. Synthesis of naphthopyran and naphthopyranpyrimidine derivatives **4–7**.



The structure of compounds **4–7** were established from their spectral data. The IR spectrum of compound **4** showed absorptions at 3,400 (NH), 2,202 cm^{-1} (CN), 1,612 cm^{-1} (C=O), while compound **5** showed absorptions at 3,464 (NH), 1,650 cm^{-1} (C=O) and compound **7** showed υ at 3,458, 3,380 (NH_2). The ^{1}H-NMR of compounds **4–7** showed chemical shifts δ_H at 3.78–3.82 (s, 3H, OCH_3), 5.66–5.70 (s, 1H, pyran CH). The mass spectra of compounds **6** and **7** provided additional evidence for the proposed structures.

Reaction of **3b** with triethyl orthoformate gave the corresponding 2-ethoxymetheneamino derivative **8**. Hydrazinolysis of **8** in EtOH at room temperature yielded 10-amino-10,11-dihydro-11-imino-3-methoxy-12-(*p*-chlorophenyl)-12*H*-naphtho[2,1-*b*]pyrano[2,3-*d*]pyrimidine (**9**). During treatment of **8** with phenylhydrazine, an nonisolable addition product **A** was formed first, followed by elimination of the nonisolable ethyl formatephenylhydrazone to give the enaminonitrile **3b**. Ammonolysis of **8** in MeOH at room temperature afforded compound **7** and aminolysis of **8** with methylamine and/or dimethylamine gave the corresponding 10-methyl-pyranopyrimidine and *N,N*-dimethylamino-methylene derivatives **10** and **11**, respectively (Scheme 3).

The structure of compounds **8–11** were established from their spectral data. The IR spectrum of compound **8** showed absorptions at 2,203 cm^{-1} (C≡N), while compound **9** showed υ at 3,316, 3,270 (NH_2), 3,209 cm^{-1} (NH) and compound **11** showed an absorption at 2,204 cm^{-1} (C≡N). The ^{1}H-NMR of compounds **8–11** showed chemical shifts δ_H at 3.77–3.80 (s, 3H, OCH_3) and 5.29–5.58 (s, 1H, pyran CH).

Scheme 3. Reaction of **3b** with triethyl orthoformate and of ammionium derivatives.



Reaction of **9** with formic acid or triethyl orthoformate, acetylchloride and benzoyl chloride afforded the corresponding triazolopyrimidine derivatives **12–14**, while cyclocondensation of **9** with ethyl cyanoacetate gave the corresponding 2-cyanomethyl derivative **15**. Treatment of **9** with ethyl chloroformate in dry benzene afforded traizolo-2-one derivative **16**. Reaction of **9** with benzaldehyde gave 10-benzalamino-10,11-dihydro-11-imino-3-methoxy-12-(*p*-chlorophenyl)-12*H*-naphtho-[2,1-*b*]-pyrano[2,3-*d*]pyrimidine (**17**) instead of the expected triazolopyrimidine derivative **14** (Scheme 4).

Scheme 4. Synthesis of naphthopyrantriazolopyridimide derivatives **12–17**.



The structure of compounds **12–17** were established from their spectral data. The IR spectrum of compound **15** showed an absorption at 2,200 cm^{-1} (C≡N), compound **16** showed absorptions at 3,200 cm^{-1} (NH) and 1,638 cm^{-1} (C=O), while compound **17** showed absorptions at 3,261 cm^{-1} (NH), and 1,621 cm^{-1} (C=N). The ^{1}H-NMR of compounds **12–17** showed chemical shifts δ_H at 3.78–3.80 (s, 3H, OCH_3), 526–5.66 (s, 1H, pyran CH), 8.63–9.64 (s, 1H, pyrimidine CH).

The antimicrobial activity of the newly synthesized compounds **3–17** was evaluated against the bacterial strains *Staphylococcus aureus (NCTC-7447)*, *Bacillus cereus (ATCC-14579)*, *Escherichia coli (NCTC-10410)*, *Serratia marcescens (IMRU-70)* and the fungal strains *Aspergillus fumigatus (MTCC-3008)*, and *Candida albicans* (MTCC-227) by the disk diffusion method [26,27]. Ampicillin and ketoconazole were used as standard drugs for the bacteria and fungi, respectively. Preliminary screening of the naphthopyran derivatives and standard drugs was performed at fixed concentrations of 500 μg/mL. Inhibition was recorded by measuring the diameter of the inhibition zone at the end of 24 hours for bacteria and 72 hours for fungi. Each experiment was repeated twice. Based on the results of zone of inhibition, the minimum inhibitory concentration (MIC) of compounds **3–17** against all bacterial and fungal strains was determined by the liquid dilution method. Stock solutions of the tested compounds with 500, 250, 200, 100, 50, 25, 12.5, and 6.25 μg/mL concentrations were prepared with DMSO as solvent. The solutions of standard drugs, ampicillin and ketoconazole, were prepared in the same concentrations. The minimum inhibitory concentration at which no growth was observed was taken as the MIC value (Table 1). The comparison of the MICs (μg/mL) of potent compounds and standard drugs against tested strains are presented in Table 1. Investigation of the antibacterial screening data showed that some of the compounds were active against all four pathogenic bacteria. Ethyl 2-amino-4-(*p*-chlorophenyl)naphthopyrane-3-carboxylate (**3e**), triazolopyrimidine derivative **12**, 2-methyl-triazolopyrimidine derivative **13**, 2-phenyl-triazolopyrimidine derivative **14** and, 2-oxa-triazolopyrimidine derivative **16** exhibited good activity against *S. aureus*. Similarly 2-amino-3-cyano-4-(*p*-flourophenyl)naphthopyran (**3c**), 2-methyltriazolopyrimidine derivative **14**, triazolopyrimidine derivative **12**, triazolopyrimidine 2-ethanenitrile derivative **15** and 2-oxa-triazolopyrimidine derivative **16** exhibited good activity against *B. cereus* and 2-amino-3-cyano4-(*p*-bromophenyl)naphthopyran (**3a**), 2-amino-3-cyano-4-(*p*-chlorophenyl)naphthopyran (**3b**), 2-amino-3-cyano-4-(*p*-flourophenyl) naphthopyran **3c**, ethyl 2-amino-4-(*p*-chlorophenyl)naphthopyran-3-carboxylate (**3e**), 2-methyl-triazolopyrimidine derivative **13** and triazolopyrimidine 2-ethanenitrile derivative **16** exhibited good activity against *E. coli*, while 2-amino-3-cyano-4-(*p*-bromo/chloro/fluorophenyl)naphthopyran derivatives **3a–c**, triazolopyrimidine derivative **12**, 2-phenyltriazolopyrimidine derivative **14** and 2-oxatriazolopyrimidine derivative **16** exhibited good activity against *S. marcescens*. Aminoimino derivative **9** was inactive against *S. aureus*, while the pyranpyrimidin-11-one **6** and 10-benzalamino-pyranopyrimidine derivative **17** was inactive against *B. cereus*, the compounds ethyl 2-amino-4-(*p*-chlorophenyl)naphthopyrane-3-carboxylate (**3e**), 11-amino-pyranopyrimidine **7** and 2-ethoxy-methyleneamino derivative **8** was inactive against *E. coli*, and the 2-acetylamino-pyranopyrimidine compound **4** was inactive against *S. marcescens*. The remaining compounds showed moderate to weak antibacterial activity.

Table 1. Antimicrobial activity of the new compounds.

Compounds	Minimum inhibitory concentration (MIC) in µg/mL					
	Bacterial strains				Fungal strains	
	S. aureus	***B. cereus***	***E. coli***	***S. marcescens***	***A. fumigatus***	***C. albicans***
3a	100	100	25	50	100	100
3b	125	200	50	50	125	100
3c	100	50	25	50	50	50
3d	250	200	500	500	250	100
3e	25	100	50	100	25	50
3f	500	500	-	125	250	500
4	500	250	250	100	500	-
5	100	125	-	500	-	250
6	100	-	100	250	500	500
7	500	200	-	200	-	250
8	250	200	500	500	250	100
9	-	500	-	100	500	500
10	500	250	100	250	250	500
11	-	500	250	-	-	500
12	25	25	50	25	100	50
13	25	50	25	50	50	500
14	50	100	100	25	25	50
15	100	25	25	125	100	500
16	50	25	50	50	125	100
17	500	-	250	500	-	-
Ampcillin	6.25	6.25	6.25	6.25	6.25	6.25
Ketoconazole	-	-	-	-	31.25	31.25

The antifungal results (Table 1) revealed that the synthesized compounds showed variable degrees of inhibition against the tested fungi. The compounds 2-amino-3-cyano-4-(*p*-fluorophenyl)naphthopyran (**3c**), ethyl 2-amino-4-(*p*-chlorophenyl)naphthopyran-3-carboxylate (**3e**), and 2-phenyltriazolopyrimidine **14** possessed good antifungal activity against *A. fumigatus* and *C. albicans*, while the compounds 9-methyl-pyranopyrimidine **5**, 11-amino-pyranopyrimidine **7**, *N*,*N*-dimethylaminomethylene derivative **11** and 10-benzalaminopyranopyrimidine **17** were inactive against *A. fumigates*, and the 2-acetylaminopyranopyrimidine **4** and 10-benzalaminopyranopyrimidine **17** were inactive against *C. albicans.* The remaining compounds showed moderate to weak antifungal activity.

3. Experimental

3.1. General

Melting points were determined on a Stuart melting point apparatus and are uncorrected. IR spectra (ν, cm^{-1}) were recorded in KBr using a FT-IR 5300 spectrometer and aPerkin Elmer spectrum RXIFT-IR system. The ^{1}H-NMR at (300 MHz) and ^{13}C-NMR spectra (75 MHz) were recorded in DMSO-d_6 on a Varian Mercury VX-300 NMR spectrometer. Chemical shifts (δ) are referred to that of the solvent.

Mass spectra were measured on a Shimadzu GMMS-QP-1000 EX mass spectrometer at 70 eV. The elemental analyses were performed at the Micro Analytical Center, Cairo University, Egypt.

3.2. General Procedure: Synthesis of 4H-Pyran Derivatives **3a–f**

A mixture of substituted 4-halobenzylidenmalononitriles **2a–c** (10 mmol) and/or ethyl 4-halobenzylidenmalonates **2d–f** (10 mmol), 6-methoxy-2-naphthol (**1**) (0.17 g, 10 mmol) and piperidine (0.5 mL) in absolute EtOH (50 mL) was heated until precipitation was completed. The precipitate was collected by filtration and recrystallized from dioxane and EtOH/benzene respectively.

2-Amino-4-(p-bromophenyl)-7-methoxy-4H-naphtho[2,1-b]pyrane-3-carbonnitrile (**3a**). White crystals (dioxane); yield 88%, mp 260–262 °C; IR: 3,454, 3,315 (NH_2), 3,182 (CH-aromatic), 2,950 (CH-aliphatic), 2,192 (C≡N), 1,658 (C=C). ^{1}H-NMR δ: 3.78 (s, 3H, OCH_3), 5.27 (s, 1H, pyran CH), 6.98 (br, 2H, NH_2, exchangeable by D_2O), 7.06–7.79 (m, 9H, Ar-H). Anal. Calcd. for $C_{21}H_{15}BrN_2O_2$ (406.03): C, 61.90; H, 3.90; N, 6.84%. Found: C, 61.93; H, 3.71; N, 6.88%.

2-Amino-4-(p-chlorophenyl)-7-methoxy-4H-naphtho[2,1-b]pyrane-3-carbonnitrile (**3b**). White crystals (dioxane); yield 90%, mp 246–248 °C; IR: 3,358, 3,314 (NH_2), 3,186 (CH-aromatic), 2,932 (CH-aliphatic), 2,198 (C≡N), 1,660 (C=C). ^{1}H-NMR δ: 3.77 (s, 3H, OCH_3), 5.28 (s, 1H, pyran CH), 6.99 (br, 2H, NH_2, exchangeable by D_2O), 7.06–7.79 (m, 9H, Ar-H). ^{13}C-NMR δ: 28.4 (C-4), 56.50 (OCH_3), 59.1 (C-3), 105.9 (C-7), 117.1 (C≡N), 118.6 (C-5), 118.9 (C-9), 121.3 (C-4a), 123.6 (C-10), 127.1 (C-6), 128.2 (C-5a, C-8a), 128.7, 129.7, 131.3, 136.6, 143.8 (Ar), 150.2 (C-10), 157.2 (C-8), 160.2 (C-2). Anal. Calcd. for $C_{21}H_{15}ClN_2O_2$ (362.08): C, 69.52; H, 4.17; N, 7.72%. Found: C, 69.50; H, 4.15; N, 7.70%.

2-Amino-4-(p-flouroophenyl)-7-methoxy-4H-naphtho[2,1-b]pyran-3-carbonitrile (**3c**). White crystals (dioxane); yield 86%, mp 255–257 °C; IR: 3,466, 3,318 (NH_2), 3,192 (CH-aromatic), 2,900 (CH-aliphatic), 2,200 (C≡N), 1,662 (C=C). ^{1}H-NMR δ: 3.75 (s, 3H, OCH_3), 5.30 (s, 1H, pyran CH), 7.01 (br, 2H, NH_2, exchangeable by D_2O), 7.08–8.31 (m, 9H, Ar-H). Anal. Calcd for $C_{21}H_{15}FN_2O_2$ (346.11): C, 72.82; H, 4.37; N, 8.09%. Found: C, 72.80; H, 4.35; N, 7.99%.

Ethyl 2-amino-4-(p-bromophenyl)-7-methoxy-4H-naphtho[2,1-b]pyran-3-carboxylate (**3d**). Colourless needle-like crystals (ethanol/benzene); yield 79%, mp 167–169 °C: IR: 3,350, 3,324 (NH_2), 3,000 (CH-aromatic), 2,944 (CH-aliphatic), 1,682 (C=O), 1,618 (C=C). ^{1}H-NMR δ: 1.21 (t, 3H, CH_3, J = 7.1 Hz), 3.78 (s, 3H, OCH_3), 4.04 (q, 2H, CH_2, J = 7.1 Hz), 5.42 (s, 1H, pyran CH), 7.21(br, 2H, NH_2, exchangeable by D_2O), 7.28–7.86 (m, 9H, Ar-H). Anal. Calcd. for $C_{23}H_{20}BrNO_4$ (453.06): C, 60.81; H, 4.44; N, 3.08%. Found: C, 60.80; H, 4.41; N, 3.03%.

Ethyl 2-amino-4-(p-chlorophenyl)-7-methoxy-4H-naphtho[2,1-b]pyran-3-carboxylate (**3e**). Colourless needle crystals, (ethanol/benzene); yield 82%, mp 172–174 °C: IR: 3,458, 3,324 (NH_2), 3,010 (CH-aromatic), 2,970 (CH-aliphatic), 1,670 (C=O), 1,622 (C=C). ^{1}H-NMR δ: 1.38 (t, 3H, CH_3, J = 7.1 Hz), 3.80 (s, 3H, OCH_3), 4.31 (q, 2H, CH_2, J = 7.1 Hz), 5.40 (s, 1H, pyran CH), 6.24 (br, 2H, NH_2, exchangeable by D_2O), 6.89–7.75 (m, 9H, Ar-H). ^{13}C-NMR δ: 13.1 (CH_3-ester), 28.4 (C-4), 56.5 (OCH_3), 59.1 (C-3), 62.5 (CH_2-ester), 105.9 (C-7), 118.6 (C-5), 118.9 (C-9), 121.3 (C-4a), 123.6

(C-10), 127.1 (C-6), 128.2 (C-5a, C-8a), 128.7, 129.7, 131.3, 136.6, 143.8 (Ar),150.2 (C-10), 157.2 (C-8), 160.2 (C-2), 172 (C=O). Anal. Calcd. for $C_{23}H_{20}ClNO_4$ (409.11): C, 67.40; H, 4.92; N, 3.42%. Found: C, 67.38; H, 4.90; N, 3.39%.

Ethyl 2-amino-4-(p-flouophenyl)-7-methoxy-4H-naphtho[2,1-b]pyran-3-carboxylate (**3f**). Colourless needle-like crystals (ethanol/benzene); yield 77%, mp 186–188 °C; IR: 3,422, 3,346 (NH_2), 3,192 (CH-aromatic), 2,900 (CH-aliphatic), 1,682 (CO), 1,600 (C=C). ^{1}H-NMR δ: 1.39 (t, 3H, CH_3, J = 7.1 Hz), 3.78 (s, 3H, OCH_3), 4.30 (q, 2H, CH_2, J = 7.1 Hz), 5.54 (s, 1H, pyran CH), 6.36 (br, 2H, NH_2, exchangeable by D_2O), 7.05–8.06 (m, 9H, Ar-H). Anal. Calcd. for $C_{23}H_{20}FNO_4$ (393.14): C, 70.20; H, 5.10; N, 3.52%. Found: C, 70.22; H, 5.12; N, 3.56%.

Synthesis of 2-acetylamino-7-methoxy-4-(p-chlorophenyl)-4H-naphtho[2,1-b]pyran-3-carbonitrile (**4**). A solution of **3b** (0.36 g, 10 mmol) in Ac_2O (20 mL) was heated under reflux for 30 min. The solid product formed was filtered off and washed with cold EtOH, The solid obtained was filtered off and recrystallized from EtOH. Pale yellow crystals, yield 89%, mp 175–177 °C; IR: 3,400 (NH), 3,122 (CH-aromatic), 2,940 (CH-aliphatic), 2,202 (C≡N), 1,612 (C=O). ^{1}H-NMR δ: 2.26 (s, 3H, CH_3), 3.82 (s, 3H, OCH_3), 5.70 (s, 1H, pyran CH), 7.22–7.83 (m, 9H, Ar-H), 12.49 (br, 1H, NH, exchangeable by D_2O). Anal. Calcd. for $C_{23}H_{17}ClN_2O_3$ (404.09): C, 68.23; H, 4.23; N, 6.92%. Found: C, 68.20; H, 4.19; N, 6.90%.

Synthesis of 10,11-dihydro-3-methoxy-9-methyl-12-(p-chlorophenyl)-12H-naphtho-[2,1-b]pyrano[2,3-d]pyrimidine-11-one (**5**). Method A: A solution of **3b** (0.36 g, 10 mmol) in Ac_2O (20 mL) was heated under reflux for 3 hours. The precipitate was filtered off, washed with cold EtOH. The solid obtained was filtered off and recrystallized from DMF; Method B: Gaseous dry HCl was bubbled through the mixture of **3e** (0.40 g, 10 mmol) and CH_3CN (30 mL) for 4–6 hours. The reaction mixture was poured into ice water and made alkaline with 10% aqueous ammonium hydroxide to give **5**. White crystals, yield 85%, mp 290–292 °C: IR: 3,464 (NH), 3,001 (CH-aromatic), 2,980 (CH-aliphatic), 1,650 (C=O), 1,620 (C=C). ^{1}H-NMR δ: 2.28 (s, 3H, CH_3), 3.78 (s, 3H, OCH_3), 5.66 (s, 1H, pyran CH), 6.89–8.01 (m, 10H, Ar-H and NH). Anal. Calcd. for $C_{23}H_{17}ClN_2O_3$ (404.09): C, 68.23; H, 4.23; N, 6.92%. Found: C, 68.20; H, 4.19; N, 6.90%.

Synthesis of 10,11-dihydro-3-methoxy-9-phenyl-12-(p-chlorophenyl)-12H-naphtho[2,1-b]-pyrano[2,3-d]pyrimidine-11-one (**6**). A solution of **3b** (0.36 g, 10 mmol) in benzoyl chloride (20 mL) was heated under reflux for 6 hours. The excess of benzoyl chloride was removed under reduced pressure and the residue was poured into cold water. The precipitate was collected by filtration, washed with CCl_4 (10 mL) to remove the formed benzoic acid and the residue was dried. The solid obtained was filtered off and recrystallized from DMF. Yellow crystals, yield 80%, mp > 360 °C; IR: 3,433 (NH), 3,012 (CH-aromatic), 2,892 (CH-aliphatic), 1,640 (C=O), 1,572 (C=C). MS *m/z* (%) = 466 (M+, 47.7), 326 (100), 250 (16.4), 129 (10.9). Anal. Calcd. for $C_{28}H_{19}ClN_2O_3$ (466.11): C, 72.03; H, 4.10; N, 6.01%. Found: C, 72.01; H, 4.02; N, 5.88%.

*Synthesis of 11-amino-3-methoxy-12-(p-*chlorophenyl*))-12H-naphtho[2,1-b]pyrano[2,3-d]pyrimidine* (**7**). Method A: A solution of **3b** (0.36 g, 10 mmol) in formamide (20 mL) was heated under reflux for

6 hours. The solid obtained was filtered off and recrystallized from benzene; Method B: Gaseous NH_3 was bubbled through **8** (0.41 g, 10 mmol) in MeOH for 1 hour. The solid formed was collected to give **7**. White crystals (benzene); yield 75%, mp 317–319 °C; IR: 3,458, 3,380 (NH_2), 3,174 (CH-aromatic), 2,901 (CH-aliphatic), 1,658 (C=C). MS *m/z* (%) = 389 (M+, 25.6), 249 (100), 223 (10.1), 181 (1.1). Anal. Calcd. for $C_{22}H_{16}ClN_3O_3$ (389.09): C, 67.78; H, 4.14; N, 10.78%. Found: C, 67.72; H, 4.10; N, 10.74%.

Synthesis of 2-ethoxymethyleneamino-7-methoxy-4-(p-chlorophenyl)-4H-naphtho-[2,1-b]pyrane-3-carbonitrile (**8**). A mixture of **3b** (0.36 g, 10 mmol) and triethyl orthoformate (2 mL) in acetic anhydride (10 mL) was refluxed for 2 hours. After cooling, the precipitated product was filtered off and washed several times with cold EtOH. The solid obtained was filtered off and recrystallized from benzene. Colourless crystals, yield 77%, mp 211–213 °C; IR: 2,980 (CH-aromatic), 2,835 (CH-aliphatic), 2,204 (CN), 1,612 (C=N). ^{1}H-NMR δ: 1.27 (t, 3H, CH_3, J = 7.1 Hz), 3.78 (s, 3H, OCH_3), 4.40 (q, 2H, CH_2, J = 7.1 Hz), 5.58 (s, 1H, pyran CH), 7.24–785 (m, 9H, Ar-H), 8.67 (s, 1H, N = CH). Anal. Calcd. for $C_{24}H_{19}ClN_2O_3$ (418.11): C, 68.82; H, 4.57; N, 6.69%. Found: C, 68.80; H, 4.51; N, 6.62%.

3.3. General Procedure: Synthesis of Pyranopyrimidine Derivatives **9** *and* **10**

A mixture of **8** (0.41 g, 10 mmol), hydrazine hydrate (5 mL, 99%) or methylamine (10 mmol) in absolute ethanol (50 mL) was stirred for 1 hour at room temperature. The solid obtained was filtered off and recrystallized from dioxane.

10-Amino-10,11-dihydro-11-imino-3-methoxy-12-(p-chlorophenyl)-12H-naphtho[2,1-b]pyrano[2,3-d] pyrimidine (**9**). White crystals, yield 81%, mp 256–258 °C; IR: 3,316, 3,270 (NH_2), 3,209 (NH), 2,936 (CH-aromatic), 2,899 (CH-aliphatic), 1,647 (C=N). ^{1}H-NMR δ: 3.80 (s, 3H, OCH_3), 5.66 (s, 1H, pyran CH), 5.87 (br, 2H, NH_2, exchangeable by D_2O), 7.15–7.79 (m, 10H, Ar-H and NH), 8.04 (s, 1H, pyrimidine CH). Anal. Calcd. for $C_{22}H_{17}ClN_4O_2$ (404.10): C, 65.27; H, 4.23; N, 13.84%. Found: C, 65.25; H, 4.20; N, 13.82%.

10,11-Dihydro-11-imino-3-methoxy-10-methyl-12-(p-chlorophenyl)-12H-naphtho[2,1-b]pyrano[2,3-d] pyrimidine (**10**). White crystals, yield 80%, mp 255–257 °C; IR: 3,376 (NH), 3,006 (CH-aromatic), 2,980, 2,830 (CH-aliphatic), 1,620 (C=N). ^{1}H-NMR δ: 3.32 (s, 3H, N-CH_3), 3.77 (s, 3H, OCH_3), 5.29 (s, 1H, pyran CH), 6.98–7.80 (m, 10H, Ar-H and NH), 8.01 (s, 1H, pyrimidine CH). Anal. Calcd for $C_{23}H_{18}ClN_3O_2$ (403.11): C, 68.40; H, 4.49; N, 10.40%. Found: C, 68.20; H, 4.41; N, 10.38%.

Synthesis of 7-methoxy-2-(N,N-dimethylaminomethylene)-4-(p-chlorophenyl)-4H-naphtho[2,1-b]-pyrane-3-carbonitrile (**11**). A mixture of **8** (0.41 g, 10 mmol) and dimethylamine (5 mL) in ethanol was stirred for 1 hour. The white solid formed was filtered, washed with cold EtOH and recrystallized from benzene. White crystals, yield 79%, mp 218–220 °C; IR: 2,924 (CH-aliphatic), 2,190 (C≡N), 1,616 (C=N). ^{1}H-NMR δ: 2.97 (s, 3H, N-CH_3), 3.13 (s, 3H, N-CH_3), 3.78 (s, 3H, OCH_3), 5.40 (s, 1H, pyran CH), 7.18–7.83 (m, 9H, Ar-H), 8.42 (s, 1H, N=CH). Anal. Calcd. for $C_{24}H_{20}ClN_3O_2$ (417.12): C, 68.98; H, 4.82; N, 10.06%. Found: C, 68.81; H, 4.66; N, 9.98%.

3.4. General Procedure: Synthesis of Pyranotriazolopyrimidine Derivatives **12–16**

A mixture of **9** (0.40 g, 10 mmol), triethyl orthoformate, formic acid, acetyl chloride or benzoyl chloride (0.01 mol) in dry benzene (20 mL), was refluxed for 3 hours. The solid obtained was filtered off and recrystallized from dioxane.

11-Methoxy-14-(p-chlorophenyl)-14H-naphtho[2,1-b]pyrano[3,2-b][1,2,4]triazolo[1,5-c]pyrimidine (**12**). White crystals, yield 80%, mp 260–262 °C; IR: 3,064 (CH-aromatic), 2,984, 2,844 (CH-aliphatic), 1,602 (C=C). ^{1}H-NMR δ: 3.78 (s, 3H, OCH_3), 5.42 (s, 1H, pyran CH), 7.00–7.91 (m, 9H, Ar-H), 8.63 (s, 1H, pyrimidine CH), 9.51 (s, 1H, triazolo CH). Anal. Calcd. for $C_{23}H_{15}ClN_4O_2$(414.09): C, 66.59; H, 3.61; N, 13.50%. Found: C, 66.50; H, 3.53; N, 13.10%.

2-Methyl-11-methoxy-14-(p-chlorophenyl)-14H-naphtho[2,1-b]pyrano[3,2-e][1,2,4]triazolo[1,5-c] pyrimidine (**13**). White crystals, yield 85%, mp 283–285 °C; IR: 3,074 (CH-aromatic), 2,936, (CH-aliphatic), 1,622 (C=C). ^{1}H-NMR δ: 2.40 (s, 3H, CH_3), 3.79 (s, 3H, OCH_3), 5.43 (s, 1H, pyran CH), 6.99–7.99 (m, 9H, Ar-H), 9.51 (s, 1H, pyrimidine CH). Anal. Calcd. for $C_{24}H_{17}ClN_4O_2$ (428.10): C, 67.21; H, 3.96; N, 13.06%. Found: C, 67.10; H, 3.66; N, 12.99%.

2-Phenyl-11-methoxy-14-(p-chlorophenyl)-14H-naphtho[2,1-b]pyrano[3,2-e][1,2,4]triazolo[1,5-c] pyrimidine (**14**). Pale yellow crystals, yield 70%, mp 281–283 °C; IR: 3,058 (CH-aromatic), 2,920 (CH-aliphatic), 1,626 (C=C). ^{1}H-NMR δ: 3.78 (s, 3H, OCH_3), 5.32 (s, 1H, pyran CH), 7.03–7.91 (m, 14H, Ar-H), 8.70 (s, 1H, pyrimidine CH). Anal. Calcd for $C_{29}H_{19}ClN_4O_2$ (490.12): C, 70.95; H, 3.87; N, 11.41%. Found: C, 70.65; H, 3.66; N, 11.27%.

Synthesis of 11-methoxy-14-(p-chlorophenyl)-14H-naphtho[2,1-b]pyrano[3,2-e][1,2,4]triazolo[1,5-c]pyrimidine-2-ethanenitrile (**15**). A mixture of **9** (0.40 g, 10 mmol) with ethyl cyanoacetate (10 mmol) in absolute ethanol (30 mL), was refluxed for 3 hours. The solid obtained was filtered off and recrystallized from dioxane. White crystals, yield 70%, mp 291–293 °C; IR: 2,934 (CH-aliphatic), 2,200 (C≡N), 1,604 (C=C). ^{1}H-NMR δ: 3.80 (s, 3H, OCH_3), 4.50 (s, 2H, CH_2), 5.40 (s, 1H, pyran CH), 6.99–7.94 (m, 9H, Ar-H), 9.64 (s, 1H, pyrimidine CH). Anal. Calcd. for $C_{25}H_{16}ClN_5O_2$ (453.10): C, 66.16; H, 3.52; N, 15.42%. Found: C, 66.01; H, 3.41; N, 15.23%.

Synthesis of 11-methoxy-14-(p-chlorophenyl)-2oxa-2H,3H,14H-naphtho[2,1-b]pyrano[3,2-e][1,2,4]-triazolo[1,5-c]pyrimidine (**16**). A mixture of **9** (0.40 g, 10 mmol) with ethyl chloroformate (10 mmol) in dry benzene (30 mL) was refluxed for 1 hour. The solid obtained was filtered off and recrystallized from dioxane. White crystals, yield 76%, mp 310–312 °C: IR: 3,200 (NH), 2,988, 2,930 (CH-aliphatic), 1,638 (C=O), ^{1}H-NMR δ: 3.80 (s, 3H, OCH_3), 5.28 (s, 1H, pyran CH), 6.80–7.88 (m, 10H, Ar-H and NH), 9.64 (s, 1H, pyrimidine CH). Anal. Calcd. for $C_{23}H_{15}ClN_4O_3$ (431.08): C, 64.12; H, 3.50; N, 13.00%. Found: C, 63.86; H, 3.45; N, 12.81%.

Synthesis of 10-benzalamino-10,11-dihydro-11-imino-3-methoxy-12-(p-chlorophenyl)12H-naphtho-[2,1-b]pyrano-[2,3-d]pyrimidine (**17**). A mixture of **9** (0.40 g, 1.0 mmol), with benzaldehyde (10 mmol), piperidine (0.5 mL) and dioxane (30 mL) was refluxed for 6 hours. The precipitate was filtered off and washed several times with cold EtOH. The solid was recrystallized from dioxane.

White crystals, yield 71%, mp 255–257 °C; IR: 3,261 (NH), 2,988, 2,930 (CH-aliphatic), 1,652 (C=N), ^{1}H-NMR δ: 3.79 (s, 3H, OCH_3), 5.66 (s, 1H, pyran CH), 6.80–7.88 (m, 15H, Ar-H and NH), 8.27 (s, 1H, N=CH), 9.64 (s, 1H, pyrimidine CH). Anal. Calcd. for $C_{29}H_{21}ClN_4O_2$(492.14): C, 70.66; H, 4.29; N, 11.37%. Found: C, 70.54; H, 4.04; N, 11.21%.

3.5. Antimicrobial Assay

Inoculums of the bacterial and fungal culture were prepared. To a series of tubes containing 1 mL each of naphthopyran compound solution with different concentrations and 0.2 mL of the inoculums was added. A further 3.8 mL of sterile water was added to each of the test tubes. These test tubes were incubated for 24 hours at 37 °C and observed for the presence of turbidity. This method was repeated by changing naphthopyran compounds for the standard drugs ampicillin and ketoconazole for comparison.

4. Conclusions

Our interest in the synthesis of such compounds was to focus on their study as antimicrobial agents as a part of our program which aimed at the development of new heterocyclic compounds as more potent antimicrobial agents. In this paper we revealed the synthesis of some new naphthopyran, naphthopyranopyrimidine and naphthopyranotriazolopyrimidine derivatives and the antimicrobial evaluation of all the novel compounds. The structures of these compounds were elucidated on the basis of IR, ^{1}H-NMR, ^{13}C-NMR and MS data. Evaluation of the new compounds established that **3a–e**, **12–14** and **16** showed improved antimicrobial activity.

References and Notes

1. Eid, F.A.; Abd El-Wahab, A.H.F.; El-Hagali, G.A.M.; Khafagy, M.M. Syhthesis and antimicrobial evaluation of naphtho[2,1-*b*]pyrano[2,3-*d*]pyrimidine and pyrano[3,2-*e*][1,2,4] triazolo[1,5-*c*]pyrimidine derivative. *Acta Pharm.* **2004**, *54*, 13–26.
2. Abd El-Wahab, A.H.F. Synthesis of some new pyrano[2,3-*d*][1,2,4]triazolo[1,5-*c*]pyrimidine and pyrimido[1,6-*b*]triazine derivatives. *Acta Pharm*. **2003**, *58*, 701–720.
3. El-Agrody, A.M.; Abd El-Latif, M.S.; El-Hady, N.A.; Fakery, A.H.; Bedair, A.H. Heteroaromatization with 4-Hyroxycoumarin. Part II: Synthesis of some new pyrano[2,3-*d*] [1,2,4]triazolo[1,5-*c*]pyrimidine and pyrimido[1,6-*b*]triazine derivatives. *Molecules* **2001**, *6*, 519–527.
4. Bedair, A.H.; El-Haddy, N.A.; Abd El-Latif, M.S.; Fakery, A.H.; El-Agrody, A.M. 4-Hyroxycoumarin in heterocycloic synthesis. Part III: Synthesis of some new pyrano[2,3-*d*] pyrimidine, 2-substituted[1,2,4]triazolo[1,5-*c*]pyrimidine and pyrimido[1,6-*b*]triazine derivatives. *Farmaco* **2000**, *55*, 708–714.
5. El-Agrody, A.M.; El-Hakim, M.H.; Abd El-Latif, M.S.; Fakery, A.H.; El-Sayed, E.M.; El-Ghareab, K.A. Synthesis pyrano[2,3-*d*]pyrimidine and pyrano[3,2-*e*][1,2,4]triazolo[2,3-*c*] pyrimidine derivatives with promising antibacterial activities. *Acta Pharm*. **2000**, *50*, 111–120.

6. Taylor, R.N.; Cleasby, A.; Singh, O.; Skarzynski, T. Dihydropyrancarboxamide related to zanamivir, a new series of inhibitors of influrnza virus sialidase. 2-Crystallographic and molecular modeling study of complex 4-amino-4*H*-pyran-6-carboxamides and sialidase from influenza virus types. *J. Med. Chem*. **1998**, *41*, 798–807.
7. Hirmoto, K.; Nasuhara, A.; Michiloshi, K.; Kato, T.; Kikugawa, K. DNA strand-breaking activity and mutagenicity of 2,3-dihydro-3,5-dihydroxy-6-methyl-4*H*-pyran-4-one (DDMP). A Maillard reaction product of glucose and glycine. *Mutat. Res*. **1997**, *395*, 47–56.
8. Martinez, A.G.; Marco, L.J. Friedlander reaction on 2-amino-3-cyano-4*H*-pyrans, synthesis of derivatives of 4*H*-pyran[2,3-*b*]quinoline, new tacrine analogues. *Bioorg. Med. Chem. Lett.* **1997**, *7*, 3165–3170.
9. Dell, C.P.; Smith, C.W. Antiproliferative derivatives of 4*H*-naphtho[1,2-b]pyran and process for their preparation. EP537949, 21 April 1993.
10. Bianchi, G.; Tava, A. Synthesis of (2R)(+)-2,3-dihydro-2,6-dimethyl-4*H*-pyran-one, a homologue of pheromones of a species in the hepialidae family. *Agric. Biol. Chem.* **1987**, *51*, 2001–2002.
11. Eiden, F.; Denk, F. Synthesis and CNS activity of pyrane derivatives 6,8-dioxabicyclo(3,2,1) octanes. *Arch. Pharm. (Weinheim)* **1991**, *324*, 353–354.
12. Shishoo, C.J.; Devani, M.B.; Ullas, G.V.; Ananthan, S.; Bahadit, V.S. Studies in the synthesis and interconversion of isomeric triazolopyrimidinees. *J. Heterocycl. Chem.* **1981**, *18*, 43–46.
13. Noda, K.; Nakagawa, A.; Nakajima, Y.; Ide, H. Pyrido[2,3-*d*]-s-triazolo[4,3-*c*]pyrimidine derivatives. *Japan Kokai*. **1977**, *194*, 7785; [*Chem. Abstr*. **1978**, *88*, P50908q].
14. Brown, D.J., Evans, R.F., Cowden, W.B., Fenn, M.D., Eds. *The Chemistry of Heterocyclic Compounds, the Pyrimidines*; John Wiley & Sons: New York, NY, USA, 1994; Volume 23.
15. Matloobi, M.; Kappe, C.O. Microwave assisted solution- and solid-phase synthesis of amino-4-arylpyrimidine derivatives. *J. Comb. Chem*. **2007**, *9*, 275–284.
16. Gadhachanda, V.R.; Wu, B.; Wang, Z.; Kuhen, K.L.; Caldwell, J.; Zondler, H.; Walter, H.; Havenhand, M.; He, Y. 4-Amino-pyrimidines as novel HIV-1 inhibitors. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 260–265.
17. Agarwal, A.; Srivastava, K.; Puri, S.K.; Chauhan, P.M.S. Synthesis of 4-pyrido-6-aryl-2-substituted amino pyrimidines as a new class of antimalarial agents. *Bioorg. Med. Chem*. **2005**, *13*, 6226–6232.
18. Chang, L.C.W.; Spanjersberg, R.F.; Frijtag, D.K.; Mulder-K, T.; Hout, G.V.; Beukers, M.W.; Brussee, J.; Jzerman, A.P. 2,4,6-Trisubstituted pyrimidines as a new class of selective adenosine A1 receptor antagonists. *J. Med. Chem*. **2004**, *47*, 6529–6540.
19. Capdeville, R.; Buchdunger, E.; Zimmermann, J.; Matter, A. Glivec (STI571, imatinib), a rationally developed, targeted anticancer drug. *Nat. Rev. Drug Discov*. **2002**, *1*, 493–502.
20. Sondhi, S.M.; Dinodia, M.; Rani, R.S.; Shukla, R.; Raghubir, R. Synthesis, anti-inflammatory and analgesic activity evaluation of some pyrimidine derivatives. *Indian J. Chem.* **2009**, *48*, 273–281.
21. Shanker, K.; Gupta, G.P.; Singh, S. Synthesis of novel pyrimidinediones and thiazolidinones as Cardiovascular agents. *Indian J. Chem*. **1985**, *24*, 1094–1097.
22. Ozeki, K.; Ichikawa, T.; Takehara, H.; Tanimura, K.; Sato, M.; Yaginuma, H. Studies on antiallergy agents. III. Synthesis of 2-anilino-1, 6-dihydro-6-oxo-5-pyrimidine carboxylic acids and related compounds; *Chem. Pharm. Bull*. **1989**, *37*, 1780–1787.

23. Elagamey, A.G.A.; El-Taweel, F.M.A.; Sowellim, S.Z.A.; Sofan, M.A.; Elnagdi, M.H. Nitriles in heterocyclic synthesis, a novel route for the synthesis of naphthodipyrans, pyridines, 2*H*- and 4*H*-pyrans. *Collect. Czech. Chem. Commun*. **1990**, *55*, 524–534.
24. Elngdi, M.H.; Elghandour, A.H.H.; Ibrahim, M.K.A.; Hafez, I.S.A. Studies with polyfunctionally substituted heterocyclic, synthesis of new pyrimidines, naphtha[1,2-*b*]pyrans, pyrazolo[3,4-*b*] pyrimidines and pyrazolo[1,5-*a*]pyrimidines. *Z. Naturforsch*. **1992**, *47*, 572–578.
25. Dave, K.G.; Shishoo, C.J.; Devani, M.B.; Kalyanaraman, R.; Ananthan, S.; Ullas, G.V.; Bahadit, V.S. Reaction of nitriles under acidic conditions. Part I: A general method of synthesis of condensed pyridines. *J. Heterocycl. Chem.* **1980**, *17*, 1497–1501.
26. European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID). Determination of minimum inhibitory concentrations (MICs) of antibacterial agents by agar dilution. *Clin. Microbiol. Infect.* **2000**, *6*, 509.
27. National Committee for Clinical Laboratory Standards. *Methods for Dilution AntimicrobialSusceptibility Tests for Bacteria That Grow Aerobically*, 5th ed., Approved Standard M7-A5; NCCLS: Wayne, PA, USA, 2000.

© 2012 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (http://creativecommons.org/licenses/by/3.0/).