

Review

Temporal Trends and Correlates of Physical Activity, Sedentary Behaviour, and Physical Fitness among School-Aged Children in Sub-Saharan Africa: A Systematic Review

Stella K. Muthuri ^{1,2,*}, **Lucy-Joy M. Wachira** ³, **Allana G. Leblanc** ^{1,2}, **Claire E. Francis** ¹,
Margaret Sampson ¹, **Vincent O. Onywera** ^{1,4} and **Mark S. Tremblay** ^{1,2,4}

¹ Children's Hospital of Eastern Ontario Research Institute, 401 Smyth Road, Ottawa, ON K1H 8L1, Canada; E-Mails: alleblanc@cheo.on.ca (A.G.L.); cefrancis89@hotmail.com (C.E.F.); msampson@cheo.on.ca (M.S.); mtremblay@cheo.on.ca (M.S.T.)

² Population Health, Faculty of Graduate and Postdoctoral Studies, University of Ottawa, 1 Stewart Street, Ottawa, ON K1N 7M9, Canada

³ Department of Physical and Health Education, Kenyatta University, P.O. Box 43844, Nairobi, 00100, Kenya; E-Mail: mwlucyjoy@yahoo.com

⁴ Department of Recreation Management and Exercise Science, Kenyatta University, P.O. Box 43844, Nairobi, 00100, Kenya; E-Mail: vonywera@gmail.com

* Author to whom correspondence should be addressed; E-Mail: smuthuri@hotmail.com; Tel.: +1-613-737-7600 (ext. 4191); Fax: +1-613-738-4800.

Received: 9 January 2014; in revised form: 7 March 2014 / Accepted: 10 March 2014 /

Published: 20 March 2014

Abstract: Recent physical activity (PA) and fitness transitions, identified as behavioural shifts from traditionally active lifestyles to more industrialised and sedentary lifestyles, have been observed among school-aged children. There is a wealth of supporting evidence of such behavioural transitions in high income countries; however, a paucity of data on lower income countries exists. These transitions pose a particular threat to the welfare of children by accelerating the onset of chronic diseases. This systematic review investigated the evidence for a PA and fitness transition among Sub-Saharan Africa's school-aged children. Temporal trends and correlates of PA, SB, and fitness were examined. Studies were identified by searching the MEDLINE, Embase, Africa Index Medicus, Global Health, Geobase, and EPPI-Centre electronic databases, and were included if they measured outcomes of interest in apparently healthy samples of children (5–17 years). A total of 71 articles met the inclusion criteria (40 informed PA, 17 informed SB,

and 37 informed fitness). Vast heterogeneity in study methodology complicated analysis of transitions over time and no temporal trends were immediately discernible. However, higher socioeconomic status, urban living, and female children were found to engage in lower levels of PA, higher SB, and performed worse on aerobic fitness measures compared to lower socioeconomic status, rural living, and male children. Data revealed that urbanization was associated with a trend towards decreased PA, increased SB, and decreased aerobic fitness over time. Representative, temporally sequenced data examining a PA and fitness transition are lacking in this region (PROSPERO Registration Number: CRD42013004399).

Keywords: motor activity; sedentary lifestyle; physical fitness; child; adolescent; Sub-Saharan Africa

1. Introduction

Global surveillance efforts have revealed a behavioural shift from traditionally active lifestyles, to more industrialised and sedentary lifestyles [1]. The resultant decline in physical activity levels, coupled with increasing sedentary behaviours over time, is referred to as a “physical activity transition” [1]. The World Health Organization (WHO) classifies physical inactivity as the fourth leading cause of global mortality, and a major determinant for various chronic diseases [2]. Indeed, maintaining an adequate level of physical activity and reducing the amount of time spent in sedentary pursuits is important for the prevention of chronic disease morbidity and mortality [1]. Global physical activity transitions may pose a particular threat to the welfare of children and youth due to the possibility of long-term co-morbidities.

While communicable diseases will likely remain the predominant health problem for the populations in Sub-Saharan Africa (SSA) in the coming years, there is growing concern about rapid increases in non-communicable diseases (NCDs) such as heart disease, diabetes, and hypertension, particularly in urban areas [3]. SSA is currently undergoing rapid socio-cultural developments and urbanization, which have led to the replacement of an economy based on manual labour, to one dominated by industry and mechanized manufacturing [4,5]. This has also resulted in shifts in habitual and occupational physical activity from high-energy expenditure activities (e.g., active transport, manual labour) to low-energy expenditure activities or sedentary behaviours (e.g., motorized transport, desk work) [1]. A transition to lower levels of physical activity is thought to be one of the main contributors to the increasing burden of preventable NCDs in SSA [1,4,6–8]. Of great concern, are the long-term health consequences on children and youth, who are also influenced by this physical activity transition.

A large scale study of secular trends of children and adolescents between 1980 and 2000 using data from 11 mainly developed countries around the world revealed that aerobic fitness (measured by shuttle run tests) had rapidly declined, with the most marked decrease occurring in older age groups, and the rate of decline was similar for boys and girls [9]. This study provided supporting evidence for a decline in physical fitness, particularly aerobic fitness, among children and youth in developed countries.

Whereas physical activity and fitness transitions are well monitored and known to have taken place in high income countries such as Canada, the United States, or the United Kingdom; there is little information on possible physical activity and fitness transitions in SSA, particularly in the school-aged population. Indeed, SSA may be at the early stages of these transitions, and possibly amenable to early intervention strategies aimed at preserving healthy active behaviours. Therefore, the objective of this systematic review was to examine the evidence for a physical activity and fitness transition occurring among SSA's school-aged children and youth. Specifically, temporal trends and correlates of physical activity, sedentary behaviour, and physical fitness in SSA countries were examined. It is important to note that these terms are not interchangeable, particularly among children and youth, as described in Table 1.

Table 1. Outcome measures and their descriptions.

Outcome	Description
Physical Activity	Adequate participation in energy expending activities (e.g., walking, cycling, dancing) provides a wide spectrum of health benefits including reductions in risk for a variety of diseases, improvements in functional ability, and promotes psychological well-being [3,10,11]. Global physical activity guidelines recommend that children and youth, 5–17 years of age, should accumulate at least 60 min of moderate to vigorous intensity physical activity daily [12].
Sedentary Behaviours	These activities (e.g., prolonged sitting, seated screen time, motorized transportation) are characterized by sitting or reclined posture, little physical movement, and low energy expenditure (<1.5 metabolic equivalent tasks) [13]. Canadian sedentary behaviour guidelines developed by the Canadian Society for Exercise Physiology state that for health benefits, children aged 5–17 years should minimize the time they spend being sedentary each day by limiting recreational screen time to no more than 2 h per day, limiting motorized transport, extended sitting, and time spent indoors throughout the day [14].
Physical Fitness	Includes a set of health or skill related attributes that individuals possess in order to perform physical activity [15]. Health related components of physical fitness include cardiorespiratory endurance/aerobic (measured by cycle ergometer, shuttle run, distance run, Harvard step test <i>etc.</i>), musculoskeletal endurance & fitness (measured by pull ups, bent-arm hangs, sit ups <i>etc.</i>), muscular strength (measured by hand grip, trunk lift <i>etc.</i>), body composition measures (measured by body mass index, body fat percentage, waist circumference <i>etc.</i>), flexibility (measured by sit-and-reach <i>etc.</i>), and anaerobic power (measured by dash/sprint runs, jumps <i>etc.</i>) [15].

2. Methods

2.1. Study Inclusion Criteria

All published, peer-reviewed studies were included if they reported using subjective or objective measures of physical activity, sedentary behaviour, or physical fitness in children and youth aged 5 to 17 years, with no chronic conditions, and living in SSA. The relevant outcome measures are described in Table 1.

2.2. Study Exclusion Criteria

In order to obtain information on a general population living under typical conditions, intervention studies were excluded unless they conducted baseline measurements. No date limits were imposed, but due to feasibility, studies in languages other than English or French were excluded. Studies were also excluded if they did not include one or more of the relevant health indicators.

2.3. Search Strategy

Studies were identified using the following electronic databases: Ovid MEDLINE (1948 to Week 4, May 2013), Ovid Embase (1974 to Week 21, 2013), Africa Index Medicus (database dates not available, searched on 3 June 2013), Global Health (1973 to 3 June 2013 through the CAB direct interface), Geobase (1884 to 3 June 2013 through the Engineering Village interface), and EPPI-Centre database of health promotion research (Bibliomap) (dates of coverage not available, searched 3 June 2013). The search strategy for this systematic review was completed in tandem with another publication examining the evidence for an overweight/obesity transition among school-aged children and youth in SSA [16]; hence, the inclusion of these terms in the search strategy. The search strategy was created and run by a research librarian. The complete search strategy used for MEDLINE is presented in Table 2. References were exported and de-duplicated using Reference Manager Software (Version 11, Thompson Reuters, San Francisco, CA, USA). Titles and abstracts of potentially relevant articles were screened by two independent reviewers, and full text articles were obtained for those meeting initial screening criteria. The full text articles were then screened in duplicate for inclusion in the review. This review is registered with the international prospective register of systematic reviews PROSPERO network (registration number: CRD42013004399) [17].

Table 2. MEDLINE search strategy; Ovid interface.

Order	Search Terms
1	exp “Africa South of the Sahara”/
2	(sub-sahar * or east afric * or south afric * or keny * or (south adj3 sahar *)).mp.
3	1 or 2
4	sedentar\$.tw.
5	Sedentary Lifestyle/
6	((chair or sitting or car or automobile or auto or bus or indoor or in-door or screen or computer) adj time).tw.
7	low energy expenditure.tw.
8	(computer game * or video game * or ((television adj watch *) or tv watch *)).tw.
9	television/ or computers/ or video games/
10	(screen based entertainment or screen-based entertainment or screen time).tw.
11	physical inactivit *.tw.
12	bed rest.mp.
13	sitting.tw.
14	exp obesity/
15	(obesity * or obese).tw.
16	exp overweight/
17	(overweight or over weight).tw.

Table 2. Cont.

Order	Search Terms
18	exp Body Fat Distribution/
19	exp body composition/
20	Waist Circumference/
21	waist circumference.tw.
22	Skinfold Thickness/
23	(skin folds or skin fold *).tw.
24	(body composition * or BMI or body mass index).tw.
25	exp “body weights and measures”/
26	(bio-impedance analysis or BIA).tw.
27	Absorptiometry, Photon/
28	(absorptiometry or densitometry or photodensitometry or DXA or DEXA).tw.
29	Physical Fitness
30	(physical conditioning or physical fitness).tw.
31	musculoskeletal fitness.tw.
32	physical endurance/
33	cardiovascular fitness.tw.
34	motor activit\$.tw.
35	physical exertion/
36	aerobic exercise.tw.
37	exp sports
38	play/
39	exp physical education/
40	musculoskeletal physiological processes/ or exercise/ or movement/ or locomotion/ or running/ or swimming/ or walking/ or motor activity/
41	or/4–40
42	(child * or adolescent * or youth * or pediatric * or paediatric *).tw.
43	3 and 41 and 42

Notes: The search strategy for this systematic review was completed in tandem with a sister publication examining the evidence for an overweight/obesity transition among school-age children and youth in Sub Saharan Africa [16]; hence, the inclusion of these terms in the search strategy.

2.4. Data Extraction, Synthesis and Quality Assessment

Data extraction was completed using a standardized data extraction template, and study quality was assessed using a modified Downs and Black instrument [18]. The Downs and Black checklist for measuring quality of evidence was selected for use due to its suitability for quality assessment of original research articles (beyond the typical gauges used for quality assessment of evidence from systematic reviews and meta-analyses) [18]. Due to limitations in study design, questions selected from the Downs and Black quality assessment instrument excluded any questions that referred to intervention and trial study methodology, leaving ten out of a possible 27 questions, as represented in Table 3.

Table 3. Modified Downs and Black checklist [18].

Reporting
Objective Clearly Stated—Question 1 from full checklist ($Y = 1/N = 0$)
Main Outcomes Clearly Described—Question 2 ($Y = 1/N = 0$)
Patient Characteristics Clearly Defined—Question 3 ($Y = 1/N = 0$)
Main Findings Clearly Defined—Question 6 ($Y = 1/N = 0$)
Random Variability in Estimates Provided—Question 7 ($Y = 1/N = 0$)
Actual Probability Values Reported—Question 10 ($Y = 1/N = 0$)
External Validity
Sample Targeted Representative of Population—Question 11 ($Y = 1/N = 0$)
Sample Recruited Representative of Population—Question 12 ($Y = 1/N = 0$)
Internal Validity/Bias
Statistical Tests Used Appropriately—Question 18 ($Y = 1/N = 0$)
Primary Outcomes Valid/Reliable—Question 20 ($Y = 1/N = 0$)

3. Results

Figure 1 shows the PRISMA flow diagram with numbers of included and excluded articles at each step of the review process, while Table 4 provides a summary of all studies included in this systematic review. A total of 2,657 records were identified through database searches and other sources. Following duplicate removal, 2,242 were screened for eligibility, and 663 were selected for a full-text review. Of these, a total of 71 articles met inclusion criteria, comprising a total sample of 77,515 participants from 17 SSA countries. Reasons for exclusion included: irrelevant population (e.g., studies that did not involve children 5–17 years of age with no pre-existing condition) (181 articles); population living in a country outside of SSA (10); irrelevant outcomes (334); and, excluded study design (67 articles).

Table 4. Descriptive characteristics of included studies.

First Author [reference]	Year	Study Design	Country	Age (Years)	Sample (n)	Outcome (Measures)	D&B Score
Sloan [19]	1967	Cross sectional	South Africa	15–17	393	PF (anaerobic fitness, musculoskeletal fitness)	7
Areskog [20]	1969	Cross sectional	Ethiopia	9–14	153	PF (aerobic fitness, musculoskeletal fitness)	7
Stephenson [21]	1985	Cross sectional	Kenya	7–15	12	PF (aerobic fitness)	7
Corlett [22]	1986	Cross sectional	Botswana	6–11	289	PF (musculoskeletal fitness)	6
Ndamba [23]	1986	Cross sectional	Zimbabwe	8–15	147	PF (aerobic fitness)	7
Corlett [24]	1988	Cross sectional	Botswana	7–12	612	PF (musculoskeletal fitness)	7
Benefice [25]	1992	Cross sectional	Senegal	9–14	100	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness)	7
Proctor [26]	1996	Cross sectional	Cameroon	9–14	119	PA (self-report)	7
Benefice [27]	1996	Cross sectional	Senegal	5–13	348	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness)	7

Table 4. *Cont.*

First Author [reference]	Year	Study Design	Country	Age (Years)	Sample (n)	Outcome (Measures)	D&B Score
Benefice [28]	1998	Cross sectional	Senegal	5–13	348	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness)	7
Prista [29]	1998	Cross sectional	Mozambique	8–15	593	PA (self-report) & PF (aerobic fitness, musculoskeletal fitness)	8
Benefice [30]	1999	Cross sectional	Senegal	12–13	221	PA (direct measures)	8
Benefice [31]	2001	Cross sectional	Senegal	13	40	PA (direct measures)	8
Benefice [32]	2001	Cross sectional	Senegal	13	40	PA (direct measures)	8
Garnier [33]	2001	Cross sectional	Senegal	13–15	80	PA (direct measures) & SB (direct measures)	8
Prista [34]	2003	Cross sectional	Mozambique	6–17	2,316	PA (self-report) & PF (musculoskeletal fitness)	8
McVeigh [35]	2004	Cross sectional	South Africa	9	386	PA (self-report) & SB (self- report)	7
McVeigh [36]	2004	Cross sectional	South Africa	10	386	PA (self-report)	7
Micklesfield [37]	2004	Cross sectional	South Africa	7–11	198	PA (self-report)	7
Larsen [38]	2004	Cross sectional	Kenya	15–17	11	PA (self-report)	7
Benefice [39]	2004	Cross sectional	Senegal	13–15	40	SB (direct measures)	7
Monyeki [40]	2004	Longitudinal	South Africa	7	85	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness, balance/flexibility)	8
Monyeki [41]	2005	Longitudinal	South Africa	7–14	855	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness)	8
Benefice [42]	2005	Cross sectional	Senegal	10–13	99	PA (direct)	8
Aandstad [43]	2006	Cross sectional	Tanzania	9–10	156	PA (self-report) & PF (aerobic fitness)	7
Garnier [44]	2006	Cross sectional	Senegal	13–15	80	SB (direct measures)	7
Djarova [45]	2006	Cross sectional	Zimbabwe	6–14	49	PA (self-report) & PF (musculoskeletal fitness)	6
Onyewadume [46]	2006	Cross sectional	Botswana	11–14	30	PF (musculoskeletal fitness)	8
Micklesfield [47]	2007	Cross sectional	South Africa	9	64	PA (self-report) & SB (self-report) & PF (musculoskeletal fitness)	7
Monyeki [48]	2007	Longitudinal	South Africa	7–14	702	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness)	8

Table 4. Cont.

First Author [reference]	Year	Study Design	Country	Age (Years)	Sample (n)	Outcome (Measures)	D&B Score
Bovet [49] ^{a,b,c}	2007	Cross sectional	Seychelles	12–15	4,343	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness)	9
Travill [50]	2007	Cross sectional	South Africa	8–17	720	PF (anaerobic fitness, musculoskeletal fitness, balance/flexibility)	7
Monyeki [51]	2008	Longitudinal	South Africa	7–13	1,817	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness, balance/flexibility)	7
Lennox [52]	2008	Cross sectional	South Africa	15	318	PA (self-report) & SB (self-report) & PF (musculoskeletal fitness, balance/flexibility)	8
Prista [53]	2009	Cross sectional	Mozambique	6–16	256	PA (direct measures)	8
Berntsen [54]	2009	Cross sectional	Tanzania	9–10	190	PF (aerobic fitness)	8
Peltzer [55]	2009	Secondary analysis	Namibia, Kenya, Uganda, Zimbabwe	13–15	12,740	PA (self-report) & SB (self-report)	9
Peltzer [56]	2010	Secondary analysis	Botswana, Kenya, Namibia, Senegal, Swaziland, Uganda, Zambia, and Zimbabwe	13–15	24,593	PA (self-report) & SB (self-report)	9
Harmse [57]	2010	Cross sectional	South Africa	13–17	221	PF (aerobic fitness)	7
Senbanjo [58]	2010	Cross sectional	Nigeria	5–14	392	PA (self-report)	8
Truter [59]	2010	Cross sectional	South Africa	9–13	280	PF (aerobic fitness)	7
Odunaiya [60]	2010	Cross sectional	Nigeria	14–16	608	PA (self-report)	7
Ansa [61]	2010	Cross sectional	Nigeria	10–17	964	PA (self-report)	8
Adeniyi [62]	2011	Cross sectional	Nigeria	13–17	1,100	PA (self-report)	7
Peltzer [63] ^c	2011	Secondary analysis	Ghana & Uganda	13–15	5,613	PA (self-report) & SB (self-report)	9
Naude [64]	2011	Cross sectional	South Africa	12–16	162	PA (self-report) & SB (self-report)	5
Croteau [65]	2011	Cross sectional	Kenya	8–12	72	PA (direct measures)	8
Muller [66]	2011	Cross sectional	Côte d'Ivoire	7–15	17	PF (aerobic fitness)	7
Dapi [67]	2011	Cross sectional	Cameroon	12–16	227	PA (self-report)	8
Puckree [68]	2011	Cross sectional	South Africa	10–12	120	PA (self-report) & SB (self-report)	7


Table 4. *Cont.*

First Author [reference]	Year	Study Design	Country	Age (Years)	Sample (n)	Outcome (Measures)	D&B Score
Armstrong [69] ^c	2011	Cross sectional	South Africa	6–13	10,295	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness, balance/flexibility)	10
Adamo [70]	2011	Cross sectional	Kenya	9–13	179	PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness)	7
Naidoo [71]	2012	Cross sectional	South Africa	7–10	170	PF (aerobic fitness)	7
Musa [72]	2012	Cross sectional	Nigeria	9–15	3,243	PF (aerobic fitness)	7
Monyeki [73]	2012	Longitudinal	South Africa	14	256	PF (anaerobic fitness, musculoskeletal fitness)	8
Onywera [74]	2012	Cross sectional	Kenya	9–12	169	PA (direct measures & self-report) & SB (self-report)	7
Ojiambo [75]	2012	Cross sectional	Kenya	12–16	200	PA (direct measures & self-report) & SB (direct measures & self-report)	7
Richards [76]	2012	Cross sectional	Uganda	11–14	31	PA (direct measures)	8
Micklesfield [77]	2012	Cross sectional	South Africa	11–15	381	PA (self-report) & SB (self-report)	6
Monyeki [78]	2012	Cross sectional	South Africa	14	256	PF (anaerobic fitness, musculoskeletal fitness)	8
Monyeki [79]	2012	Cross sectional	South Africa	14–15	153	PA (self-report) & PF (aerobic fitness)	8
Truter [80]	2012	Cross sectional	South Africa	9–13	280	PF (anaerobic fitness)	7
Bovet [81] ^{a,b}	2012	Cross sectional	Seychelles	9–16	8,462	PA (self-report)	9
Pienaar [82]	2012	Longitudinal	South Africa	13–17	87	PA (self-report)	8
Toriola [83]	2012	Longitudinal	South Africa	14	283	PA (self-report) & SB (self-report) & PF (aerobic fitness, anaerobic fitness, musculoskeletal fitness, balance/flexibility)	8
Craig [84]	2012	Secondary analysis	South Africa	7, 11, 15	89	PA (direct measures)	9
Ojiambo [85]	2012	Cross sectional	Kenya	13–16	200	PA (direct measures) & SB (direct measures)	7
Malete [86]	2013	Cross sectional	Botswana	13–16	756	PA (self-report) & SB (self-report)	7

Table 4. Cont.

First Author [reference]	Year	Study Design	Country	Age (Years)	Sample (n)	Outcome (Measures)	D&B Score
Onywera [87]	2013	Cross sectional	Kenya	9–13	179	PA (direct measures)	7
Heroux [88]	2013	Cross sectional	Kenya	9–13	179	PF (musculoskeletal fitness)	7
			Total (n)	77,515	Average D&B score	7.5	

Notes: D & B score (Downs & Black score) [18]; PA = physical activity, SB = sedentary behaviour, PF = physical fitness. ^a = Article indicated targeting a sample size representative of the population of interest (2 articles). ^b = Article indicated recruiting a sample size representative of the population of interest (2 articles). ^c = Article indicated that the sample size was nationally representative (3 articles).

Figure 1. PRISMA flow chart of search strategy results.

Notes: PRISMA flow diagram [89]: ^a Reasons for exclusion included irrelevant population (181 studies), irrelevant geographic location/not Sub Saharan Africa (10 studies), irrelevant outcomes (334 studies), and irrelevant content/study design e.g. intervention programs or review articles (67 studies).

Data quality assessment revealed that the average score for all included studies was 7.5 out of a possible 10, pointing to a fairly high quality of evidence of the individual articles. The earliest relevant record captured was published in 1967, with two articles published between 1960 and 1969,

none between 1970 and 1979, four articles between 1980 and 1989, six articles between 1990 and 1999, 26 articles between 2000 and 2009, and 33 articles published between 2010 and June 2013. This substantial increase in the publishing rate may be representative of a growing interest in this field of research. There was also good representation of the four regions of SSA with 20 (28.2%) articles from East African countries, 30 (42.3%) from South African countries, 16 (22.5%) from West African countries, two (2.8%) from Central African countries, and three (4.2%) that were combined.

3.1. Physical Activity

As shown in Table 5, 36 studies comprising 62,188 participants, presented in 40 articles, examined physical activity outcome(s). Of the 36 studies, 26 (72.2%) studies [26,29,34,35,37,38,43,45,47,52,55,56,58,60–64,67,68,77,79,81–83,86] used solely subjective measurements of physical activity (e.g., self-report or interviewer administered questionnaires), and 10 (27.8%) studies [30–33,42,53,65,74–76,84] used direct/objective measures (e.g., accelerometers, pedometers) or a combination of direct and subjective measures. Table 5 indicates the type the type of measure used and the main findings for each of the 36 studies. Despite the heterogeneity in the types of physical activity measures used in the included studies, some findings stood out from the data taken together.

Table 5. Findings from studies reporting on physical activity outcomes.

First Author [reference]	Year	Country	Age (years)	(n) Total	(n) Male	(n) Female	Main Finding(s) *
Subjective Measures (Self-report or interviewer administered questionnaires)							
Proctor [26]	1996	Cameroon	9–14	119	65	54	Physical activity among rural children was more than twice that of urban children, and was mostly work related and heavier intensity.
Prista [29]	1998	Mozambique	8–15	593	277	316	Underprivileged children had higher levels of physical activity due to survival activities (e.g., walking).
Prista [34]	2003	Mozambique	6–17	2,316	1,094	1,222	Low SES children and adolescents had higher levels of physical activity due to higher demands of survival activities and playing, but spent less time in formal sports than their more privileged peers.
McVeigh [35]	2004	South Africa	9	386	202	184	Inverted bell shape curve for association between percent physical activity levels and SES. The highest SES had the highest MVPA. Also discusses differences between different ethnicities.
Micklesfield [37]	2004	South Africa	7–11	198	0	198	Discussed differences between different ethnicities.

Table 5. *Cont.*

First Author [reference]	Year	Country	Age (years)	(n) Total	(n) Male	(n) Female	Main Finding(s) *
Subjective Measures (Self-report or interviewer administered questionnaires)							
Larsen [38]	2004	Kenya	15–17	11	11	0	Village boys were significantly more habitually active (e.g., running, field work) than town boys.
Aandstad [43]	2006	Tanzania	9–10	156	87	69	88% of rural children walked to school, 71% for more than 15 min, and 34% participated in outdoor games after school on most or all days.
Djarova [45]	2006	Zimbabwe	6–14	49			Mean duration of activities during the day (hours): Sleeping 8.2, walking 2.4, sitting 3.9, standing 2.0, running 1.0, sports 2.5, playing 4.0.
Micklesfield [47]	2007	South Africa	9	64	36	28	Mean hours of participation in school physical education 1.1, school sports 1.4, informal activities 7.7, and weekend activities 8.
Lennox [52]	2008	South Africa	15	318	137	181	Lower SES school had higher activity levels, particularly, walking to school.
Peltzer [55]	2009	Namibia, Kenya, Uganda, Zimbabwe	13–15	12,740	7,517	5,223	Physical activity on ≥ 3 days (for ≥ 60 min) ranged from 24.4% to 36.0%. Active transport on ≥ 5 days (for ≥ 60 min) ranged from 19.8% to 31.1%.
Peltzer [56]	2010	Botswana, Kenya, Namibia, Senegal, Swaziland, Uganda, Zambia, and Zimbabwe	13–15	24,593	10,575	14,018	A range of 9.0% to 17.7% of children reported being physically active on ≥ 5 days (for ≥ 60 min). The average was 14.2% in the total sample, 16.6% in males, and 12.0% in females. 18.9% of the total sample was physically active on ≥ 3 days (for ≥ 60 min).

Table 5. Cont.

First Author [reference]	Year	Country	Age (years)	(n) Total	(n) Male	(n) Female	Main Finding(s) *
Subjective Measures (Self-report or interviewer administered questionnaires)							
Senbanjo [58]	2010	Nigeria	5–14	392	202	190	Frequency of leisure time physical activity on ≥ 3 days in 5–9 year old males (3.2%), females (6.0%), in 10–14 year old males (20.1%), females (12.3%). Leisure time physical activity was higher in older children, in males, and in children of mothers with higher education level.
Odunaiya [60]	2010	Nigeria	14–16	608			64.1% of the total sample participated in moderate and high physical activity.
Ansa [61]	2010	Nigeria	10–17	964	472	492	75% of the total sample participated in outdoor sports. No SES effect found (not significant).
Adeniyi [62]	2011	Nigeria	13–17	1,100	538	562	Females had higher risk of low physical activity than boys.
Peltzer [63]	2011	Ghana & Uganda	13–15	5,613	2,738	2,875	78.5% of males and 84.9% of females were physically active for less than 60 min per day on at least 5 days per week.
Naude [64]	2011	South Africa	12–16	162			33% of the total sample participated in organized sports.
Dapi [67]	2011	Cameroon	12–16	227	108	119	Boys had significantly higher levels of physical activity than girls. No SES effect found (not significant).
Puckree [68]	2011	South Africa	10–12	120	48	72	92% of the total sample participated in extracurricular sporting activities.
Micklesfield [77]	2012	South Africa	11–15	381			Informal activity was significantly lower in older children. MVPA was significantly lower in girls than boys at all ages. Low SES was associated with higher walking for transport and lower MVPA in schools or clubs.
Monyeki [79]	2012	South Africa	14–15	153	0	153	26.8% of girls participated in moderate and high physical activity.
Bovet [81]	2012	Seychelles	9–16	8,462	4,239	4,223	There was higher leisure time physical activity in private compared to public schools, but higher active transport (walking) in public compared to private schools.

Table 5. Cont.

First Author [reference]	Year	Country	Age (years)	(n) Total	(n) Male	(n) Female	Main Finding(s) *
Subjective Measures (Self-report or interviewer administered questionnaires)							
Pienaar [82]	2012	South Africa	13–17	87	28	59	There was increased leisure time physical activity over time, higher house-chore related physical activity on weekends compared to weekdays, and higher exercise related activity on week days compared to weekends.
Toriola [83]	2012	South Africa	14	283	111	172	Boys had higher physical activity in the high/vigorous level compared to girls.
Malete [86]	2013	Botswana	13–16	756	464	292	Low SES and rural children had more minutes of physical activity than high SES or urban children.
Direct/Objective Measures							
Benefice [30]	1999	Senegal	13	221	0	221	Instrument: CSA accelerometer. Findings: Percentage of time spent during 12 hours of a day in MVPA (>940 counts/min corresponding to >6 METs) was 21.7% in the dry season, 29.3% in the rainy season, and 24.3% (corresponding to 2.9 h) in total.
Benefice [31]	2001	Senegal	13	(40 at each visit)	120		Instrument: CSA accelerometer. Findings: Time spent in moderate and vigorous activity was 29.12% (corresponding to 4.4 h) at the 1st visit in 1997, 25.71% (corresponding to 3.9 h) at the 2nd visit in 1998, and 25.54% (corresponding to 3.8 h) at the 3rd visit in 1999. Daily physical activity was expressed as a physical activity level unit (ratio of energy expenditure to BMR). Three units of BMR corresponded to 890 counts/min. MVPA was therefore a BMR corresponding to >1,890 counts/min and was calculated as a percent of 15 h or time between 7 a.m. and 10 p.m.
Garnier [33]	2001	Senegal	13–15	80	0	80	Instrument: CSA accelerometer. Findings: Migrants to the city spent more time engaged in moderate to heavy activity than did non-migrants in the rural area (9.3 h/24 h vs. 6.1 h/24 h). Intensity thresholds were defined according to Benefice & Cames, whereby MVPA was >610 counts/min or ≥3 METs [30].

Table 5. Cont.

First Author [reference]	Year	Country	Age (years)	(n) Total	(n) Male	(n) Female	Main Finding(s) *
Direct/Objective Measures							
Benefice [42]	2005	Senegal	10–13	99	46	53	Instrument: Cardio-frequencemeters. Findings: In adolescent girls, body composition was a significant predictor of activity levels.
Prista [53]	2009	Mozambique	6–16	256	139	117	Instrument(s): Actigraph accelerometer and self-report measures. Findings: Software converted actigraph counts into relative energy exposure using the regression equation developed by Freedson <i>et al.</i> for children 6–18 years of age [90]. The average minutes of time spent in MVPA (≥ 3 METs) in boys and girls 6–8 years of age was 232.2 and 235.5, in boys and girls 9–11 years of age was 202.0 and 210.3, and in boys and girls 12–16 years of age was 157.4 and 158.7. This corresponds to range of 2.6–3.9 hours for the total sample.
Croteau [65]	2011	Kenya	8–12	72	29	43	Instrument(s): Pedometers (step count data) and self-report measures. Findings: Daily steps in total sample $14,558 \pm 3,993$. Boys ($16,262 \pm 4,698$) were significantly more active than girls ($13,463 \pm 3,051$). No significant effect for age was found. Observed daily steps are higher than those observed in the USA and samples in other developed countries.
Onywera [74]	2012	Kenya	9–12	169	85	84	Instrument(s): Pedometers (step count data) and self-report measures. Findings: Rural children were more physically active than urban children. 87% rural children and 42% urban children used active transport to get to school.
Ojiambo [75]	2012	Kenya	12–16	200	99	101	Instrument: Actigraph accelerometer. Physical activity levels were assessed using cut-points developed by Puyau <i>et al.</i> whereby MVPA $>3,200$ counts/min [91]. The mean minutes of MVPA were 54 (corresponding to 0.9 h) in the total sample, 68 in rural males, 62 in rural females, 50 in urban males, and 37 in urban females.

Table 5. Cont.

First Author [reference]	Year	Country	Age (years)	(n) Total	(n) Male	(n) Female	Main Finding(s) *
Direct/Objective Measures							
Richards [76]	2012	Uganda	11–14	31	31	0	Instrument: Actigraph accelerometer. Findings: Mean minutes per day of MVPA were 114.09 (corresponding to 1.9 h), with only 4.32 minutes in the vigorous range. (MVPA cut-points not indicated).
Craig [84]	2013	South Africa	7–15	89	46	43	Instrument: Actigraph accelerometer. Findings: Total physical activity was generally high (mean accelerometer counts per minute ranged 485–1017), but MVPA was low with less than 1% of the total sample meeting the MVPA guidelines using the Puyau <i>et al.</i> cut points [91]. MVPA in boys comprised 1.7% of waking hours (12 min/day) at age 7, 1.3% (10 min/day) at age 11, and 0.6% (4 min/day) at age 15. MVPA in girls comprised 1.5% (11 min/day) at age 7, 0.9% (6 min/day) at age 11, and 0.2% (1 min/day) at age 15 years.

Notes: SES (socioeconomic status); MVPA (moderate-to-vigorous physical activity); Counts/min (counts per min); BMR (basal metabolic rate); METs (metabolic equivalents). Table excludes references [32,36,85,87] which used an identical study sample as used in another included manuscript. * Main finding(s) from the article as relates to physical activity outcomes.

3.1.1. Physical Activity Transition

Of the articles that used direct measures, seven studies [30,31,33,53,75,76,84] used accelerometry to measure physical activity. Only one study examined the changing trends over time. Benefice *et al.* found a decline in the amount of time spent in moderate-to-vigorous intensity physical activity (MVPA) from 29.12% (corresponding to 4.4 h) in 1997, to 25.71% (3.9 h) in 1998 and 25.54% (3.8 h) in 1999 [31]. Studies that used accelerometry used varying MVPA cut-points to categorise their samples making it impossible to comment on trends over time. As such, the general trend towards decreasing proportions of MVPA from earlier studies (1999–2009) with sample means of ≥ 2.6 h of MVPA [30,31,33,53] to later studies (2012) with sample means of ≤ 1.9 h of MVPA [75,76,84] as observed in Table 5, may be related to more stringent cut-points rather than decreasing levels of physical activity in these samples. In general, a lack of population-representative samples and heterogeneity of measurement protocols did not allow for a clear assessment of physical activity level trends over time.

One large scale, multi-country study using subjective measures with a sample of 24,593 participants reported that 14.2% (16.6% of males, 12.0% of females) of children were physically active on 5 days or more, for at least 60 min/day during leisure time [56]. The proportion was 18.9% of children on 3 days or more, for at least 60 min/day during leisure time [56]. Step count data determined that the

observed average number of daily steps was higher than those of the United States and samples in other developed countries [65].

3.1.2. Socioeconomic (SES) and Urban/Rural Differences

Of the 14 studies using subjective measures to examine the association between physical activity and SES, 11 found that lower SES and rural children had higher levels of physical activity compared to higher SES and urban children, or children who had mothers with a higher educational level [26,29,34,38,43,52,58,74,77,81,86], while three found contradicting evidence or no SES differences [35,61,67]. Lower SES and rural children engaged in higher levels of active transportation (e.g., walking and running to school) [43,52,74,77,81], spent more time in activities of daily living (e.g., house chores, work related, habitual activity) [26,29,34,38], but, spent less time engaged in organized sports or formal activities compared to their higher SES and urban peers [77,81]. Directly measured physical activity (both accelerometry and pedometer data) also found that rural children were more physically active than urban children [74,75].

3.1.3. Sex Differences

Of the six studies using subjective data collection methods and that examined the differences in physical activity trends in boys and girls, five studies found higher levels of physical activity (e.g., informal activity, leisure time physical activity, MVPA) in boys compared to girls irrespective of age [56,63,67,77,83], while one study found that this was only true in the higher age-group (10–14 years), but not in lower age-group (5–9 years) [58]. Similarly, directly measured physical activity (both accelerometry and pedometer data) found that boys were significantly more active than girls [65,84].

3.2. Sedentary Behaviours

As shown in Table 6, a total of 17 studies examining sedentary behaviours, comprising 33,525 participants, were included in this review. Of these, the majority (12 studies) used self-report measures [35,47,52,55,56,63,64,68,74,77,83,86], and five studies used direct measurements [33,39,44,75,85]. Specific measurement tools varied across studies, but total weekday and/or weekend television viewing time was most often studied. All studies examining sedentary behaviour were published after 2000, with the majority published in the last three years (*i.e.*, since 2010).

Table 6. Findings from studies reporting on sedentary behaviour outcomes.

First Author [reference]	Year	Country	Age (Years)	(n) Total	(n) Male	(n) Female	Main Finding(s) *
Subjective Measures (Self-report or interviewer administered questionnaires)							
McVeigh [35]	2004	South Africa	9	386	202	184	Sedentary activity (h/day) among white girls 8.61 (± 0.54), black girls 9.03 (± 0.34), white boys 10.48 (± 0.79), and black boys 10.63 (± 0.33).

Table 6. *Cont.*

First Author [reference]	Year	Country	Age (Years)	(n) Total	(n) Male	(n) Female	Main Finding(s)*
Subjective Measures (Self-report or interviewer administered questionnaires)							
Micklesfield [47]	2007	South Africa	9	64	36	28	Average min per day spent on television watching was 123 ± 8.3 .
Lennox [52]	2008	South Africa	15	318	137	181	TV h during the week in school 1 “(lower SES) & school 2 (higher SES) among boys 1.92 (1.43) & 2.24 (1.27), in school 1 & 2 among girls 1.76 (1.32) & 2.53 (1.38). TV hrs during the weekend in school 1 & 2 among boys 3.31 (2.10) & 3.62 (2.50), in school 1 & 2 among girls 2.74 (1.94) & 3.28 (2.38).
Peltzer [55]	2009	Namibia, Kenya, Uganda, Zimbabwe	13–15	12,740	7,517	5,223	Proportion of children reporting less than 3 h of sitting/day was 56.6% for Kenya, 64.3% for Namibia, 70.3% for Uganda, and 56.3% for Zimbabwe.
Peltzer [56]	2010	Botswana, Kenya, Namibia, Senegal, Swaziland, Uganda, Zambia, and Zimbabwe	13–15	24,593	10,575	14,018	Overall time spent sitting on a usual day for <1 h was 39.4%, 1–2 h was 32.7%, 3–4 h was 17.5%, ≥ 5 h was 11.2%.
Peltzer [63]	2011	Ghana and Uganda	13–15	5,613	2,738	2,875	Sedentary behavior (≥ 3 h/day) in males was 27.1% (23.6–30.5), in females was 26.9% (24.3–29.6).
Naude [64]	2011	South Africa	12–16	162			Mean total weekly time (min) spent watching TV or playing on the computer was 1,001 (± 570).
Puckree [68]	2011	South Africa	10–12	120	48	72	All sampled children reported watching >4 h of TV on weekdays and 6–8 h on weekends.
Onywera [74]	2012	Kenya	9–12	169	85	84	30% of rural children spent >2 h/week on screen time compared with 50% of urban children, and 62.5% of the rural children spent 0 h/week playing screen games compared with 13.1% of urban children.

Table 6. Cont.

First Author [reference]	Year	Country	Age (Years)	(n) Total	(n) Male	(n) Female	Main Finding(s)*
Subjective Measures (Self-report or interviewer administered questionnaires)							
Micklesfield [77]	2012	South Africa	11–15	381			Sedentary activity was significantly higher in older girls compared to younger counterparts ($p < 0.05$). Increasing pubertal status was associated with an increase in sedentary activity. Lower SES at the maternal, household, and community level was significantly associated with less sedentary activity.
Toriola [83]	2012	South Africa	14	283	111	172	Overall, 18%, 23%, and 40% watched TV for >3 h, 2–3 h, <1 h per day, respectively. More girls (19%) than boys (16%) watched TV for >3 h/day.
Malete [86]	2013	Botswana	13–16	756	464	292	Time spent sitting (min/week) was 2,612.38 (± 910.87) or 6.2 h/day.
Direct/Objective Measures							
Garnier [33]	2001	Senegal	13–15	80	0	80	Instrument: CSA accelerometer. Findings: Sleeping and inactivity of migrants (who move for work) was 6 h and 42 min, and of non-migrants 8 h and 29 min.
Benefice [39]	2004	Senegal	13–15	40 (in each year)	40 (in each year)		Instrument: CSA accelerometer. Findings: Resting time (min)—defined as the number of zero counts during the day—was 93 (± 54) in the year 1997, 111 (± 65) in 1998, and 103 (± 51) in 1999.
Garnier [44]	2006	Senegal	13–15	80	40	40	Instrument: CSA accelerometer. Findings: Girls were more inactive (4 h and 23 min) than boys 2 h and 49 min).
Ojiambo [75]	2012	Kenya	12–16	200	99	101	Instrument: Actigraph accelerometer. Findings: Min (%) sedentary time in total sample was 584 ± 113 (72%); in rural males was 555 ± 67 (65%), in rural females was 539 ± 91 (66%), in urban males was 678 ± 95 (78%), and in urban females was 694 ± 81 (80%).
Ojiambo [85]	2012	Kenya	13–16	200	98	102	Instrument: Actigraph accelerometer. Findings: Daily mean sedentary time (min) was 619 ± 109 , or approximately 78% of the monitored time.

Notes: SES (socioeconomic status); TV (television). * Main finding(s) from the article as relates to sedentary behaviour outcomes.

3.2.1. Sedentary Behaviour Transition

In South Africa, self-reported time spent watching television was inconsistent. Recent work by Puckree *et al.* showed that 100% of participants aged 10–12 years watched more than 4 h of television on weekdays, and 6–8 h of television on weekends [68]. Toriola and Monyeki reported that 40% of 14 year old participants watched less than 1 hour of television per day [83]. In other studies, the self-reported mean television viewing time was between 2 and 2.5 h per day for 9–16 year old children, with weekend television viewing slightly higher at 2.7 to 3.6 h, and television viewing generally higher in older children [47,52,64]. Data from Ghana and Uganda found that 27% of children aged 13–15 years spent more than 3 h per day in sedentary pursuits [63], while in Botswana, 14 year olds spent an average of 6.2 hours per day sitting [86]. Senegalese youth reported only 1.33 to 1.41 h of sedentary time per day [39].

One multi-country comparative study found that in children 13–15 years of age, the proportion reporting less than 3 h of sitting per day was 29.7% in Uganda, 25.7% in Namibia, 43.4% in Kenya, and 43.7% in Zimbabwe [55]. Another large scale multi-country study using data from eight African countries (Botswana, Kenya, Namibia, Senegal, Swaziland, Uganda, Zambia, and Zimbabwe) found that 39.4% of children aged 13–15 years spent less than one hour per day sitting, and an additional 32.7% spent 1–2 h per day sitting when not in school or doing homework [56].

Notably, the short time span within which the studies reporting on sedentary behaviour were published made it impossible to examine transitions over time, since temporal trends are often reasonably stable over short periods of time.

3.2.2. Sex Differences

Accelerometry measured sedentary time showed that Senegalese girls spent more time in sedentary behaviours than boys (4.23 h vs. 2.49 h) [44]. Investigation of self-reported television viewing also found that more girls (19%) than boys (16%) watched television for more than 3 hours daily [83]. Most studies however either found no significant difference or did not report on the difference in sedentary behaviours between boys and girls [35,52,63,75].

3.2.3. Age Differences

One included study examined age differences and found that sedentary activity was significantly higher in older girls compared to their younger counterparts [77]. This study also showed that increasing pubertal status, which is not a proxy for age, was associated with an increase in sedentary behaviours, but that the most significant determinant of sedentary behaviour was SES [77].

3.2.4. SES and Urban/Rural Differences

Higher SES and urban living children in Kenya were found to spend significantly more time in sedentary pursuits than their lower SES and rural counterparts, with approximately 50% of the urban children, and only 30% of the rural children reporting spending over 2 hours each week on screen time activities [74,75]. This is consistent with the findings of Lennox *et al.* showing that South African children in higher SES schools spent more time watching television than children in lower SES

schools on both weekdays and weekends [52]. Additionally, Micklesfield *et al.* found that higher maternal, household, and community level SES were significantly associated with increased sedentary time [77]. In contrast, Malete *et al.*, found that public school students and those living in rural villages (lower SES) reported significantly more minutes of sitting than students in private schools or students from cities (higher SES); which is an unexpected finding given that the former group also reported more vigorous and total physical activity than the latter [86].

3.3. Physical Fitness

Thirty six included studies, presented in 37 papers, comprising 30,452 participants, examined physical fitness measures. The majority of these studies were published in the past three years, but studies were included from as early as 1967 [19]. Almost half (47.2%) of the 36 fitness related studies included South African participants. All studies used direct measures of physical fitness. Studies are presented based on whether they examined aerobic fitness, anaerobic fitness, musculoskeletal fitness and strength, or balance and flexibility, with many of the studies examining more than one aspect of physical fitness.

3.3.1. Aerobic Fitness

Twenty one studies presented data on aerobic fitness. Aerobic fitness was measured through shuttle run [29,41,43,48,49,51,52,57,66,69–72,79,83,88], maximal cycle ergometer test [20,54], maximal aerobic capacity [52,59], distance run [23,29,41,48,51], and the Harvard Step test [21]. Boys were found to have higher aerobic capacity and performed considerably better on measures of aerobic fitness than girls [23,41,49,54,59,66,72,83,88].

Two studies found that lower SES and rural children performed better on measures of aerobic fitness than higher SES and urban children [52,70]. In contrast, one study found the opposite to be true when comparing girls in the townships (lower SES) to girls in the towns (higher SES) [79]. It is noteworthy that in the latter study, township girls were also found to have significantly higher prevalence of underweight than the town girls. Aandstad *et al.* found that Tanzanian children had better aerobic fitness test results when compared to a Norwegian reference group [43].

Only one study examined changes over time in aerobic fitness tests. In their study of 702 rural South African children, Monyeki *et al.* found significant improvements in aerobic fitness results (shuttle run and 1,600 m run tests) from 2001 to 2002 though this was only across one year [48]. However, when data from studies using similar measures of aerobic fitness (*i.e.*, shuttle run [41,48,51,69] and 1,600 m run [41,48,51]) were examined, we found largely unaltered trends in aerobic fitness over time, with boys performing better than girls as shown in Table 7.

3.3.2. Anaerobic Fitness

Seventeen studies presented data on anaerobic fitness. This was measured though timed dashes/sprints [19,25,27,28,40,41,48–50,70] or long jumps [25,27,28,40,41,48–51,69,73,78,83]. Boys performed better in all measures of anaerobic fitness than girls [41,49,83]. Anaerobic test parameters in Senegalese children were found to be inferior to those of well-nourished Western children [25,28]. No temporal trend evidence was available.

Table 7. Findings from studies reporting on comparable physical fitness outcomes.

First Author [reference]	Year	Country	Age (Years)	(n)	(n)	(10 × 5m) Shuttle Run (sec)	Run 1,600		Grip Strength (kg)		Sit & Reach (cm)	
				Boys	Girls		B	G	B	G	B	G
Corlett [24]	1988	Botswana	7 (Urb)	18	17				11.9	11.6		
									(0.3)	(1.0)		
			7 (Rur)	19	31				7.4	6.3		
									(1.3)	(1.1)		
			8 (Urb)	26	29				12.4	11.6		
									(0.8)	(0.5)		
			8 (Rur)	25	24				9.2	7.8		
									(1.1)	(1.6)		
			9 (Urb)	18	30				13.4	12.7		
									(0.9)	(0.8)		
			9 (Rur)	32	37				9.8	8.4		
									(1.1)	(1.0)		
Monyeki [48] *	2001	South Africa	10 (Urb)	28	20				15.6	14.0		
									(1.2)	(0.6)		
			10 (Rur)	43	25				12.4	10.0		
									(2.2)	(1.7)		
			11 (Urb)	24	18				17.4	15.1		
									(1.3)	(1.1)		
			11 (Rur)	30	38				13.7	12.8		
									(1.7)	(2.0)		
			12 (Urb)	17	16				20.2	16.8		
									(1.4)	(1.5)		
			12 (Rur)	36	32				16.5	15.2		
									(2.3)	(2.8)		
Monyeki [40]	2004	South Africa	7–10	152	133	22.9	23.2	489.8	531.2			
						(2.1)	(1.9)	(42.0)	(62.9)			
			7–10	228	189	21.9	21.8	475.3	514.6			
						(2.1)	(2.1)	(48.0)	(57.7)			
Monyeki [41]	2005	South Africa	11–15	152	133	22.0	22.2	452.9	511.8			
						(1.8)	(1.7)	(40.5)	(59.8)			
			11–15	228	189	21.2	21.5	460.7	529.3			
						(1.9)	(1.6)	(53.8)	(79.6)			

Table 7. Cont.

First Author [reference]	Year	Country	Age (Years)	(n) Boys	(n) Girls	(10 × 5m) Shuttle Run (sec)		Run 1,600 (m)		Grip Strength (kg)		Sit & Reach (cm)		
				B	G	B	G	B	G	B	G	B	G	
Onyewadume [46]	2006	Botswana	11–14	15	15	22.0	22.8	456	510	34.6	24.1	14.4	15.8	
						(2.2)	(1.9)	(39.6)	(53.0)	(7.3)	(5.1)	(5.1)	(4.6)	
						11	69	74	21.8	22.1	450	498	15.7	16.9
								(1.9)	(1.7)	(41.0)	(50.3)	(5.1)	(5.1)	
						12	87	67	21.8	21.9	450	510	13.8	17.8
								(1.6)	(1.5)	(40.4)	(44.1)	(4.8)	(5.0)	
Monyeki [51]	2008	South Africa	7	80	61	23.8	23.7	549.2	570.7	29.0	31.1	12.8	17.6	
						(2.6)	(2.9)	(94.6)	(69.0)			(5.3)	(4.2)	
						8	86	82	23.0	23.3	523.6	563.6		
								(2.3)	(2.6)	(76.3)	(87.2)			
						9	107	89	22.7	22.3	500.2	535.0		
								(2.2)	(2.0)	(78.9)	(80.9)			
Lennox [52]	2008	South Africa	10	102	108	22.3	22.4	496.7	527.6	27.5	29.1	School 1		
						(1.8)	(1.7)	(84.0)	(67.6)					
						11	111	94	21.9	21.8	479.5	526.9		
								(2.1)	(1.8)	(64.3)	(74.2)			
						12	99	82	21.7	21.9	475.5	523.0		
								(1.7)	(1.9)	(83.1)	(82.2)			
Armstrong [69]	2011	South Africa	13	49	42	21.4	21.6	463.9	501.9	27.6	28.6			
						(1.4)	(1.3)	(96.9)	(128.6)					
						15 (RR)	116	136						
						15 (RL)	116	136						
						15 (RR)	21	45						
						15 (RL)	21	45						
Toriola [83]	2012	South Africa	6–13	5,611	4,684	22.8	23.8			21.4	23.7			
Heroux [88]	2013	Kenya	14	111	172					42.2	48.5			
										(9.1)	(7.3)			
[88]	2013	Kenya	9–13	86	93					34.7	31.1			
										(32.0,	(28.7			
										37.3)	, 3.5)			

Notes: B (boys); G (girls); Urb (urban); Rur (rural); RR (reach-right); RL (reach-left). Where reported, variance included in brackets as (standard deviation) or (confidence intervals). School 1 had lower socioeconomic status than school 2 [52]. (*) Study was published in 2007, but data was collected in the years indicated, *i.e.*, 2001 and 2002.

3.3.3. Musculoskeletal Fitness and Strength

Twenty five studies examined musculoskeletal fitness and strength. This was measured via bent arm hang [41,48,51,73,78,83], grip strength [20,24,25,27,29,45–47,50,52,68,70,88], timed push-ups [46,49], timed sit-ups [19,29,40,41,46,48,49,51,52,69,73,78,83], or throwing [19,25,27,28,49,69]. Boys performed better than girls in measures of bent arm, sit-ups, and push-ups [41,83].

As shown in Table 7, when data from studies using a similar measure of grip strength (kilograms) were examined, we found a trend towards increasing scores in both boys and girls over time [24,46,88]. It is however important to note that geographical differences between Botswana [24,46] and Kenya [88] may be clouding these findings. Further, even within the studies from Botswana, there was little overlap among the age groups, further complicating temporal trend analysis. From these data, it was also determined that boys performed better than girls in grip strength measurements [24,46,88]. Grip strength seemingly increased with age [50], and was found to be higher in higher SES and urban children compared to lower SES and rural children [24,70]. In general, measures of skeletal muscle fitness and strength were found to be lower in SSA children compared to various Western reference samples (e.g., United States, the Netherlands) [22,25,28,50].

3.3.4. Balance and Flexibility

Eight studies presented data on balance and flexibility. This was measured via flamingo balance [41,51] and sit-and-reach test [22,40,41,50,52,69,83]. Girls were more flexible and performed better in balance tests than boys [41,50,83]. As shown in Table 7, when data from studies examining sit and reach (cm) performance were assessed, we found a trend towards increasing measures of flexibility in both sexes, with girls performing consistently better than boys [40,41,52,69,83]. It is however important to note that these apparent increases may be a function of the way in which sit-and reach performance was scored, since we observe an approximate 3-fold increase in scores over an 8 year period among South African children [40,41,52,69,83].

4. Discussion

This is the first systematic review to comprehensively explore the evidence for a physical activity and fitness transition, and their correlates, among SSA's children and youth.

4.1. Physical Activity Transition

Subjective methods of assessing physical activity are commonly used to measure types, frequencies, durations, and levels of physical activity. This is likely due to low costs associated with measurement [92]. Previous work however has shown that there is little association between self-report and directly measured physical activity; largely as a result of poor recall in children [92–95]. A majority of the included studies examining physical activity used subjective methods of data collection. A smaller proportion of studies used direct methods to assess physical activity, thereby providing higher accuracy and validity [96]. The means of measured MVPA reported in studies that used accelerometry to monitor physical activity ranged from 1 min/day to 9.3 h/day [30,31,33,53,75,76,84]. The only included study that examined changes in reported time

spent in MVPA found a decline in MVPA of 30 minutes per day from 1997 to 1999 [31]. Regrettably, those using direct methods also used vastly different measurement devices and cut-points, making it impossible to conduct legitimate quantitative analyses of the results.

Assessment of a physical activity time trend was challenging given that many of the subjective instruments used have yet to be properly validated and are fraught with various limitations, while the objective measures used were mainly cross-sectional and differed in methodology [97]. Owing to the vast heterogeneity of these results, little could be said definitively on whether there was indeed evidence of a physical activity transition over time. It is imperative that future work examine the duration and intensity of physical activity on population representative samples using common measurement techniques and sampling procedures.

Similarly, a clear transition to increasing time spent in sedentary behaviours was not apparent. The majority of studies reporting on sedentary behaviour outcomes were published after 2010, further complicating analysis of trends over time. The reported means of time spent in sedentary pursuits ranged from 1.3 hours to 6 hours on weekdays, and were as high as 8 h on weekends [39,47,52,64,68,86]. These means are problematic given that the Canadian sedentary behaviour guidelines (the first published evidence-based guidelines on sedentary behaviour) recommend that children 5–17 years should limit their recreational sedentary screen time to no more than 2 h per day [14]. The urbanization trend suggests an increase in sedentary behaviours over time as the data revealed higher television viewing times among urban and higher SES children than rural and lower SES children.

Quantitative and narrative analyses of studies examining physical fitness measures revealed a tendency towards increased fitness measures, particularly in grip strength and flexibility over time in SSA, though the inconsistencies in measures used and representativeness of the samples makes direct comparisons across studies tenuous. It has been postulated that body composition indices should be interpreted differently in undernourished populations, that is, higher body mass index measures could be seen as a measure of muscle mass instead of a measure of fatness [41,48]. With this in mind, whereas previously the prevalence of underweight in SSA's school-aged children was high, these numbers are falling, giving rise to higher proportions of well-nourished children who may have greater muscle mass and perform better in various fitness measures. Improvements in fitness parameters in SSA's school-aged children may also be as a result of an increase in the proportion of children participating in formal/organized sports or activities, which has enabled these children to improve certain physical fitness skills that may not be learned through informal activities, random play, and self-directed activities engaged in more by the rural living and lower SES children. Despite these apparent improvements, when compared to Western reference groups, SSA children performed better in aerobic fitness measures but worse in anaerobic fitness tests or measures of musculoskeletal fitness and strength [22,25,28,43].

4.2. SES Differences

This systematic review found that lower SES and rural living children had higher levels of physical activity compared to higher SES and urban living children. Higher activity levels in lower SES children are associated with higher demands of informal/survival activities such as household chores

and walking from place to place (active transport), rather than the formal/organized sporting activities engaged in by their more privileged peers. Higher SES and urban living children were also found to spend more time in sedentary pursuits than lower SES and rural living children; consistent with the finding that lower SES children were more active. This finding is likely due to lower access to motorized transport and higher need for their contribution to household and other tasks in lower SES settings. Lastly, lower SES children were also found to perform better in aerobic fitness measures compared to their higher SES peers. Superior aerobic fitness performance in lower SES and rural children may indeed be as a direct result of their higher participation in habitual active transport (e.g., walking, running), when compared to higher SES and urban children. In the one study where lower SES girls performed worse than their higher SES peers, this may be explained by the high prevalence of underweight that was also found in the low SES group, resulting in lower muscle mass as previously described; hence, poor performance in aerobic fitness tests [41,48,79]. The same may be said about the finding that higher SES children perform better in grip strength measures than lower SES children.

4.3. Sex Differences

This review found significantly higher levels of physical activity in boys compared to girls irrespective of age. Similarly, boys engaged in less sedentary behaviours (e.g., television viewing) than girls. Boys were also found to have considerably better aerobic, anaerobic, musculoskeletal fitness, and strength measures than girls; however, girls performed better on balance and flexibility measures of fitness. Besides biology and socio-cultural roles, this superior physical and functional ability of boys may be explained by their higher motivation to participate in physical activities.

The use of high quality methodology to capture and synthesize the studies in this review offers strength to the findings and conclusions in this manuscript. Additionally, decisions were made *a priori* to limit possible bias, and review processes were conducted in duplicate to ensure a higher level of accuracy. However, this review had several limitations including the vast heterogeneity in study sampling and methodology, which complicated quantitative analysis and direct comparison. Further, any trends observed over time may be reflective of the growth or differences in the ages of participants rather than group differences. It is also unclear if any material relevant for this review may have been published in un-indexed journals and hence not captured by the literature search.

5. Concluding Remarks

Owing to the vast heterogeneity in measurement devices, methodologies, and cut-points used by studies included in this review, little could be said definitively on the evidence for a physical activity and fitness transition over time. More generally however, the data revealed that urbanization was associated with a developing trend towards decreasing physical activity, increasing sedentary behaviours, and decreasing fitness measures (particularly aerobic fitness) over time. This was shown by engagement in lower amounts and levels of physical activity, higher television viewing and other sedentary pursuits, and lower aerobic fitness levels among urban and higher SES children compared to their rural and lower SES peers. As such, proactive strategies to prevent decreased physical activity

and fitness and increased sedentary behaviours in the context of a probable physical activity transition in children from SSA appears warranted.

This systematic review also revealed a critical lack of representative, temporally sequenced data on physical activity, sedentary behaviours, and physical fitness measures in SSA's school-aged children and youth, which is a largely understudied and vulnerable group [98]. It is our recommendation that future work entail concerted efforts in carrying out nationally representative surveys, using comparable or common measurement techniques, sampling procedures, and cut-points, in order to effectively monitor physical activity transitions over time in this region. Further, considering that costs and ease of accelerometry use have improved over the last decade, we recommend that objective monitoring of physical activity and sedentary time be used in future studies. For instance, multi-country surveys would best strengthen the knowledge base in this field of research.

Acknowledgements

The authors are grateful to Alison McFarlane for her contributions towards manuscript formatting.

Author Contributions

Stella Muthuri, Mark Tremblay, Margaret Sampson, Lucy-Joy Wachira, Allana Leblanc, and Vincent Onywera developed the systematic review research question and objectives. Margaret Sampson created the search strategy and provided guidance on review methodology. Stella Muthuri, Claire Francis, Lucy-Joy Wachira, and Allana Leblanc screened and extracted the evidence. Stella Muthuri and Mark Tremblay led data analysis, synthesis, and writing of the manuscript. All authors contributed to interpretation of the results, edited, reviewed, and approved the final manuscript.

Conflicts of Interest

The authors declare that they have no competing interests.

References

1. Katzmarzyk, P.T.; Mason, C. The physical activity transition. *J. Phys. Act. Health* **2009**, *6*, 269–280.
2. World Health Organization (WHO). *Global Health Risks: Mortality and Burden of Disease Attributable to Selected Major Risks*. WHO: Geneva, Switzerland, 2009.
3. Unwin, N.; Setel, P.; Rashid, S.; Mugusi, F.; Mbanya, J.; Kitange, H.; Hayes, L.; Edwards, R.; Aspray, T.; Alberti, K.G. Noncommunicable diseases in Sub-Saharan Africa: Where do they feature in the health research agenda? *Bull. World Health Organ.* **2001**, *79*, 947–953.
4. Omran, A. The epidemiologic transition theory: A preliminary update. *J. Trop. Pediatr.* **1983**, *29*, 305–316.
5. Popkin, B.M.; Gordon-Larsen, P. The nutrition transition: Worldwide obesity dynamics and their determinants. *Int. J. Obes. Relat. Metab. Disord.* **2004**, *28*, S2–S9.
6. Popkin, B. The nutrition transition and its health implications in lower income countries. *Public Health Nutr.* **1998**, *1*, 5–21.

7. Tremblay, M.S.; Onywera, V.; Adamo, K.B. A Child’s Right to Healthy, Active Living—Building Capacity in Sub-Saharan Africa to Curb the Impeding Physical Activity Transition: The KIDS-CAN Research Alliance. In *20th Anniversary of the Convention on the Rights of the Child*; Bennett, S., Pare, M., Eds.; University of Ottawa Press: Ottawa, ON, Canada, 2010; pp. 97–110.
8. World Health Organization. *Obesity: Preventing and Managing the Global Epidemic*; WHO: Geneva, Switzerland, 2000.
9. Tomkinson, G.R.; Léger, L.A.; Olds, T.S.; Cazorla, G. Secular trends in the performance of children and adolescents (1980–2000): An analysis of 55 studies of the 20m shuttle run test in 11 countries. *Sport. Med.* **2003**, *33*, 285–300.
10. Powell, K.; Pratt, M. Physical activity and health. *BMJ* **1996**, *313*, 126–127.
11. Powell, K.E.; Paluch, A.E.; Blair, S.N. Physical activity for health: What kind? How much? How intense? On top of what? *Annu. Rev. Public Health* **2011**, *32*, 349–365.
12. World Health Organization. *Global Recommendations on Physical Activity for Health*; WHO: Geneva, Switzerland, 2010.
13. Tremblay, M.; Colley, R.; Saunders, T.; Healey, G.; Owen, N. Physiological and health implications of a sedentary lifestyle. *Appl. Physiol. Nutr. Metab.* **2010**, *35*, 1–16.
14. Canadian Society for Exercise Physiology. Canadian Physical Activity Guidelines. Available online: www.csep.ca/guidelines (accessed on 20 August 2013).
15. Caspersen, C.J.; Powell, K.E.; Christenson, G.M. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep.* **1985**, *100*, 126–131.
16. Muthuri, S.K.; Francis, C.E.; Wachira, L.M.; LeBlanc, A.G.; Sampson, M.; Onywera, V.O.; Tremblay, M.S. Evidence of an overweight/obesity transition among school-aged children and youth in Sub-Saharan Africa: A systematic review. *PLoS One* **2014**, in press.
17. The PROSPERO Network. Available online: <http://www.crd.york.ac.uk/prospero/> (accessed on 18 March 2014).
18. Downs, S.H.; Black, N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J. Epidemiol. Community Health* **1998**, *52*, 377–384.
19. Sloan, A.W.; Wiggett, R. Inter-racial comparison of tests of physical fitness on high school children at Cape Town. *J. Sports Med. Phys. Fitness* **1967**, *7*, 192–197.
20. Areskog, N.H.; Selinus, R.; Vahlquist, B. Physical work capacity and nutritional status in Ethiopian male children and young adults. *Amer. J. Clin. Nutr.* **1969**, *22*, 471–479.
21. Stephenson, L.S.; Latham, M.C.; Kurz, K.M.; Miller, D.; Kinoti, S.N.; Oduori, M.L. Urinary iron loss and physical fitness of Kenyan children with urinary schistosomiasis. *Am. J. Trop. Med. Hyg.* **1985**, *34*, 322–330.
22. Corlett, J.T. Minimum muscular fitness of urban Tswana children. *Child. Care Health Dev.* **1985**, *11*, 37–43.
23. Ndamba, J. Schistosomiasis: Its effects on the physical performance of school children in Zimbabwe. *Cent. Afr. J. Med.* **1986**, *32*, 289–293.
24. Corlett, J.T. Strength development of Tswana children. *Hum. Biol.* **1988**, *60*, 569–577.
25. Benefice, E. Physical activity and anthropometric and functional characteristics of mildly malnourished Senegalese children. *Ann. Trop. Paediatr.* **1992**, *12*, 55–66.

26. Proctor, M.H.; Moore, L.L.; Singer, M.R.; Hood, M.Y.; Nguyen, U.S.; Ellison, R.C. Risk profiles for non-communicable diseases in rural and urban schoolchildren in the Republic of Cameroon. *Ethn. Dis.* **1996**, *6*, 235–243.
27. Benefice, E.; Malina, R. Body Size, Body composition and motor performances of mild-to-moderately undernourished Senegalese children. *Ann. Hum. Biol.* **1996**, *23*, 307–321.
28. Benefice, E. Growth and Motor Performances of Rural Senegalese Children. In *Physical Fitness and Nutrition during Growth*; Parizkova, J., Hills, A.P., Eds.; Karger Medical and Scientific Publishers: Brisbane, QLD, Australia, 1998.
29. Prista, A. Nutritional Status, Physical Fitness and Physical Activity in Children and Youth in Maputo (Mozambique). In *Physical Fitness and Nutrition during Growth*; Parizkova, J., Hills, A.P., Eds.; Karger Medical and Scientific Publishers: Brisbane, QLD, Australia, 1998, pp. 117–131.
30. Benefice, E.; Came, C. Physical activity patterns of rural Senegalese adolescent girls during the dry and rainy seasons measured by movement registration and direct observation methods. *Eur. J. Clin. Nutr.* **1999**, *53*, 636–643.
31. Benefice, E.; Garnier, D.; Ndiaye, G. High levels of habitual physical activity in west African adolescent girls and relationship to maturation, growth, and nutritional status: Results from a 3-year prospective study. *Amer. J. Human Biol.* **2001**, *13*, 808–820.
32. Benefice, E.; Garnier, D.; Ndiaye, G. Assessment of physical activity among rural Senegalese adolescent girls: Influence of age, sexual maturation, and body composition. *J. Adolesc. Health* **2001**, *28*, 319–327.
33. Garnier, D.; Benefice, E. Habitual physical activity of Senegalese adolescent girls under different working conditions, as assessed by a questionnaire and movement registration. *Ann. Hum. Biol.* **2001**, *28*, 79–97.
34. Prista, A.; Maia, J.A.; Damasceno, A.; Beunen, G. Anthropometric indicators of nutritional status: implications for fitness, activity, and health in school-age children and adolescents from Maputo, Mozambique. *Amer. J. Clin. Nutr.* **2003**, *77*, 952–959.
35. McVeigh, J.A.; Norris, S.A.; Cameron, N.; Pettifor, J.M. Associations between physical activity and bone mass in black and white South African children at age 9. *J. Appl. Physiol.* **2004**, *97*, 1006–1012.
36. McVeigh, J.A.; Norris, S.A.; de Wet, T. The relationship between socio-economic status and physical activity patterns in South African children. *Acta Paediatr.* **2004**, *93*, 982–988.
37. Micklesfield, L.K.; Zielonka, E.A.; Charlton, K.E.; Katzenellenbogen, L.; Harkins, J.; Lambert, E.V. Ultrasound bone measurements in pre-adolescent girls: Interaction between ethnicity and lifestyle factors. *Acta Paediatr.* **2004**, *93*, 752–758.
38. Larsen, H.B.; Christensen, D.L.; Nolan, T.; Sondergaard, H. Body dimensions, exercise capacity and physical activity level of adolescent Nandi boys in western Kenya. *Ann. Hum. Biol.* **2004**, *31*, 159–173.
39. Benefice, E.; Garnier, D.; Ndiaye, G. Nutritional status, growth and sleep habits among Senegalese adolescent girls. *Eur. J. Clin. Nutr.* **2004**, *58*, 292–301.
40. Monyeki, M.A.; Toriola, A.L.; Monyeki, K.D.; Brits, S.J.; Pienaar, A.E. Body size, body composition and physical fitness of 7-years-old Ellisras rural children, South Africa: Ellisras Longitudinal Study (ELS). *Afri. J. Phys. Health Educ. Recreat. Dance* **2004**, *10*, 154–162.

41. Monyeki, M.A.; Koppes, L.L.; Kemper, H.C.; Monyeki, K.D.; Toriola, A.L.; Pienaar, A.E.; Twisk, J.W. Body composition and physical fitness of undernourished South African rural primary school children. *Eur. J. Clin. Nutr.* **2005**, *59*, 877–883.
42. Benefice, E.; Ndiaye, G. Relationships between anthropometry, cardiorespiratory fitness indices and physical activity levels in different age and sex groups in rural Senegal (west Africa). *Ann. Hum. Biol.* **2005**, *32*, 366–382.
43. Aandstad, A.; Berntsen, S.; Hageberg, R.; Klasson-Heggebo, L.; Anderssen, S.A. A comparison of estimated maximal oxygen uptake in 9 and 10 year old schoolchildren in Tanzania and Norway. *BJSM Online* **2006**, *40*, 287–292.
44. Garnier, D.; Benefice, E. Reliable method to estimate characteristics of sleep and physical inactivity in free-living conditions using accelerometry. *Ann. Epidemiol.* **2006**, *16*, 364–369.
45. Djarova, T.; Dube, S.; Tivchev, G.; Chivengo, A. Nutritional profiles, physical development and daily activities of African children in zimbabwe with insulin-dependent diabetes mellitus. *S. Afr. J. Sci.* **2006**, *102*, 4–6.
46. Onyewadume, I.U. Fitness of black African early adolescents with and without mild mental retardation. *APAQ* **2006**, *23*, 277–292.
47. Micklesfield, L.K.; Levitt, N.S.; Carstens, M.T.; Dhansay, M.A.; Norris, S.A.; Lambert, E.V. Early life and current determinants of bone in South African children of mixed ancestral origin. *Ann. Hum. Biol.* **2007**, *34*, 647–655.
48. Monyeki, M.A.; Koppes, L.L.; Monyeki, K.D.; Kemper, H.C.; Twisk, J.W. Longitudinal relationships between nutritional status, body composition, and physical fitness in rural children of South Africa: The Ellisras longitudinal study. *Amer. J. Human Biol.* **2007**, *19*, 551–558.
49. Bovet, P.; Auguste, R.; Burdette, H. Strong inverse association between physical fitness and overweight in adolescents: A large school-based survey. *Int. J. Behav. Nutr. Phys. Act.* **2007**, *4*, doi:10.1186/1479-5868-4-24.
50. Travill, A.L. Growth and physical fitness of socially disadvantaged boys and girls aged 8–17 years living in the Western Cape, South Africa. *Afri. J. Phys. Health Educ. Recreat. Dance* **2007**, *13*, 279–293.
51. Monyeki, K.D.; Kemper, H.C.; Makgae, P.J. Relationship between fat patterns, physical fitness and blood pressure of rural South African children: Ellisras longitudinal growth and health study. *J. Hum. Hypertens.* **2008**, *22*, 311–319.
52. Lennox, A.; Pienaar, A.E.; Wilders, C. Physical fitness and the physical activity status of 15-year-old adolescents in a semi-urban community. *S. Afr. J. Res Sport Phys. Educ. Recreat.* **2008**, *30*, 59–73.
53. Prista, A.; Nhantumbo, L.; Silvio, S.; Lopes, V.; Maia, J.; Andre, S.; Vinagre, J.; Conn, C.A.; Beunen, G. Physical activity assessed by accelerometry in rural african school-age children and adolescents. *Pediatr. Exerc. Sci.* **2009**, *21*, 384–399.
54. Berntsen, S.; Lodrup Carlsen, K.C.; Hageberg, R.; Aandstad, A.; Mowinckel, P.; Anderssen, S.A.; Carlsen, K.H. Asthma symptoms in rural living Tanzanian children; prevalence and the relation to aerobic fitness and body fat. *Allergy* **2009**, *64*, 1166–1171.
55. Peltzer, K. Health behavior and protective factors among school children in four African countries. *Int. J. Behav. Med.* **2009**, *16*, 172–180.

56. Peltzer, K. Leisure time physical activity and sedentary behavior and substance use among in-school adolescents in eight African countries. *Int. J. Behav. Med.* **2010**, *17*, 271–278.
57. Harmse, B.; Kruger, H.S. Significant differences between serum crp levels in children in different categories of physical activity: The PLAY study. *Cardiovasc. J. Afr.* **2010**, *21*, 316–322.
58. Senbanjo, I.O.; Oshikoya, K.A. Physical activity and body mass index of school children and adolescents in Abeokuta, Southwest Nigeria. *World J. Pediatr.* **2010**, *6*, 217–222.
59. Truter, L.; Pienaar, A.E.; Du Toit, D. Relationships between overweight, obesity and physical fitness of nine- to twelve-year-old South African children. *S. Afr. Fam. Pract.* **2010**, *52*, 227–233.
60. Odunaiya, N.A.; Ayodele, O.A.; Oguntibeju, O.O. Physical activity levels of senior secondary school students in Ibadan, Western Nigeria. *West. Indian Med. J.* **2010**, *59*, 529–534.
61. Ansa, V.O.; Anah, M.U.; Odey, F.A.; Mbu, P.N.; Agbor, E.I. Relationship between parental socio-economic status and casual blood pressure in Coastal Nigerian adolescents. *West Afr. J. Med.* **2010**, *29*, 146–152.
62. Adeniyi, A.F.; Okafor, N.C.; Adeniyi, C.Y. Depression and physical activity in a sample of Nigerian adolescents: Levels, relationships and predictors. *Child. Adolescent Psychiatr. Mental Health* **2011**, *5*, doi:10.1186/1753-2000-5-16.
63. Peltzer, K.; Pengpid, S. Overweight and obesity and associated factors among school-aged adolescents in Ghana and Uganda. *Int. J. Environ. Res. Public Health* **2011**, *8*, 3859–3870.
64. Naude, C.E.; Senekal, M.; Laubscher, R.; Carey, P.D.; Fein, G. Growth and weight status in treatment-naive 12–16 year old adolescents with alcohol use disorders in Cape Town, South Africa. *Nutr. J.* **2011**, *10*, doi:10.1186/1475-2891-10-87.
65. Croteau, K.; Schofield, G.; Towle, G.; Suresh, V. Pedometer-determined physical activity of western Kenyan children. *J. Phys. Act. Health* **2011**, *8*, 824–828.
66. Muller, I.; Coulibaly, J.T.; Furst, T.; Knopp, S.; Hattendorf, J.; Krauth, S.J.; Stete, K.; Righetti, A.A.; Glinz, D.; Yao, A.K.; et al. Effect of Schistosomiasis and soil-transmitted helminth infections on physical fitness of school children in Cote d'Ivoire. *PLoS Negl. Trop. Dis.* **2011**, *5*, doi:10.1371/journal.pntd.0001239.
67. Dapi, L.N.; Hornell, A.; Janlert, U.; Stenlund, H.; Larsson, C. Energy and nutrient intakes in relation to sex and socio-economic status among school adolescents in urban Cameroon, Africa. *Public Health Nutr.* **2011**, *14*, 904–913.
68. Puckree, T.; Naidoo, P.; Pill, P.; Naidoo, T. Underweight and overweight in primary school children in Ethekwini district in KwaZulu-Natal, South Africa. *Afr. J. Primary Health Care Fam. Med.* **2011**, *3*, 1–6.
69. Armstrong, M.E.; Lambert, E.V.; Lambert, M.I. Physical fitness of South African primary school children, 6 to 13 years of age: Discovery vitality health of the nation study. *Percept. Mot. Skills* **2011**, *113*, 999–1016.
70. Adamo, K.B.; Sheel, A.W.; Onywera, V.; Waudo, J.; Boit, M.; Tremblay, M.S. Child obesity and fitness levels among Kenyan and Canadian children from urban and rural environments: A KIDS-CAN research alliance study. *Int. J. Pediatr. Obes.* **2011**, *6*, 225–232.
71. Naidoo, T.; Konkol, K.; Biccard, B.; Dudose, K.; McKune, A.J. Elevated salivary c-reactive protein predicted by low cardio-respiratory fitness and being overweight in African children. *Cardiovasc. J. Afr.* **2012**, *23*, 501–506.

72. Musa, D.I.; Williams, C.A. Cardiorespiratory fitness, fatness, and blood pressure associations in Nigerian youth. *Med. Sci. Sports Exerc.* **2012**, *44*, 1978–1985.
73. Monyeki, M.A.; Neetens, R.; Moss, S.J.; Twisk, J. The relationship between body composition and physical fitness in 14 year old adolescents residing within the tlokwe local municipality, South Africa: The PAHL study. *BMC Public Health* **2012**, *12*, doi:10.1186/1471-2458-12-374.
74. Onywera, V.O.; Adamo, K.B.; Sheel, A.W.; Waudo, J.N.; Boit, M.K.; Tremblay, M. Emerging Evidence of the Physical Activity Transition in Kenya. *J. Phys. Act. Health* **2012**, *9*, 554–562.
75. Ojiambo, R.M.; Easton, C.; Casajus, J.A.; Konstabel, K.; Reilly, J.J.; Pitsiladis, Y. Effect of urbanization on objectively measured physical activity levels, sedentary time, and indices of adiposity in Kenyan adolescents. *J. Phys. Act. Health* **2012**, *9*, 115–123.
76. Richards, J.; Foster, C. The Impact of a Sport-for-Development Programme on the Physical Activity Levels of Young Adolescent Boys in Gulu, Northern Uganda. In Proceedings of the Be Active 2012 Conference, Sydney, Australia, 26–28 September 2012; pp. S257–S258.
77. Micklesfield, L.; Pedro, T.; Twine, R.; Kinsman, J.; Pettifor, J.; Tollman, S.; Kahn, K.; Norris, S. Physical Activity Patterns and Determinants in Rural South African Adolescents. In Proceedings of the Be Active 2012 Conference, Sydney, Australia, 26–28 September 2012; p. S251.
78. Monyeki, M.; Neetens, R.; Moss, S.; Twisk, J. The Relationship of Body Composition with Physical Fitness in the 14 Adolescents Residing within the Tlokwe Local Municipality, South Africa: The PAHL Study. In Proceedings of the Be Active 2012 Conference, Sydney, Australia, 26–28 September 2012; pp. S170–S171.
79. Monyeki, M.; Mamabolo, R. Relationships between Physical Activity with BMI and Percentage Body Fat among Girls by Locality in Tlokwe Local Municipality, South Africa: The PAHL-study. In Proceedings of the Be Active 2012 Conference, Sydney, Australia, 26–28 September 2012; p. S42.
80. Truter, L.; Pienaar, A.E.; Toit, D. The relationship of overweight and obesity to the motor performance of children living in South Africa. *S. Afr. Fam. Pract.* **2012**, *54*, 429–435.
81. Bovet, P.; Paccaud, F.; Chiolero, A. Socio-economic status and obesity in children in Africa. *Obes. Rev.* **2012**, *13*, doi:10.1111/j.1467-789X.2012.01023.x.
82. Pienaar, A.E.; Salome, K.H.; Steyn, H.S.; Naude, D. Change over three years in adolescents' physical activity levels and patterns after a physical activity intervention: PLAY study. *J. Sports Med. Phys. Fitness* **2012**, *52*, 300–310.
83. Toriola, O.M.; Monyeki, M.A. Health-related fitness, body composition and physical activity status among adolescent learners: The PAHL study. *Afr. J. Phys. Health Educ. Recreat. Dance* **2012**, *18*, 795–811.
84. Craig, E.M.; Reilly, J.J.; Bland, R.M. How Best to Assess Unhealthy Weight Status? In Proceedings of Archives of Disease in Childhood Conference: Annual Conference of the Royal College of Paediatrics and Child Health, Glasgow, Scotland, 22–24 May 2012; pp. 45–46.
85. Ojiambo, R.M.; Magutah, K.; Thairu, K.; Takahashi, R.; Wilunda, C. The relation between physical activity and indicators of body fatness in Kenyan adolescents. *J. Appl. Med. Sci.* **2013**, *2*, 43–59.
86. Malete, L.; Motlhoiwa, K.; Shaibu, S.; Wrotniak, B.H.; Maruapula, S.D.; Jackson, J.; Compher, C.W. Body image dissatisfaction is increased in male and overweight/obese adolescents in Botswana. *J. Obes.* **2013**, *2013*, doi:10.1155/2013/763624.

87. Onywera, V.O.; Heroux, M.; Jaureguiulloa, E.; Adamo, K.B.; Taylor, J.L.; Janssen, I.; Tremblay, M.S. Adiposity and physical activity among children in countries at different stages of the physical activity transition: Canada, Mexico and Kenya. *Afr. J. Phys. Health Educ. Recreat. Dance* **2013**, *19*, 132–142.
88. Heroux, M.; Onywera, V.O.; Tremblay, M.S.; Adamo, K.B.; Taylor, J.L.; Ulloa, E.J.; Janssen, I. The relation between aerobic fitness, muscular fitness, and obesity in children from three countries at different stages of the physical activity transition. *ISRN Obesity* **2013**, *2013*, doi:10.1155/2013/134835.
89. Moher, D.; Liberati, A.; Tetzlaff, J.; Altman, D.G.; The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med.* **2009**, *6*, doi:10.1371/journal.pmed.1000097.
90. Freedson, P.S.; Melanson, E.; Sirard, J. Calibration of the Computer Science and Applications, Inc. accelerometer. *Med. Sci. Sport. Exercise* **1998**, *30*, 777–781.
91. Puyau, M.R.; Adolph, A.L.; Firoz, A.V.; Butte, N.F. Validation and calibration of physical activity monitors in children. *Obes. Res.* **2002**, *10*, 150–157.
92. Adamo, K.B.; Prince, S.A.; Tricco, A.C.; Connor-Gorber, S.; Tremblay, M.S. A comparison of indirect versus direct measures for assessing physical activity in the pediatric population: A systematic review. *Int. J. Pediatr. Obes.* **2009**, *4*, 2–27.
93. Armstrong, N.; Bray, S. Physical activity patterns defined by continuous heart rate monitoring. *Arch. Dis. Child.* **1991**, *66*, 245–247.
94. Bailey, R.; Olson, J.; Pepper, S.L.; Porszasz, J.; Barstow, T.J.; Cooper, D.M. The level and tempo of children’s physical activities: An observational study. *Med. Sci. Sports Exerc.* **1995**, *27*, 1033–1041.
95. LeBlanc, A.; Janssen, I. Difference between self-reported and accelerometer measured moderate-to-vigorous physical activity in youth. *Pediatr. Exerc. Sci.* **2010**, *22*, 523–534.
96. Shephard, R.J. Limits to the measurement of habitual physical activity by questionnaires. *Br. J. Sport. Med.* **2003**, *37*, 197–206.
97. Wareham, N.J. Commentary: Measuring physical activity in Sub-Saharan Africa. *Int. J. Epidemiol.* **2001**, *30*, 1369–1370.
98. Hossain, P.; Kawar, B.; El Nahas, M. Obesity and diabetes in the developing world: A growing challenge. *NEJM* **2007**, *356*, 213–215.