

Supplementary Materials: Epidemiology of Burns in Rural Bangladesh: An Update

Siran He, Olakunle Alonge *, Priyanka Agrawal, Shumona Sharmin, Irteja Islam, Saidur Rahman Mashreky and Shams El Arifeen

Table S1: Unadjusted and adjusted analysis of non-fatal burn injury by age group, sex and SES quintiles for children under 9 years of age.

	Unadjusted			Adjusted		
	OR	95% CI	P value	OR	95% CI	P value
Age group						
<1 yr	Reference group			Reference group		
1–4 yrs	4.36	3.37–5.63	<0.001	4.31	3.34–5.58	<0.001
5–9 yrs	1.41	1.09–1.85	<0.001	1.41	1.08–1.83	0.011
Sex						
Male	Reference group			Reference group		
Female	1.00	0.91–1.1	0.981	1.00	0.91–1.09	0.980
SES quintiles						
Lowest	Reference group			Reference group		
Low	0.85	0.74–0.97	0.018	0.86	0.75–0.99	0.041
Middle	0.73	0.63–0.84	<0.001	0.74	0.64–0.85	<0.001
High	0.74	0.64–0.84	<0.001	0.74	0.64–0.85	<0.001
Highest	0.67	0.58–0.78	<0.001	0.66	0.57–0.76	<0.001



© 2017 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).