

Supplementary table: Analysis results for the other factors.

Table S1. Age (years).

| Baseline difference between groups | Interaction between factors and visits | | ≤35 (n=20) | | | | | | >35 (n=18) | | | | | | | |
|------------------------------------|--|-------|------------|-------|-------------------|------|---------------------------|------|---------------------------|-------|-------------------|------|---------------------------|------|---------------------------|-------|
| | | | Baseline | | 6-month follow-up | | Difference between visits | | Baseline | | 6-month follow-up | | Difference between visits | | | |
| | Mann-Whitney U | p | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P |
| PCL: total scores | -1.00 | 0.318 | 2.90 | 0.097 | 3.40 | 4.64 | 1.60 | 3.02 | -2.19 | 0.028 | 1.72 | 2.40 | 1.44 | 1.98 | -0.81 | 0.416 |
| PCL: re-experience | -0.45 | 0.656 | 0.46 | 0.503 | 1.10 | 1.45 | 0.45 | 1.00 | -2.21 | 0.027 | 0.83 | 1.10 | 0.44 | 0.78 | -1.31 | 0.191 |
| PCL: avoidance | -1.28 | 0.201 | 3.78 | 0.060 | 1.15 | 1.93 | 0.35 | 1.35 | -1.78 | 0.075 | 0.39 | 0.78 | 0.50 | 0.99 | -1.00 | 0.317 |
| PCL: hyperarousal | -1.29 | 0.198 | 1.70 | 0.200 | 1.15 | 1.63 | 0.80 | 1.40 | -1.40 | 0.161 | 0.50 | 0.99 | 0.50 | 0.99 | 0.00 | 1.000 |

Table S2. Marital status.

| Baseline difference between groups | Interaction between factors and visits | | Marriage (n=32) | | | | | | Other than marriage (n=6) | | | | | | | |
|------------------------------------|--|-------|-----------------|-------|-------------------|------|---------------------------|------|---------------------------|-------|-------------------|------|---------------------------|------|---------------------------|-------|
| | | | Baseline | | 6-month follow-up | | Difference between visits | | Baseline | | 6-month follow-up | | Difference between visits | | | |
| | Mann-Whitney U | p | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P |
| PCL: total scores | -0.34 | 0.737 | 2.34 | 0.135 | 2.31 | 2.93 | 1.53 | 2.54 | -1.80 | 0.073 | 4.17 | 7.05 | 1.50 | 2.81 | -1.34 | 0.180 |
| PCL: re-experience | -0.09 | 0.931 | 1.15 | 0.290 | 0.91 | 1.09 | 0.47 | 0.92 | -2.08 | 0.038 | 1.33 | 2.16 | 0.33 | 0.82 | -1.41 | 0.157 |
| PCL: avoidance | -0.17 | 0.865 | 3.13 | 0.085 | 0.69 | 1.20 | 0.50 | 1.27 | -0.85 | 0.398 | 1.33 | 2.80 | 0.00 | 0.00 | -1.34 | 0.180 |
| PCL: hyperarousal | -0.35 | 0.728 | 0.22 | 0.639 | 0.72 | 1.14 | 0.56 | 1.01 | -0.91 | 0.366 | 1.50 | 2.35 | 1.17 | 2.04 | -1.00 | 0.317 |

Table S3. Years of working experience.

| ≤13 (n=19) | >14 (n=19) |
|------------|------------|
|------------|------------|

| Baseline difference between groups | Interaction between factors and visits | | | | | | | | | | | | Baseline 6-month follow-up | | | | Difference between visits | | | | Baseline 6-month follow-up | | | | Difference between visits | | | |
|------------------------------------|--|-------|------|-------|------|------|------|------|---------------------------|-------|------|------|----------------------------|------|---------------------------|-------|---------------------------|--|--|--|----------------------------|--|--|--|---------------------------|--|--|--|
| | Mann-Whitney U | P | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | | | | | | | | | | | | |
| PCL: total scores | -0.52 | 0.602 | 2.51 | 0.122 | 3.21 | 4.69 | 1.42 | 1.98 | -2.31 | 0.021 | 2.00 | 2.62 | 1.63 | 3.06 | -0.84 | 0.404 | | | | | | | | | | | | |
| PCL: re-experience | -0.27 | 0.787 | 2.86 | 0.099 | 1.11 | 1.52 | 0.26 | 0.56 | -2.41 | 0.016 | 0.84 | 1.01 | 0.63 | 1.12 | -0.92 | 0.357 | | | | | | | | | | | | |
| PCL: avoidance | -0.44 | 0.658 | 1.18 | 0.284 | 0.89 | 1.79 | 0.26 | 0.73 | -1.51 | 0.131 | 0.68 | 1.25 | 0.58 | 1.50 | -0.43 | 0.671 | | | | | | | | | | | | |
| PCL: hyperarousal | -1.52 | 0.128 | 0.95 | 0.337 | 1.21 | 1.65 | 0.89 | 1.41 | -1.19 | 0.236 | 0.47 | 0.96 | 0.42 | 0.96 | -0.58 | 0.564 | | | | | | | | | | | | |

Table S4. Personality: Anxiety.

| Baseline difference between groups | Interaction between factors and visits | | | | | | | | | | | | No (n=13) | | | | Yes (n=25) | | | |
|------------------------------------|--|-------|------|-------|------|------|------|------|---------------------------|-------|------|------|-----------|------|---------------------------|-------|------------|--|--|--|
| | Mann-Whitney U | P | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | | | | |
| PCL: total scores | -2.04 | 0.042 | 2.19 | 0.148 | 1.15 | 2.19 | 1.00 | 1.58 | -0.55 | 0.581 | 3.36 | 4.25 | 1.80 | 2.92 | -2.22 | 0.026 | | | | |
| PCL: re-experience | -2.19 | 0.028 | 1.25 | 0.271 | 0.38 | 0.87 | 0.15 | 0.38 | -1.13 | 0.257 | 1.28 | 1.37 | 0.60 | 1.04 | -2.27 | 0.023 | | | | |
| PCL: avoidance | -1.23 | 0.217 | 0.74 | 0.394 | 0.23 | 0.44 | 0.15 | 0.38 | -1.00 | 0.317 | 1.08 | 1.80 | 0.56 | 1.42 | -1.24 | 0.215 | | | | |
| PCL: hyperarousal | -1.12 | 0.261 | 3.47 | 0.071 | 0.54 | 1.05 | 0.69 | 1.11 | -1.00 | 0.317 | 1.00 | 1.53 | 0.64 | 1.29 | -2.06 | 0.039 | | | | |

Table S5. Personality: Perfectionism.

| Baseline difference between groups | Interaction between factors and visits | | | | | | | | | | | | No (n=29) | | | | Yes (n=9) | | | |
|------------------------------------|--|-------|------|-------|------|------|------|------|---------------------------|-------|------|------|-----------|------|---------------------------|-------|-----------|--|--|--|
| | Mann-Whitney U | P | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | | | | |
| PCL: total scores | -3.37 | 0.001 | 4.04 | 0.052 | 1.41 | 2.51 | 0.83 | 1.61 | -1.31 | 0.190 | 6.44 | 4.77 | 3.78 | 3.67 | -1.85 | 0.064 | | | | |

| | | | | | | | | | | | | | | | | |
|--------------------|-------|-------|------|-------|------|------|------|------|-------|-------|------|------|------|------|-------|-------|
| PCL: re-experience | -3.44 | 0.001 | 6.31 | 0.017 | 0.55 | 0.91 | 0.28 | 0.65 | -1.47 | 0.142 | 2.33 | 1.41 | 1.00 | 1.32 | -2.06 | 0.040 |
| PCL: avoidance | -2.67 | 0.008 | 0.88 | 0.355 | 0.41 | 0.95 | 0.17 | 0.60 | -1.51 | 0.131 | 2.00 | 2.35 | 1.22 | 2.05 | -0.53 | 0.596 |
| PCL: hyperarousal | -3.13 | 0.002 | 2.43 | 0.128 | 0.45 | 0.99 | 0.38 | 0.82 | 0.00 | 1.000 | 2.11 | 1.76 | 1.56 | 1.81 | -1.89 | 0.059 |

Table S6. Personality: Introvert/socially inactive.

| Baseline difference between groups | Interaction between factors and visits | | No (n=24) | | | | | | Yes (n=14) | | | | | | | |
|------------------------------------|--|-------|-----------|-------------------|---------------------------|------|------|------|---------------------------|-------------------|---------------------------|------|------|------|---------------------------|-------|
| | | | Baseline | 6-month follow-up | Difference between visits | | | | Baseline | 6-month follow-up | Difference between visits | | | | | |
| | Mann-Whitney U | p | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P |
| PCL: total scores | -0.24 | 0.812 | 0.12 | 0.735 | 2.25 | 0.60 | 1.29 | 0.36 | -1.81 | 0.071 | 3.21 | 1.33 | 1.93 | 0.95 | -1.53 | 0.125 |
| PCL: re-experience | -0.82 | 0.410 | 0.56 | 0.461 | 0.83 | 0.24 | 0.42 | 0.18 | -1.63 | 0.102 | 1.21 | 0.39 | 0.50 | 0.25 | -1.98 | 0.047 |
| PCL: avoidance | -0.79 | 0.429 | 0.03 | 0.853 | 0.75 | 0.24 | 0.42 | 0.18 | -1.29 | 0.196 | 0.86 | 0.55 | 0.43 | 0.43 | -0.54 | 0.593 |
| PCL: hyperarousal | -0.33 | 0.739 | 0.05 | 0.819 | 0.67 | 0.22 | 0.46 | 0.18 | -0.91 | 0.366 | 1.14 | 0.48 | 1.00 | 0.43 | -1.00 | 0.317 |

Table S7. Number of previous deployments to disaster field operations.

| Baseline difference between groups | Interaction between factors and visits | | ≥ 1 (n=20) | | | | | | 0 (n=18) | | | | | | | |
|------------------------------------|--|-------|------------|-------------------|---------------------------|------|------|------|---------------------------|-------------------|---------------------------|------|------|------|---------------------------|-------|
| | | | Baseline | 6-month follow-up | Difference between visits | | | | Baseline | 6-month follow-up | Difference between visits | | | | | |
| | Mann-Whitney U | p | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P |
| PCL: total scores | -0.43 | 0.667 | 0.41 | 0.528 | 2.80 | 3.49 | 2.00 | 3.03 | -1.01 | 0.310 | 2.39 | 4.20 | 1.00 | 1.81 | -2.25 | 0.024 |
| PCL: re-experience | -1.16 | 0.245 | 0.02 | 0.887 | 1.15 | 1.23 | 0.65 | 1.09 | -1.63 | 0.102 | 0.78 | 1.35 | 0.22 | 0.55 | -1.98 | 0.047 |
| PCL: avoidance | -0.69 | 0.489 | 0.90 | 0.350 | 0.90 | 1.41 | 0.75 | 1.55 | -0.26 | 0.796 | 0.67 | 1.68 | 0.06 | 0.24 | -1.84 | 0.066 |
| PCL: hyperarousal | -0.73 | 0.466 | 0.07 | 0.794 | 0.75 | 1.33 | 0.60 | 1.10 | -0.38 | 0.705 | 0.94 | 1.47 | 0.72 | 1.36 | -1.41 | 0.157 |

Table S8. How many days after the earthquake did you arrive at the field?

| Baseline difference between groups | Interaction between factors and visits | ≥2 (n=7) | | | | | | | 1 (n=31) | | | | | | | |
|------------------------------------|--|----------|-------------------|---------------------------|------|------|----------|---------------------------|---------------------------|-------|------|------|------|---------------------------|-------|-------|
| | | Baseline | 6-month follow-up | Difference between visits | | | Baseline | 6-month follow-up | Difference between visits | | | Mean | SD | Wilcoxon Signed-Rank Test | P | |
| Mann-Whitney U | P | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | |
| PCL: total scores | -0.49 | 0.621 | 0.04 | 0.833 | 3.14 | 6.23 | 1.86 | 3.18 | -0.54 | 0.593 | 2.48 | 3.15 | 1.45 | 2.43 | -2.28 | 0.023 |
| PCL: re-experience | -0.10 | 0.918 | 0.06 | 0.812 | 1.14 | 1.86 | 0.71 | 1.25 | -0.82 | 0.414 | 0.94 | 1.15 | 0.39 | 0.80 | -2.56 | 0.010 |
| PCL: avoidance | -0.07 | 0.945 | 0.91 | 0.346 | 1.29 | 2.63 | 0.43 | 1.13 | -0.45 | 0.655 | 0.68 | 1.19 | 0.42 | 1.20 | -1.21 | 0.227 |
| PCL: hyperarousal | -1.03 | 0.305 | 0.41 | 0.525 | 0.71 | 1.89 | 0.71 | 1.89 | 0.00 | 1.000 | 0.87 | 1.28 | 0.65 | 1.05 | -1.31 | 0.190 |

Table S9. How many hours did you participate in the field operation for the 2016 Taiwan earthquake?

| Baseline difference between groups | Interaction between factors and visits | ≤60 (n=17) | | | | | | | >60 (n=20) | | | | | | | |
|------------------------------------|--|------------|-------------------|---------------------------|------|------|----------|---------------------------|---------------------------|-------|------|------|------|---------------------------|-------|-------|
| | | Baseline | 6-month follow-up | Difference between visits | | | Baseline | 6-month follow-up | Difference between visits | | | Mean | SD | Wilcoxon Signed-Rank Test | P | |
| Mann-Whitney U | P | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | |
| PCL: total scores | -0.05 | 0.962 | 0.26 | 0.612 | 2.88 | 4.55 | 2.12 | 3.35 | -0.81 | 0.419 | 2.35 | 3.23 | 1.10 | 1.59 | -2.22 | 0.026 |
| PCL: re-experience | -0.40 | 0.692 | 0.36 | 0.552 | 1.12 | 1.45 | 0.71 | 1.21 | -1.21 | 0.227 | 0.90 | 1.17 | 0.25 | 0.44 | -2.36 | 0.018 |
| PCL: avoidance | -0.93 | 0.355 | 0.01 | 0.911 | 1.12 | 1.96 | 0.82 | 1.67 | -0.11 | 0.916 | 0.45 | 1.00 | 0.10 | 0.31 | -1.89 | 0.059 |
| PCL: hyperarousal | -1.18 | 0.237 | 0.48 | 0.495 | 0.65 | 1.41 | 0.59 | 1.37 | -0.58 | 0.564 | 1.00 | 1.41 | 0.75 | 1.12 | -0.96 | 0.336 |

Table S10. Tasks: Managing injured patients.

| Baseline difference between groups | Interaction between factors and visits | No (n=11) | | | | | | | Yes (n=27) | | | | | | |
|------------------------------------|--|-----------|-------------------|---------------------------|----|------|----------|---------------------------|---------------------------|------|----|------|----|---------------------------|---|
| | | Baseline | 6-month follow-up | Difference between visits | | | Baseline | 6-month follow-up | Difference between visits | | | Mean | SD | Wilcoxon Signed-Rank Test | P |
| Mann-Whitney U | P | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P |
| | | | | | | | | | | | | | | | |

| | | | | | | | | | | | | | | | | |
|--------------------|-------|-------|------|-------|------|------|------|------|-------|-------|------|------|------|------|-------|-------|
| PCL: total scores | -2.67 | 0.008 | 1.27 | 0.267 | 0.36 | 0.20 | 0.09 | 0.09 | -1.13 | 0.257 | 3.52 | 0.80 | 2.11 | 0.54 | -2.06 | 0.039 |
| PCL: re-experience | -1.84 | 0.066 | 0.29 | 0.594 | 0.36 | 0.20 | 0.00 | 0.00 | -1.63 | 0.102 | 1.22 | 0.27 | 0.63 | 0.19 | -2.12 | 0.034 |
| PCL: avoidance | -2.58 | 0.010 | 1.48 | 0.232 | 0.00 | 0.00 | 0.09 | 0.09 | -1.00 | 0.317 | 1.11 | 0.33 | 0.56 | 0.26 | -1.55 | 0.121 |
| PCL: hyperarousal | -2.87 | 0.004 | 0.75 | 0.392 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 1.000 | 1.19 | 0.29 | 0.93 | 0.26 | -1.31 | 0.190 |

Table S11. Tasks: Managing dead people.

| Baseline difference between groups | Interaction between factors and visits | | | | | | | | | | No (n=11) | | | | | | Yes (n=26) | | | | | |
|------------------------------------|--|-------|------|-------|----------|------|-------------------|------|---------------------------|-------|-----------|------|-------------------|------|---------------------------|-------|---------------------------|--|--|--|--|--|
| | | | | | Baseline | | 6-month follow-up | | | | Baseline | | 6-month follow-up | | | | Difference between visits | | | | | |
| | Mann-Whitney U | p | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | | | | | | |
| PCL: total scores | -1.66 | 0.098 | 0.75 | 0.391 | 1.09 | 0.58 | 0.64 | 0.36 | -1.29 | 0.197 | 3.22 | 0.81 | 1.89 | 0.55 | -1.93 | 0.053 | | | | | | |
| PCL: re-experience | -1.45 | 0.146 | 0.71 | 0.405 | 0.55 | 0.31 | 0.27 | 0.27 | -1.34 | 0.180 | 1.15 | 0.26 | 0.52 | 0.17 | -2.22 | 0.026 | | | | | | |
| PCL: avoidance | -1.15 | 0.249 | 0.00 | 0.990 | 0.36 | 0.28 | 0.00 | 0.00 | -1.34 | 0.180 | 0.96 | 0.33 | 0.59 | 0.26 | -0.85 | 0.398 | | | | | | |
| PCL: hyperarousal | -2.16 | 0.030 | 3.16 | 0.084 | 0.18 | 0.18 | 0.36 | 0.28 | -1.41 | 0.157 | 1.11 | 0.29 | 0.78 | 0.25 | -1.93 | 0.054 | | | | | | |

Table S12. Tasks: Managing dead children (under 8 years of age).

| Baseline difference between groups | Interaction between factors and visits | | | | | | | | | | No (n=22) | | | | | | Yes (n=16) | | | | | |
|------------------------------------|--|-------|------|-------|----------|------|-------------------|------|---------------------------|-------|-----------|------|-------------------|------|---------------------------|-------|---------------------------|--|--|--|--|--|
| | | | | | Baseline | | 6-month follow-up | | | | Baseline | | 6-month follow-up | | | | Difference between visits | | | | | |
| | Mann-Whitney U | p | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | | | | | | |
| PCL: total scores | -1.18 | 0.238 | 0.30 | 0.588 | 2.18 | 0.86 | 1.32 | 0.62 | -1.75 | 0.080 | 3.19 | 0.87 | 1.81 | 0.49 | -1.54 | 0.124 | | | | | | |
| PCL: re-experience | -0.64 | 0.520 | 0.03 | 0.874 | 0.86 | 0.27 | 0.36 | 0.20 | -2.33 | 0.020 | 1.13 | 0.33 | 0.56 | 0.20 | -1.44 | 0.150 | | | | | | |
| PCL: avoidance | -1.19 | 0.236 | 0.00 | 0.982 | 0.73 | 0.37 | 0.36 | 0.28 | -0.81 | 0.416 | 0.88 | 0.30 | 0.50 | 0.26 | -1.19 | 0.236 | | | | | | |
| PCL: hyperarousal | -1.85 | 0.064 | 2.67 | 0.111 | 0.59 | 0.28 | 0.59 | 0.28 | 0.00 | 1.000 | 1.19 | 0.37 | 0.75 | 0.27 | -1.51 | 0.131 | | | | | | |

Table S13. Tasks: Managing dead persons who were apparently pregnant.

| Baseline difference between groups | Interaction between factors and visits | | No (n=33) | | | | | | Yes (n=5) | | | | | | | |
|------------------------------------|--|-------|-----------|-------|-------------------|------|---------------------------|------|---------------------------|-------|-------------------|------|---------------------------|------|---------------------------|-------|
| | | | Baseline | | 6-month follow-up | | Difference between visits | | Baseline | | 6-month follow-up | | Difference between visits | | | |
| | Mann-Whitney U | p | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P |
| PCL: total scores | -2.47 | 0.013 | 4.23 | 0.047 | 2.12 | 0.63 | 1.39 | 0.45 | -1.58 | 0.114 | 5.80 | 1.62 | 2.40 | 1.03 | -2.04 | 0.041 |
| PCL: re-experience | -2.66 | 0.008 | 5.33 | 0.027 | 0.79 | 0.22 | 0.42 | 0.16 | -1.75 | 0.081 | 2.20 | 0.37 | 0.60 | 0.24 | -2.06 | 0.039 |
| PCL: avoidance | -1.42 | 0.157 | 1.03 | 0.318 | 0.70 | 0.26 | 0.42 | 0.22 | -0.85 | 0.398 | 1.40 | 0.75 | 0.40 | 0.24 | -1.34 | 0.180 |
| PCL: hyperarousal | -2.35 | 0.019 | 3.34 | 0.076 | 0.64 | 0.22 | 0.55 | 0.20 | -0.91 | 0.366 | 2.20 | 0.73 | 1.40 | 0.68 | -1.00 | 0.317 |

Table S14. Tasks: Managing people whose families had died in the disaster.

| Baseline difference between groups | Interaction between factors and visits | | No (n=12) | | | | | | Yes (n=26) | | | | | | | |
|------------------------------------|--|-------|-----------|-------|-------------------|------|---------------------------|------|---------------------------|-------|-------------------|------|---------------------------|------|---------------------------|-------|
| | | | Baseline | | 6-month follow-up | | Difference between visits | | Baseline | | 6-month follow-up | | Difference between visits | | | |
| | Mann-Whitney U | p | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P |
| PCL: total scores | -1.02 | 0.307 | 1.53 | 0.223 | 1.33 | 0.54 | 1.08 | 0.56 | -0.64 | 0.524 | 3.19 | 0.85 | 1.73 | 0.55 | -2.24 | 0.025 |
| PCL: re-experience | -0.80 | 0.422 | 0.15 | 0.703 | 0.67 | 0.28 | 0.25 | 0.25 | -1.09 | 0.276 | 1.12 | 0.27 | 0.54 | 0.18 | -2.30 | 0.021 |
| PCL: avoidance | -0.72 | 0.469 | 2.32 | 0.136 | 0.42 | 0.23 | 0.58 | 0.34 | -1.00 | 0.317 | 0.96 | 0.34 | 0.35 | 0.23 | -1.78 | 0.075 |
| PCL: hyperarousal | -1.82 | 0.069 | 0.85 | 0.362 | 0.25 | 0.18 | 0.25 | 0.25 | 0.00 | 1.000 | 1.12 | 0.31 | 0.85 | 0.26 | -1.40 | 0.161 |

Table S15. Tasks: Managing victims' families who were emotionally distraught.

| No (n=17) | Yes (n=21) |
|-----------|------------|
|-----------|------------|

| Baseline difference between groups | Interaction between factors and visits | | | | | | | | | | | | Interaction between baseline and 6-month follow-up | | | | | | | | | | | | | | | | | | | | | | |
|------------------------------------|--|-------|------|-------|------|------|-------------------|---------------------------|-------|-------|------|------|--|---------------------------|-------|-------|--|--|----------|--|--|--|--|--|-------------------|--|--|--|--|--|---------------------------|--|--|--|--|
| | Baseline | | | | | | 6-month follow-up | | | | | | Difference between visits | | | | | | Baseline | | | | | | 6-month follow-up | | | | | | Difference between visits | | | | |
| Mann-Whitney U | P | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | | | | | | | | | | | | | | | | | | | | |
| PCL: total scores | -2.57 | 0.010 | 4.39 | 0.043 | 0.94 | 0.41 | 0.88 | 0.41 | -0.17 | 0.864 | 3.95 | 0.98 | 2.05 | 0.66 | -2.49 | 0.013 | | | | | | | | | | | | | | | | | | | |
| PCL: re-experience | -2.19 | 0.029 | 1.92 | 0.174 | 0.47 | 0.21 | 0.24 | 0.18 | -0.85 | 0.395 | 1.38 | 0.31 | 0.62 | 0.21 | -2.49 | 0.013 | | | | | | | | | | | | | | | | | | | |
| PCL: avoidance | -1.75 | 0.081 | 3.46 | 0.071 | 0.29 | 0.17 | 0.41 | 0.24 | -1.00 | 0.317 | 1.19 | 0.41 | 0.43 | 0.29 | -1.78 | 0.075 | | | | | | | | | | | | | | | | | | | |
| PCL: hyperarousal | -2.89 | 0.004 | 2.74 | 0.107 | 0.18 | 0.13 | 0.24 | 0.18 | -0.58 | 0.564 | 1.38 | 0.36 | 1.00 | 0.31 | -1.73 | 0.084 | | | | | | | | | | | | | | | | | | | |

Table S16. Feelings of guilt and self-blame.

| Baseline difference between groups | Interaction between factors and visits | | | | | | | | | | | | Interaction between baseline and 6-month follow-up | | | | | | | | | | | | |
|------------------------------------|--|-------|------|-------|------|------|-----------|---------------------------|-------|-------|------|------|--|---------------------------|-------|----------------|---|---|-----------|------|----|------|----|---------------------------|---|
| | No (n=27) | | | | | | Yes (n=9) | | | | | | No (n=27) | | | | | | Yes (n=9) | | | | | | |
| Mann-Whitney U | P | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P | Mann-Whitney U | P | F | p | Mean | SD | Mean | SD | Wilcoxon Signed-Rank Test | P |
| PCL: total scores | -2.76 | 0.006 | 3.89 | 0.057 | 1.48 | 2.19 | 0.93 | 1.73 | -1.96 | 0.051 | 6.11 | 5.64 | 3.44 | 3.81 | -1.12 | 0.261 | | | | | | | | | |
| PCL: re-experience | -1.75 | 0.080 | 1.08 | 0.307 | 0.74 | 1.06 | 0.33 | 0.83 | -1.93 | 0.054 | 1.78 | 1.72 | 0.89 | 1.05 | -1.30 | 0.194 | | | | | | | | | |
| PCL: avoidance | -2.78 | 0.005 | 5.06 | 0.031 | 0.37 | 0.84 | 0.30 | 0.82 | -0.41 | 0.680 | 2.11 | 2.42 | 0.78 | 1.99 | -1.38 | 0.168 | | | | | | | | | |
| PCL: hyperarousal | -3.30 | 0.001 | 1.31 | 0.261 | 0.37 | 0.84 | 0.30 | 0.82 | -1.00 | 0.317 | 2.22 | 1.86 | 1.78 | 1.64 | -0.74 | 0.461 | | | | | | | | | |