

Supplementary

Diatomite Photonic Crystals for Facile On-Chip Chromatography and Sensing of Harmful Ingredients from Food

Xianming Kong ¹, Qian Yu ^{1,*}, Erwen Li ², Rui Wang ¹, Qing Liu ³ and Alan Wang ^{2,*}

¹ College of Chemistry, Chemical Engineering and Environment Engineering, Liaoning Shihua University, Fushun 113001, Liaoning, China; xmkong@lnpu.edu.cn (X.K.); rwang@lnpu.edu.cn (R.W.)

² School of Electrical Engineering and Computer Science, Oregon State University, Corvallis, OR 97331, USA; lie@oregonstate.edu

³ Key Laboratory of Low Carbon Energy and Chemical Engineering, College of Chemical and Environmental Engineering, Shandong University of Science and Technology, Qingdao 266590, Shandong, China; qliu@sdust.edu.cn

* Correspondence: qyu@lnpu.edu.cn (Q.Y.); wang@eecs.oregonstate.edu (A.W.); Tel.: +1-541-737-4247 (A.W.)

Received: 6 March 2018; Accepted: 29 March 2018; Published: 31 March 2018




Figure S1. The UV-VIS absorption spectrum of the prepared Au colloids.




Figure S2. Microscopic image of the cross-section of the commercial silica gel TLC chip.