



## Article Recyclable Polymer-Supported N-Hydroxyphthalimide Catalysts for Selective Oxidation of Pullulan

## Madalina Elena Culica <sup>1</sup>, Kornela Kasperczyk <sup>2</sup>, Raluca Ioana Baron <sup>1</sup>, Gabriela Biliuta <sup>1</sup>, Ana Maria Macsim <sup>1</sup>, Andrada Lazea-Stoyanova <sup>3</sup>, Beata Orlinska <sup>2</sup> and Sergiu Coseri <sup>1,\*</sup>

- <sup>1</sup> "Petru Poni" Institute of Macromolecular Chemistry of Romanian Academy, 41 A, Gr. Ghica Voda Alley, Iasi 700487, Romania; culica.madalina@icmpp.ro (M.E.C.); baron.raluca@icmpp.ro (R.I.B.); biliuta.gabriela@icmpp.ro (G.B.); ana.iurascu@icmpp.ro (A.M.M.)
- <sup>2</sup> Department of Chemical Organic Technology and Petrochemistry, Silesian University of Technology, Krzywoustego 4, Gliwice 44 100, Poland; kornela.kasperczyk@gmail.com (K.K.); Beata.Orlinska@polsl.pl (B.O.)
- <sup>3</sup> National Institute for Lasers, Plasma and Radiation Physics, 409 Atomistilor Street, Magurele 77125, Romania; andrada@infim.ro
- \* Correspondence: coseris@icmpp.ro ; Tel.: +40-232-217-454; Fax: +40-232-2112-99

Received: 26 August 2019; Accepted: 30 October 2019; Published: date



Figure S1. The <sup>13</sup>C-NMR spectra of PI - APS.











Figure S6. The <sup>13</sup>C-NMR spectra of PII - BMPS.