

Table S1. The effect of basic media on rooting rates in *K. paniculata*.

Basic Medium	IBA (mg L ⁻¹)	Frequency of Rhizogenesis + SE (%)
DKW	0.5	0.00 + 0.00 ^d
DKW	1.0	0.00 + 0.00 ^d
DKW	1.5	0.00 + 0.00 ^d
DKW	2.0	3.33 + 0.06 ^d
DKW	2.5	6.67 + 0.06 ^{cd}
1/2DKW	0.5	7.33 + 0.06 ^{cd}
1/2DKW	1.0	14.00 + 0.05 ^{bcd}
1/2DKW	1.5	16.67 + 0.12 ^{bcd}
1/2DKW	2.0	16.00 + 0.04 ^{bcd}
1/2DKW	2.5	13.00 + 0.06 ^{bcd}
1/4DKW	0.5	24.00 + 0.02 ^{bc}
1/4DKW	1.0	47.50 + 0.08 ^a
1/4DKW	1.5	52.00 + 0.12 ^a
1/4DKW	2.0	26.75 + 0.02 ^b
1/4DKW	2.5	28.00 + 0.02 ^b

Date represented mean + SE of all replicates, each with 9 cultures. Means having the same letter were not significantly different by Tukey's multiple range test ($p \leq 0.05$).