
Supplementary Information

In Vitro Evaluation of Anti-RIFT Valley Fever Virus, Antioxidant and Anti-Inflammatory Activity of South African Medicinal Plant Extracts


Garland K. More ^{1,*}, Raymond T. Makola ^{2,3} and Gerhard Prinsloo, ¹


¹ College of Agriculture and Environmental Sciences, University of South Africa, Private Bag X6, Florida 1710, South Africa; moregk@unisa.ac.za (G.K.M.); prinsg@unisa.ac.za (G.P.)

² Department of Biochemistry Microbiology and Biotechnology, School of Molecular and Life Science, University of Limpopo (Turfloop Campus) Sovenga 0727; makolaraymond4@gmail.com (R.T.M.)

³ National institution of communicable diseases, Special Viral Pathogen/ Arbovirus Unit, 1 Modderfontein Rd, Sandringham, Johannesburg 2192

* Correspondence: moregk@unisa.ac.za









Figure S1: Lethal concentration (LC₅₀ = µg/mL) of the MTT assays following 48 hours treatment with twenty plant extracts well known for their antiviral activity.

A






Figure S2: Effective concentration (EC₅₀ = µg/mL) of the DPPH (A) and ABTS (B) assays following 48 hours treatment with eight plant extracts well known for their antiviral activity.