

*Article Supplementary Materials*


**Table S1.** Nutrient contents of as-is and dried fine solids, other blend ingredients, derived plant foods, and the market alternative on a dry weight basis.

	As-is Fine Solids	Dried Fine Solids <sup>a</sup>	Potting Mix	Dried Distiller's Grain & Whey Permeate	Biochar	Plant Food A	Plant Food A1	Plant Food A2	Plant Food B	Market Alter- native
Total N (g kg <sup>-1</sup> )	50.2 <sup>c</sup>	19.5 <sup>c</sup>	5.7 <sup>b</sup>	39.8 <sup>b</sup>	17.8 <sup>b</sup>	41.6 <sup>c</sup>	41.6 <sup>c</sup>	40.4 <sup>c</sup>	40.1 <sup>c</sup>	44.9 <sup>b</sup>
Organic N (g kg <sup>-1</sup> )	38.8 <sup>e</sup>	17.7 <sup>e</sup>	5.7 <sup>d</sup>	39.7 <sup>d</sup>	17.5 <sup>d</sup>	40.1 <sup>e</sup>	40.5 <sup>e</sup>	39.7 <sup>e</sup>	38.8 <sup>e</sup>	NA
NH <sub>4</sub> -N (g kg <sup>-1</sup> )	11.3	1.8	0.0	0.1	0.3	1.5	1.1	0.7	1.3	NA
NO <sub>3</sub> -N (g kg <sup>-1</sup> )	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	NA
Total P (g kg <sup>-1</sup> )	18.3	17.1	0.6	5.3	8.3	16.0	17.1	17.4	16.1	19.3
Neutral NH <sub>4</sub> Citrate P (g kg <sup>-1</sup> )	19.4	19.5	0.4	5.3	6.8	17.3	16.5	16.0	16.9	15.7
2% Citric Acid P (g kg <sup>-1</sup> )	15.3	11.9	NA	NA	NA	11.8	10.8	9.7	11.8	9.9
Olsen P (g kg <sup>-1</sup> )	0.8	1.8	NA	NA	NA	1.9	1.9	2.1	1.6	0.7
Water Extractable P (g kg <sup>-1</sup> )	1.9	3.0	NA	NA	NA	3.2	3.4	3.4	3.5	2.3
Total K (g kg <sup>-1</sup> )	14.7	12.9	1.2	7.6	31.0	13.3	14.5	14.9	13.8	19.4
Neutral NH <sub>4</sub> Citrate K (g kg <sup>-1</sup> )	18.6	16.9	1.5	8.4	32.5	16.9	16.2	15.8	16.1	21.4
Total C (g kg <sup>-1</sup> )	405	383	329	470	550	386	413	424	388	NA
Total B (g kg <sup>-1</sup> )	0.02	0.02	0.03	0.00	0.04	0.02	0.02	0.02	0.02	0.00
Total Ca (g kg <sup>-1</sup> )	32	32	25	2	35	22	22	22	22	56
Total Cu (g kg <sup>-1</sup> )	0.64	0.66	0.04	0.00	0.03	0.67	0.70	0.69	0.68	0.04
Total Fe (g kg <sup>-1</sup> )	0.83	0.83	1.85	0.22	2.50	0.82	0.79	0.78	0.80	0.25
Total Mg (g kg <sup>-1</sup> )	12	12	5.1	1.6	15	12	13	12	12	3.5
Total Mn (g kg <sup>-1</sup> )	0.22	0.22	0.11	0.01	0.28	0.23	0.24	0.23	0.24	0.07
Total Na (g kg <sup>-1</sup> )	5.5	5.4	0.4	1.7	7.1	5.1	5.6	5.2	5.2	1.7
Total Z (g kg <sup>-1</sup> )	0.35	0.35	0.08	0.06	0.14	0.29	0.32	0.32	0.32	0.28

<sup>a</sup>Partially dried to ~45% total solids at 60°C; <sup>b</sup> Total N measured by combustion analysis; <sup>c</sup> Total N calculated as sum of total Kjeldahl N and NO<sub>3</sub>-N; <sup>d</sup> Organic N estimated as total N – (NH<sub>4</sub>-N + NO<sub>3</sub>-N); <sup>e</sup> Organic N is estimated as total Kjeldahl N – NH<sub>4</sub>-N.

**Table S2.** Bioassay germination rates, survival rates, root dry biomass, shoot dry biomass and total dry biomass by amendment and application rate (mean  $\pm$  1 SD). Groups share a letter if the difference in means was not statistically significant ( $P > 0.05$ ).  $n = 6$  trays of 16 seedlings for germination and survival and  $n = 24$  seedlings for root biomass, shoot biomass and total biomass.

Plant	Amendment	Application Rate	Germination (%)	Survival (%)	Root Biomass (mg dry)	Shoot Biomass (mg dry)	Total Biomass (mg dry)
tomato	control	0	93 $\pm$ 5 <sup>ab</sup>	98 $\pm$ 6 <sup>a</sup>	9 $\pm$ 2 <sup>d</sup>	17 $\pm$ 3 <sup>d</sup>	26 $\pm$ 5 <sup>d</sup>
tomato	Market Alternative	2	84 $\pm$ 7 <sup>b</sup>	89 $\pm$ 9 <sup>a</sup>	51 $\pm$ 11 <sup>a</sup>	139 $\pm$ 38 <sup>a</sup>	190 $\pm$ 47 <sup>a</sup>
tomato	Plant Food B	2	94 $\pm$ 4 <sup>ab</sup>	99 $\pm$ 3 <sup>a</sup>	23 $\pm$ 6 <sup>cd</sup>	63 $\pm$ 17 <sup>cd</sup>	86 $\pm$ 21 <sup>cd</sup>
tomato	Plant Food B	4	94 $\pm$ 0 <sup>ab</sup>	97 $\pm$ 6 <sup>a</sup>	30 $\pm$ 6 <sup>bc</sup>	90 $\pm$ 19 <sup>bc</sup>	120 $\pm$ 24 <sup>bc</sup>
tomato	Plant Food B	6	91 $\pm$ 5 <sup>ab</sup>	100 $\pm$ 0 <sup>a</sup>	38 $\pm$ 7 <sup>ab</sup>	115 $\pm$ 21 <sup>ab</sup>	152 $\pm$ 27 <sup>ab</sup>
tomato	Plant Food B	8	97 $\pm$ 5 <sup>a</sup>	96 $\pm$ 8 <sup>a</sup>	37 $\pm$ 9 <sup>ab</sup>	114 $\pm$ 27 <sup>ab</sup>	150 $\pm$ 35 <sup>ab</sup>
tomato	Plant Food B	10	96 $\pm$ 5 <sup>a</sup>	93 $\pm$ 0 <sup>a</sup>	37 $\pm$ 10 <sup>ab</sup>	110 $\pm$ 28 <sup>ab</sup>	147 $\pm$ 38 <sup>ab</sup>
tomato	Plant Food B	12	91 $\pm$ 3 <sup>ab</sup>	92 $\pm$ 9 <sup>a</sup>	33 $\pm$ 14 <sup>b</sup>	107 $\pm$ 44 <sup>ab</sup>	141 $\pm$ 57 <sup>ab</sup>
marigold	control	0	98 $\pm$ 3 <sup>a</sup>	100 $\pm$ 0 <sup>a</sup>	10 $\pm$ 2 <sup>c</sup>	17 $\pm$ 3 <sup>d</sup>	27 $\pm$ 5 <sup>d</sup>
marigold	Market Alternative	2	96 $\pm$ 5 <sup>a</sup>	80 $\pm$ 24 <sup>ab</sup>	51 $\pm$ 16 <sup>a</sup>	127 $\pm$ 38 <sup>a</sup>	178 $\pm$ 52 <sup>ab</sup>
marigold	Plant Food A	2	99 $\pm$ 3 <sup>a</sup>	100 $\pm$ 0 <sup>a</sup>	33 $\pm$ 8 <sup>b</sup>	69 $\pm$ 15 <sup>cd</sup>	102 $\pm$ 22 <sup>cd</sup>
marigold	Plant Food A	4	98 $\pm$ 3 <sup>a</sup>	97 $\pm$ 3 <sup>ab</sup>	44 $\pm$ 10 <sup>ab</sup>	99 $\pm$ 19 <sup>abc</sup>	143 $\pm$ 26 <sup>abc</sup>
marigold	Plant Food A	6	99 $\pm$ 3 <sup>a</sup>	92 $\pm$ 5 <sup>ab</sup>	38 $\pm$ 15 <sup>ab</sup>	88 $\pm$ 31 <sup>cb</sup>	125 $\pm$ 45 <sup>bc</sup>
marigold	Plant Food A	8	97 $\pm$ 5 <sup>a</sup>	82 $\pm$ 12 <sup>b</sup>	46 $\pm$ 13 <sup>ab</sup>	114 $\pm$ 29 <sup>ab</sup>	160 $\pm$ 40 <sup>ab</sup>
marigold	Plant Food A <sub>1</sub>	6	99 $\pm$ 3 <sup>a</sup>	90 $\pm$ 7 <sup>ab</sup>	48 $\pm$ 17 <sup>a</sup>	113 $\pm$ 33 <sup>a</sup>	161 $\pm$ 48 <sup>ab</sup>
marigold	Plant Food A <sub>2</sub>	6	100 $\pm$ 0 <sup>a</sup>	83 $\pm$ 14 <sup>ab</sup>	52 $\pm$ 12 <sup>a</sup>	121 $\pm$ 21 <sup>ab</sup>	174 $\pm$ 31 <sup>a</sup>



**Figure S1.** Disease suppression potential of as-is fine solids and the market alternative ( $n = 5$  per amendment). Negative values represented suppressive potential.