

Supplementary Materials: Dynamics and Drivers of Vegetation Phenology in Three-River Headwaters Region Based on the Google Earth Engine *Remote Sensing* 2021, *x*, Article No.

Jiyan Wang, Huaizhang Sun, Junnan Xiong, Dong He, Weiming Cheng, Chongchong Ye, Zhiwei Yong and Xianglin Huang




Figure S1. Topographic features of Three-River Headwaters region: (a) elevation; (b) slope; (c) aspect; and (d) topographic relief.




Figure S2. The monthly means for (a) soil temperature, (b) soil moisture, (c) relative humidity, (d) temperature, (e) precipitation, (f) shortwave radiation in the Yangtze (A), Yellow (B), and Lancang (C) river basins in different

periods. Horizontal lines in box plots denote the 95th, 75th, 50th, 25th, and 5th percentiles from top to bottom; the rectangles represent the average values.




Figure S3. Spatial pattern of (a, d, g, j, m, and p), standard deviation (b, e, h, k, n, and q) and temporal trend (c, f, i, l, o, and r) for the monthly mean temperature, precipitation, relative humidity, shortwave radiation, soil temperature, and soil moisture at the start of the growing season.




Figure S4. Spatial pattern of (a, d, g, j, m, and p), standard deviation (b, e, h, k, n, and q), and temporal trend (c, f, i, l, o, and r) for the monthly mean temperature, precipitation, relative humidity, shortwave radiation, soil temperature, and soil moisture at the end of the growing season.




Figure S5. Spatial pattern (a, d, g, j, m, and p), standard deviation (b, e, h, k, n, and q), and temporal trend (c, f, i, l, o, and r) for the monthly mean temperature, precipitation, relative humidity, shortwave radiation, soil temperature, and soil moisture in length of the growing season.