

Table S1. Sequences of primer sets used for qPCR experiments.

Gene	Primer sequence (5' → 3')	
	Forward Primer	Reverse Primer
<i>CXCL-10</i>	CCAAGTGCTGCCGTCATTTT	CTCAACACGTGGGCAGGATA
<i>CXCL-11</i>	GCTGCTCAAGGCTTCCTTATGT	ACTTTGTCGCAGCCGTTACT
<i>IL-12</i>	GTGGAATGGCGTCTCTGTCT	CGGGTCTGGTTTGATGATGT
<i>IL-23</i>	CCCGTATCCAGTGTGAAGATG	AGGGAGGTGTGAAGTTGCTC
<i>CCL-2</i>	ACAAGAGGATCACCAGCAGC	ATTCCTTCTTGGGGTCAGCAC
<i>CCL-22</i>	GCTGTGGCAATTCAGACCTC	TGACGGATGTAGTCCTGGCA
<i>CD-163</i>	ATGCTTCCATCCAGTGCCTC	CACAAACCAAGAGTGCCGTG
<i>Arg-1</i>	CAGCACTGAGGAAAGCTGGT	CAGACCGTGGGTCTTCACA
<i>Rpl13a</i>	CACTCTGGAGGAGAAACGGAAGG	GCAGGCATGAGGCAAACAGTC

Table S2. List of Antibodies using in Wester blot.

Antibody	Dilution	Source	Cat. No.
Mouse monoclonal anti-β-actin	1:2500	Sigma-Aldrich Corp	A2228
Rabbit polyclonal anti-STAT3	1:200	Santa Cruz Biotechnology	sc-482
Mouse monoclonal anti-pSTAT3 (Tyr ⁷⁰⁵)	1:200	Santa Cruz Biotechnology	sc-8059

Rabbit policlonal anti-NF-kB p65	1:200	Santa Cruz Biotechnology	sc-372
Rabbit policlonal anti-pNF-kB p65 (Ser ₂₇₆)	1:200	Santa Cruz Biotechnology	sc-101749
Rabbit policlonal anti-Arg-1	1:100	ThermoFisher Scientific Inc	pa5-32267
Goat policlonal anti-IL-10	1:100	Santa Cruz Biotechnology	sc-1783