

Modulation of Phase II Drug-Metabolizing Enzymes in proliferating and differentiated CaCo-2 Cells by Hop-Derived Prenylflavonoids

Kateřina Lněničková ¹, Michaela Šadibolová ², Petra Matoušková ², Barbora Szotáková ², Lenka Skálová ² and Iva Boušová ^{2,*}

¹ Faculty of Medicine and Dentistry, Palacký University, Hněvotinská 3, 775 15 Olomouc, Czech Republic; katerina.lnenickova@upol.cz (K.L.)

² Faculty of Pharmacy in Hradec Králové, Charles University, Heyrovského 1203, 500 05, Hradec Králové, Czech Republic; sadibolom@faf.cuni.cz (M.Š.); matousp7@faf.cuni.cz (P.M.); szotakova@faf.cuni.cz (B.S.); skaloval@faf.cuni.cz (L.S.)

* Correspondence: Iva.Bousova@faf.cuni.cz (I.B.); Tel.: +420-495-067-406




Figure S1. mRNA expression of phase II DMEs in proliferating (a) and differentiated (b) CaCo-2 cells after 24 h and 72 h. The normalized expression level was calculated using the $2^{-\Delta\Delta C_t}$ method with the geometric mean of GAPDH and B2M as a reference gene. Results are presented as the mean \pm SD of 3 independent experiments. Statistical analyses were performed using Student's t-test with $p < 0.05$, * different from control after 24 h.