

















































































































303. Nagahama, M.; Sakurai, J. Distribution of labeled *Clostridium perfringens* epsilon toxin in mice. *Toxicon* **1991**, *29*, 211–217.
304. Nagahama, M.; Sakurai, J. High-affinity binding of *Clostridium perfringens* epsilon-toxin to rat brain. *Infect. Immun.* **1992**, *60*, 1237–1240.
305. Zhu, C.; Ghabriel, M.N.; Blumbergs, P.C.; Reilly, P.L.; Manavis, J.; Youssef, J.; Hatami, S.; Finnie, J.W. *Clostridium perfringens* prototoxin-induced alteration of endothelial barrier antigen (EBA) immunoreactivity at the blood brain barrier (BBB). *Exp. Neurol.* **2001**, *169*, 72–82.
306. Buxton, D. The use of an immunoperoxidase technique to investigate by light and electron microscopy the sites of binding of *Clostridium welchii* type D e-toxin in mice. *J. Med. Microbiol.* **1978**, *11*, 289–292.
307. Finnie, J.W.; Blumbergs, P.C.; Manavis, J. Neuronal damage produced in rat brains by *Clostridium perfringens* type D epsilon-toxin. *J. Comp. Path.* **1999**, *120*, 415–420.
308. Miyamoto, O.; Minami, J.; Toyoshima, T.; Nakamura, T.; Masada, T.; Nagao, S.; Negi, T.; Itano, T.; Okabe, A. Neurotoxicity of *Clostridium perfringens* epsilon-toxin for the rat hippocampus via glutamanergic system. *Infect. Immun.* **1998**, *66*, 2501–2508.
309. Miyamoto, O.; Sumitami, K.; Nakamura, T.; Yamagani, S.; Miyatal, S.; Itano, T.; Negi, T.; Okabe, A. *Clostridium perfringens* epsilon toxin causes excessive release of glutamate in the mouse hippocampus. *FEMS Microbiol. Lett.* **2000**, *189*, 109–113.
310. Cole, A.R.; Gibert, M.; Popoff, M.R.; Moss, D.S.; Titball, R.W.; Basak, A. *Clostridium perfringens* ε-toxin shows structural similarity to the pore-forming toxin aerolysin. *Nat. Struct. Mol. Biol.* **2004**, *11*, 797–798.
311. Knapp, O.; Maier, E.; Benz, R.; Geny, B.; Popoff, M.R. Identification of the channel-forming domain of *Clostridium perfringens* Epsilon-toxin (ETX). *Biochim. Biophys. Acta* **2009**, *1788*, 2584–2593.
312. Cole, A. Structural studies on epsilon toxin from *Clostridium perfringens*. In *Protein Toxins of the Genus Clostridium and Vaccination*, Duchesnes, C., Mainil, J., Popoff, M.R., Titball, R. Eds.; Presses de la Faculté de Médecine Vétérinaire: Liège, Montreal, Canada, 2003; p. 95.
313. Miyata, S.; Minami, J.; Tamai, E.; Matsushita, O.; Shimamoto, S.; Okabe, A. *Clostridium perfringens* ε-toxin forms a heptameric pore within the detergent-insoluble microdomains of Madin-Darby Canine Kidney Cells and rat synaptosomes. *J. Biol. Chem.* **2002**, *277*, 39463–39468.
314. Petit, L.; Gibert, M.; Gillet, D.; Laurent-Winter, C.; Boquet, P.; Popoff, M.R. *Clostridium perfringens* epsilon-toxin acts on MDCK cells by forming a large membrane complex. *J. Bacteriol.* **1997**, *179*, 6480–6487.
315. Petit, L.; Maier, E.; Gibert, M.; Popoff, M.R.; Benz, R. *Clostridium perfringens* epsilon-toxin induces a rapid change in cell membrane permeability to ions and forms channels in artificial lipid bilayers. *J. Biol. Chem.* **2001**, *276*, 15736–15740.
316. Petit, P.; Breard, J.; Montalescol, V.; Ben El Hadj, N.; Levade, T.; Popoff, M.R.; Geny, B. Lethal toxin from *Clostridium sordellii* induces apoptotic cell death by disruption of mitochondrial homeostasis in HL-60 cells. *Cell. Microbiol.* **2003**, *5*, 761–771.
317. Miyata, S.; Matsushita, O.; Minami, J.; Katayama, S.; Shimamoto, S.; Okabe, A. Cleavage of C-terminal peptide is essential for heptamerization of *Clostridium perfringens* ε-toxin in the synaptosomal membrane. *J. Biol. Chem.* **2001**, *276*, 13778–13783.

318. Petit, L.; Gibert, M.; Gourch, A.; Bens, M.; Vandewalle, A.; Popoff, M.R. *Clostridium perfringens* Epsilon Toxin rapidly decreases membrane barrier permeability of polarized MDCK Cells. *Cell. Microbiol.* **2003**, *5*, 155–164.
319. Chassin, C.; Bens, M.; de Barry, J.; Courjaret, R.; Bossu, J.L.; Cluzeaud, F.; Ben Mkaddem, S.; Gibert, M.; Poulain, B.; Popoff, M.R.; Vandewalle, A. Pore-forming epsilon toxin causes membrane permeabilization and rapid ATP depletion-mediated cell death in renal collecting duct cells. *Am. J. Physiol. Renal. Physiol.* **2007**, *293*, F927–F937.
320. Farthing, M.J. Enterotoxins and the enteric nervous system--a fatal attraction. *Int. J. Med. Microbiol.* **2000**, *290*, 491–496.
321. Farthing, M.J.G.; Casburn-Jones, A.; Banks, M.R. Enterotoxins, enteric nerves, and intestinal secretion. *Curr. Gastroenterol. Rep.* **2004**, *6*, 177–180.
322. Hirst, T.R.; D'Souza, J.M. *Vibrio cholerae* and *Escherichia coli* thermolabile enterotoxin. In *The Sourcebook of Bacterial Protein Toxins*, 3rd ed.; Alouf, J.E., Popoff, M.R., Eds.; Elmsevier Academic Press: Amsterdam, The Netherland, 2006; pp. 270–290.
323. Holmes, R.K.; Jobling, M.G.; Conell, T.D. Cholera toxin and related enterotoxins of Gram-negative bacteria. In *Bacterial Toxins and Virulence Factors in Disease*; Moss, J., Iglewski, B., Vaughan, M., Tu, A.T., Eds.; Marcel Dekker: New York, NY, USA, 1995; pp. 225–255.
324. de Haan, L.; Hirst, T.R. Cholera toxin: A paradigm for multi-functional engagement of cellular mechanisms. *Mol. Membr. Biol.* **2004**, *21*, 77–92.
325. Nichols, B.J. A distinct class of endosome mediates clathrin-independent endocytosis to the Golgi complex. *Nat. Cell Biol.* **2002**, *4*, 374–378.
326. Fujinaga, Y.; Wolf, A.A.; Rodighiero, C.; Wheeler, H.E.; Tsai, B.; Allen, L.; Jobling, M.G.; Rapoport, T.; Holmes, R.K.; Lencer, W.I. Gangliosides that associate with lipid rafts mediate transport of cholera and related toxins from the plasma membrane to endoplasmic reticulum. *Mol. Biol. Cell* **2003**, *14*, 4783–4793.
327. Johannes, L.; Tenza, D.; Antony, C.; Goud, B. Retrograde transport of KDEL-bearing B-fragment of Shiga toxin. *J. Biol. Chem.* **1997**, *272*, 19554–19561.
328. Lundgren, O. 5-Hydroxytryptamine, enterotoxins, and intestinal fluid secretion. *Gastroenterology* **1998**, *115*, 1009–1012.
329. Turvill, J.L.; Mourad, F.H.; Farthing, M.J. Crucial role for 5-HT in cholera toxin but not *Escherichia coli* heat-labile enterotoxin-intestinal secretion in rats. *Gastroenterology* **1998**, *115*, 883–890.
330. Mourad, F.H.; O'Donnell, L.J.; Dias, J.A.; Ogutu, E.; Andre, E.A.; Turvill, J.L.; Farthing, M.J. Role of 5-hydroxytryptamine type 3 receptors in rat intestinal fluid and electrolyte secretion induced by cholera and *Escherichia coli* enterotoxins. *Gut* **1995**, *37*, 340–345.
331. Bearcroft, C.P.; Perrett, D.; Farthing, M.J. 5-hydroxytryptamine release into human jejunum by cholera toxin. *Gut* **1996**, *39*, 528–531.
332. Nilsson, O.; Cassuto, J.; Larsson, P.A.; Jodal, M.; Lidberg, P.; Ahlman, H.; Dahlstrom, A.; Lundgren, O. 5-Hydroxytryptamine and cholera secretion: A histochemical and physiological study in cats. *Gut* **1983**, *24*, 542–548.

333. Beubler, E.; Horina, G. 5-HT<sub>2</sub> and 5-HT<sub>3</sub> receptor subtypes mediate cholera toxin-induced intestinal fluid secretion in the rat. *Gastroenterology* **1990**, *99*, 83–89.
334. Buchheit, K.H. Inhibition of cholera toxin-induced intestinal secretion by the 5-HT<sub>3</sub> receptor antagonist ICS 205–930. *Naunyn Schmiedebergs Arch. Pharmacol.* **1989**, *339*, 704–705.
335. Beattie, D.T.; Smith, J.A. Serotonin pharmacology in the gastrointestinal tract: A review. *Naunyn Schmiedebergs Arch. Pharmacol.* **2008**, *377*, 181–203.
336. Cooke, H.J. Neurotransmitters in neuronal reflexes regulating intestinal secretion. *Ann. NY Acad. Sci.* **2000**, *915*, 77–80.
337. Lundgren, O. Enteric nerves and diarrhoea. *Pharmacol. Toxicol.* **2002**, *90*, 109–120.
338. Turvill, J.L.; Connor, P.; Farthing, M.J. Neurokinin 1 and 2 receptors mediate cholera toxin secretion in rat jejunum. *Gastroenterology* **2000**, *119*, 1037–1044.
339. Castagliuolo, I.; Lamont, J.T.; Letourneau, R.; Kelly, C.; O'Keane, J.C.; Jaffer, A.; Theoharides, T.C.; Pothoulakis, C. Neuronal involvement in the intestinal effects of *Clostridium difficile* Toxin A and *Vibrio cholerae* enterotoxin in rat ileum. *Gastroenterology* **1994**, *107*, 657–665.
340. Pothoulakis, C.; Lamont, J.T. Microbes and microbial toxins: paradigms for microbial-mucosa interactions II. The integrated response of the intestine to *Clostridium difficile* toxins. *Am. J. Physiol. Gastrointest. Liver Physiol.* **2001**, *280*, G178–G183.
341. Castagliuolo, I.; Riegler, M.; Pasha, A.; Nikulasson, S.; Lu, B.; Gerard, C.; Gerard, N.P.; Pothoulakis, C. Neurokinin-1 (NK-1) receptor is required in *Clostridium difficile* -induced enteritis. *J. Clin. Invest.* **1998**, *101*, 1547–1550.
342. Castagliuolo, I.; Wang, C.C.; Valenick, L.; Pasha, A.; Nikulasson, S.; Carraway, R.E.; Pothoulakis, C. Neurotensin is a proinflammatory peptide in colonic inflammation. *J. Clin. Invest.* **1999**, *103*, 843–849.
343. Xia, Y.; Hu, H.Z.; Liu, S.; Pothoulakis, C.; Wood, J.D. *Clostridium difficile* toxin A excites enteric neurones and suppresses sympathetic neurotransmission in the guinea pig. *Gut* **2000**, *46*, 481–486.
344. McClane, B.A. *Clostridium perfringens* enterotoxin. In *The Comprehensive Sourcebook of Bacterial Protein Toxins*, 3rd ed.; Alouf, J.E., Popoff, M.R., Eds.; Elsevier Academic Press: Amsterdam, The Netherlands, 2006; pp. 763–778.
345. Senda, T.; Sugimoto, N.; Horiguchi, Y.; Matsuda, M. The enterotoxin of *Clostridium perfringens* type A binds to the presynaptic nerve endings in neuromuscular junctions of mouse phrenic nerve-diaphragm. *Toxicon* **1995**, *33*, 499–506.
346. Rolfe, V.E.; Levin, R.J. Vagotomy inhibits the jejunal fluid secretion activated by luminal ileal *Escherichia coli* STa in the rat *in vivo*. *Gut* **1999**, *44*, 615–619.
347. Mourad, F.H.; Nassar, C.F. Effect of vasoactive intestinal polypeptide (VIP) antagonism on rat jejunal fluid and electrolyte secretion induced by cholera and *Escherichia coli* enterotoxins. *Gut* **2000**, *47*, 382–386.
348. Rolfe, V.; Levin, R.J. Enterotoxin *Escherichia coli* STa activates a nitric oxide-dependent myenteric plexus secretory reflex in the rat ileum. *J. Physiol.* **1994**, *475*, 531–537.
349. Eklund, S.; Jodal, M.; Lundgren, O. The enteric nervous system participates in the secretory response to the heat stable enterotoxins of *Escherichia coli* in rats and cats. *Neuroscience* **1985**, *14*, 673–681.

350. Stenfors Arnesen, L.P.; Fagerlund, A.; Granum, P.E. From soil to gut: *Bacillus cereus* and its food poisoning toxins. *FEMS Microbiol. Rev.* **2008**, *32*, 579–606.
351. Toh, M.; Moffitt, M.C.; Henrichsen, L.; Raftery, M.; Barrow, K.; Cox, J.M.; Marquis, C.P.; Neilan, B.A. Cereulide, the emetic toxin of *Bacillus cereus*, is putatively a product of nonribosomal peptide synthesis. *J. Appl. Microbiol.* **2004**, *97*, 992–1000.
352. Agata, N.; Ohta, M.; Mori, M.; Isobe, M. A novel dodecadepsipeptide, cereulide, is an emetic toxin of *Bacillus cereus*. *FEMS Microbiol. Lett.* **1995**, *129*, 17–20.
353. Horwood, P.F.; Burgess, G.W.; Oakey, H.J. Evidence for non-ribosomal peptide synthetase production of cereulide (the emetic toxin) in *Bacillus cereus*. *FEMS Microbiol. Lett.* **2004**, *236*, 319–324.
354. Agata, N.; Ohta, M.; Mori, M. Production of an emetic toxin, cereulide, is associated with a specific class of *Bacillus cereus*. *Curr. Microbiol.* **1996**, *33*, 67–69.
355. Shinagawa, K.; Konuma, H.; Sekita, H.; Sugii, S. Emesis of rhesus monkeys induced by intragastric administration with the HEp-2 vacuolation factor (cereulide) produced by *Bacillus cereus*. *FEMS Microbiol. Lett.* **1995**, *130*, 87–90.
356. Agata, N.; Mori, M.; Ohta, M.; Suwan, S.; Ohtani, I.; Isobe, M. A novel dodecadepsipeptide, cereulide, isolated from *Bacillus cereus* causes vacuole formation in HEp-2 cells. *FEMS Microbiol. Lett.* **1994**, *121*, 31–34.
357. Yokoyama, K.; Ito, M.; Agata, N.; Isobe, M.; Shibayama, K.; Horii, T.; Ohta, M. Pathological effect of synthetic cereulide, an emetic toxin of *Bacillus cereus*, is reversible in mice. *FEMS Immunol. Med. Microbiol.* **1999**, *24*, 115–120.
358. Virtanen, S.M.; Roivainen, M.; Andersson, M.A.; Ylipaasto, P.; Hoornstra, D.; Mikkola, R.; Salkinoja-Salonen, M.S. *In vitro* toxicity of cereulide on porcine pancreatic Langerhans islets. *Toxicon* **2008**, *51*, 1029–1037.
359. Andersson, M.A.; Hakulinen, P.; Honkalampi-Hamalainen, U.; Hoornstra, D.; Lhuguenot, J.C.; Maki-Paakkanen, J.; Savolainen, M.; Severin, I.; Stamatii, A.L.; Turco, L.; Weber, A.; von Wright, A.; Zucco, F.; Salkinoja-Salonen, M. Toxicological profile of cereulide, the *Bacillus cereus* emetic toxin, in functional assays with human, animal and bacterial cells. *Toxicon* **2007**, *49*, 351–367.
360. Mikkola, R.; Saris, N.E.; Grigoriev, P.A.; Andersson, M.A.; Salkinoja-Salonen, M.S. Ionophoretic properties and mitochondrial effects of cereulide: the emetic toxin of *B. cereus*. *Eur. J. Biochem.* **1999**, *263*, 112–117.
361. Teplova, V.V.; Mikkola, R.; Tonshin, A.A.; Saris, N.E.; Salkinoja-Salonen, M.S. The higher toxicity of cereulide relative to valinomycin is due to its higher affinity for potassium at physiological plasma concentration. *Toxicol. Appl. Pharmacol.* **2006**, *210*, 39–46.
362. Saris, N.E.; Andersson, M.A.; Mikkola, R.; Andersson, L.C.; Teplova, V.V.; Grigoriev, P.A.; Salkinoja-Salonen, M.S. Microbial toxin's effect on mitochondrial survival by increasing K<sup>+</sup> uptake. *Toxicol. Ind. Health.* **2009**, *25*, 441–446.
363. Krakauer, T.; Stiles, B.G. Staphylococcal enterotoxins, toxic-shock syndrome toxin-1, and streptococcal pyrogenic exotoxins: Some basic biology of bacterial superantigens. *Rec. Res. Dev. Infect. Immun.* **2003**, *1*, 21–27.

364. Jett, M.; Brinkley, W.; Neill, R.; Gemski, P.; Hunt, R. Staphylococcus aureus enterotoxin B challenge of monkeys: Correlation of plasma levels of arachidonic acid cascade products with occurrence of illness. *Infect. Immun.* **1990**, *58*, 3494–3499.
365. Alber, G.; Scheuber, P.H.; Reck, B.; Sailer-Kramer, B.; Hartmann, A.; Hammer, D.K. Role of substance P in immediate-type skin reactions induced by staphylococcal enterotoxin B in unsensitized monkeys. *J. Allergy Clin. Immunol.* **1989**, *84*, 880–885.
366. Tiegs, G.; Bang, R.; Neuhuber, W.L. Requirement of peptidergic sensory innervation for disease activity in murine models of immune hepatitis and protection by beta-adrenergic stimulation. *J. Neuroimmunol.* **1999**, *96*, 131–143.
367. Wang, X.; Wang, B.R.; Zhang, X.J.; Duan, X.L.; Guo, X.; Ju, G. Fos expression in the rat brain after intraperitoneal injection of Staphylococcus enterotoxin B and the effect of vagotomy. *Neurochem. Res.* **2004**, *29*, 1667–1674.
368. Hu, D.L.; Zhu, G.; Mori, F.; Omoe, K.; Okada, M.; Wakabayashi, K.; Kaneko, S.; Shinagawa, K.; Nakane, A. Staphylococcal enterotoxin induces emesis through increasing serotonin release in intestine and it is downregulated by cannabinoid receptor 1. *Cell Microbiol.* **2007**, *9*, 2267–2277.

© 2010 by the authors; licensee Molecular Diversity Preservation International, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).