

Supplementary Figures

Profiling Immune Escape in Hodgkin's and Diffuse large B-Cell Lymphomas Using the Transcriptome and Immunostaining

Sarah Pericart, Marie Tosolini, Pauline Gravelle, Cédric Rossi, Alexandra Traverse-Glehen, Nadia Amara, Camille Franchet, Elodie Martin, Christine Bezombes, Guy Laurent, Pierre Brousset, Jean-Jacques Fournié and Camille Laurent




Figure S1. Overall survival (OS) curves according to the four stages of IE of DLBCL patients from public microarrays data (n = 200) (log rank $p = 0.0006$).




Figure S2. Concordance between Hans algorithm [51] based on immunohistochemical finding and molecular classification based on GEP [52] in lymphoma cohort of our institution.

Table S5. Intra-class correlation coefficient between manual scoring and automated scoring for IHC: CD3, CD8, CD4, PD-1, PD-L1, TIM-3 and LAG-3.

IHC	ICC	[95% CI]
CD3	0.75	[0.52 ; 0.87]
CD8	0.81	[0.61 ; 0.90]
CD4	0.86	[0.75 ; 0.92]
PD-1	0.70	[0.49 ; 0.84]
PD-L1	0.87	[0.68 ; 0.94]
TIM3	0.72	[0.52 ; 0.85]
LAG3	0.89	[0.74 ; 0.95]