

Supplementary Material

Prognostic Value of Simple Non-Invasive Tests for the Risk Stratification of Incident Hepatocellular Carcinoma in Cirrhotic Individuals with Non-Alcoholic Fatty Liver Disease

Angelo Armandi ^{1,2,†}, Gian Paolo Caviglia ^{1,†}, Amina Abdulle ¹, Chiara Rosso ¹, Kamela Gjini ¹, Gabriele Castelnuevo ¹, Marta Guariglia ¹, Nuria Perez Diaz del Campo ¹, Daphne D'Amato ¹, Davide Giuseppe Ribaldone ¹, Giorgio Maria Saracco ¹ and Elisabetta Bugianesi ^{1,*}

¹ Department of Medical Sciences, University of Turin, 10126 Turin, Italy

² Metabolic Liver Disease Research Program, University Medical Center, Department of Internal Medicine I, Johannes Gutenberg University, 55131 Mainz, Germany

* Correspondence: elisabetta.bugianesi@unito.it

† These authors equally contributed to the manuscript.

Table S1. Cumulative probability and hazard ratios of HCC by AAR, APRI, and BARD risk category.

NITs	HCC (n)	Cumulative (%)	<i>p</i> value ¹	HR (95% CI)	<i>p</i> value ²
AAR T1	2/41	4.9%	0.204	2.03 (0.87–4.71)	0.204
AAR T2	6/40	15.0%			
AAR T3	5/41	12.1%			
APRI <0.5	1/32	3.1%	0.062	3.41 (0.89–13.03)	0.073
APRI 0.5–1.5	10/79	12.7%			
APRI >1.5	2/11	18.2%			
BARD 0–1	1/20	5.0%	0.820	1.19 (0.49–2.86)	0.703
BARD 2–3	8/66	12.1%			
BARD 4	4/36	11.1%			

¹ *p* value was calculated by log-rank test. ² *p* value was calculated by Cox regression analysis. Abbreviations: AAR, AST to ALT ratio; APRI, AST to platelets ratio index; BARD, body mass index, AST/ALT ratio, diabetes mellitus score; CI, confidence interval; HCC, hepatocellular carcinoma; HR, hazard ratio; *n*, number.

