

Review

BRCA1 and Oxidative Stress

Yong Weon Yi ¹, Hyo Jin Kang ¹ and Insoo Bae ^{1,2,*}

¹ Department of Oncology, Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, Washington, DC 20057, USA; E-Mails: yy226@georgetown.edu (Y.W.Y.); hk73@georgetown.edu (H.J.K.)

² Department of Radiation Medicine, Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, Washington, DC 20057, USA

* Author to whom correspondence should be addressed; E-Mail: ib42@georgetown.edu; Tel.: +1-202-687-5267.

Received: 19 December 2013; in revised form: 20 March 2014 / Accepted: 24 March 2014 /

Published: 3 April 2014

Abstract: The breast cancer susceptibility gene 1 (BRCA1) has been well established as a tumor suppressor and functions primarily by maintaining genome integrity. Genome stability is compromised when cells are exposed to oxidative stress. Increasing evidence suggests that BRCA1 regulates oxidative stress and this may be another mechanism in preventing carcinogenesis in normal cells. Oxidative stress caused by reactive oxygen species (ROS) is implicated in carcinogenesis and is used strategically to treat human cancer. Thus, it is essential to understand the function of BRCA1 in oxidative stress regulation. In this review, we briefly summarize BRCA1's many binding partners and mechanisms, and discuss data supporting the function of BRCA1 in oxidative stress regulation. Finally, we consider its significance in prevention and/or treatment of BRCA1-related cancers.


Keywords: BRCA; oxidative stress; reactive oxygen species (ROS); carcinogenesis; detoxification

1. Introduction

Mutations of BRCA1 are found in a high percentage of hereditary breast and ovarian cancers [1]. In the early 1990s, chromosome 17q21 was assigned to contain the locale of a gene responsible for inherited susceptibility to breast cancer in families with early-onset disease [2,3]. Subsequently, the

BRCA1 gene was cloned and analyzed to be composed of 22 coding exons distributed over ~100 kb of genomic DNA [4,5]. The full-length *BRCA1* cDNA encodes a large protein of 1863 amino acids (Figure 1) [4]. The *BRCA1* protein contains an *N*-terminal really interesting new gene (RING) finger domain and two *BRCA1* C-Terminal (BRCT) domains, which are involved in various protein-protein interactions [6–12]. The *BRCA1* protein also has a DNA-binding domain [13,14] and an SQ-cluster domain (SCD) in its central region [15]. It also has two nuclear localization signals (NLSs) [16,17] and a nuclear export signal (NES) [18]. Localization of *BRCA1* is regulated by interactions with other proteins such as the *BRCA1*-associated RING domain protein 1 (BARD1) [19,20]. The *BRCA1* gene is conserved in mammals [21,22] and the *BRCA1* protein has roles in various cellular events including cell cycle control, DNA damage signaling, maintaining genomic integrity, protein ubiquitination, and transcriptional regulation. All these responsibilities of *BRCA1* may collectively participate in tumor suppression; however, precise mechanisms of how the *BRCA1* protein functions as a tumor suppressor are still not yet fully understood [23,24].

Figure 1. The *BRCA1* protein and its binding transcription factors.



Every cell is inevitably exposed to oxidative stress, either extracellular or intracellular, during its life time [25,26]. Reactive oxygen species (ROS) are produced in a controlled manner in normal cells and act as important signaling molecules to regulate various cellular functions including transcriptional regulation and signal transduction (reviewed in [25–29]). However, uncontrolled production of ROS causes oxidative stress resulting in DNA damage [30–35] and impaired cellular functions leading to various human diseases including cancer [26,36–38]. It is interesting that recent data demonstrate that *BRCA1* is involved in oxidative stress regulation. These findings prompt the thought that *BRCA1* might exert its tumor suppressive functions through oxidative stress regulation. Given that the mechanisms of *BRCA1* tumor suppression have not yet been fully elucidated, we summarize and explore the potential role of *BRCA1* in response to ROS and carcinogenesis in the context of its known functions.

2. BRCA1 as a Tumor Suppressor

Women carrying germline mutations in the *BRCA1* gene have a 50%–80% lifetime risk of developing breast cancer and a 20%–40% lifetime risk of developing ovarian cancer [39]. Decrease or absence of the BRCA1 protein in sporadic breast and ovarian cancers suggests that BRCA1 functions as a tumor suppressor in nonhereditary tumors as well [40,41]. In addition, it has been reported that germline mutations in the *BRCA1* gene confer the risk of prostate [42–44] and pancreatic cancer [45–53].

Tumor suppressor functions of BRCA1 have been studied in mouse models. Deficiency of *Brcal*, in which *Brcal* mutations encode truncated proteins that lack the C-terminal half of the protein, results in early embryonic lethality ([54,55] and references therein). Mouse models carrying a series of allelic mutations in *Brcal* have been created to overcome this embryonic lethality [54]. Cells carrying these *BRCA1* mutations show genomic instability and defects in DNA damage repair [54]. However, the loss of *Brcal* alone is not sufficient for tumorigenesis. Rather, multiple genetic alterations, including the inactivation of p53 and activation of a number of oncogenes caused by genomic instability in *Brcal*-defective cells may cause mammary tumorigenesis [54]. Most recently, two separate groups demonstrated the importance of two domains, BRCT and RING, in the tumor suppressor function of *Brcal* in mouse models [56,57]. A RING mutation (C61G), found in human cancer, destabilizes the BRCA1/BARD1 heterodimer and abrogates ubiquitin ligase activity [58–61]. Mice homozygous for this mutation, in a mutated p53 background, were found to rapidly develop tumors than mice carrying a *Brcal*-null allele [56]. A mutation of S1598F in BRCT domain that ablates phosphoprotein recognition was also found to be defective in the control of genome integrity and this mutant *Brcal* failed to function as a tumor suppressor in three separate mouse models of *Brcal*-dependent tumorigenesis [57]. Thus, genome stability is a key for healthy cells and the main function of BRCA1 is to keep the genome stable. Since ROS cause DNA damage, then it is plausible that BRCA1 normally functions to regulate oxidative stress.

2.1. Genome Stability, Oxidative Stress and Tumor Suppression

Genomic instability observed in BRCA1-defective cells, either by loss of function or by mutations of BRCA1, results from defects in DNA damage repair [62,63]. This genomic instability may facilitate permanent oncogenic changes in the genome [24]. Various carcinogenic insults such as ultraviolet (UV) light, ionizing radiation and environmental carcinogens generate ROS in cells. Excess ROS generate oxidative stress that sustains uncontrolled cell proliferation that eventually causes genomic instability and cancer ([30–35] and references therein). In addition, accumulating data suggest that residual oxidative stresses from xenobiotics promote tumorigenesis [64].

Despite evidence that maintenance of genome integrity by BRCA1 serves as an important tumor suppressor activity, the exact biochemical activity that responsible for tumor suppression by BRCA1 still remains elusive [23,24].

2.2. Mouse Model: Genome Stability, Oxidative Stress and Tumor Suppression

A study in *Brcal*^{Δ11/Δ11} *p53*^{+/-} mice also suggests that BRCA1 functions in the oxidative stress response. These mice display spontaneous tumor formation in mammary gland and ovary as well as

lymphoma [65–68]. It has been reported that these mice exhibit increased expression of Redd1, elevated ROS and are more sensitive to oxidative stress induced lethality [69]. In this study, *Brca1^{Δ11/Δ11}p53^{+/-}* mice were treated with methyl-*N*-amyl nitrosamine (MNAN), a carcinogen that specifically induces esophageal tumors in mice and rats [69]. In this background MNAN-treated mice had a significantly increased esophagus and forestomach tumor incidence as compared with that of control mice. Although the significance of these MNAN-induced tumors is not yet understood in human BRCA1-related cancers, these results suggest that *Brca1* is involved in the regulation of ROS in carcinogenesis of the mouse.

2.3. Cell Studies: Genome Stability, Oxidative Stress and Tumor Suppression

In cell studies, BRCA1 was reported to associate with complex DNA damage such as double-strand breaks (DSBs) and non-DSB bistranded oxidatively induced clustered DNA lesions (OCDLs) [70,71]. In fact, a large protein complex, named BRCA1-associated genome surveillance complex (BASC), was known to be involved in the recognition of abnormal DNA structures or damaged DNA to repair them [72]. HCC1937 breast cancer cells, that harbor a 5382insC germ-line mutation of BRCA1, were demonstrated to have a significant accumulation of various types of DNA damage and chromatid breaks after ionizing radiation compared to MCF7 cells that are BRCA1 hemizygous [71]. And partial restoration of wild type BRCA1 in HCC1937 cells increased the overall repair of OCDLs [71].

BRCA1 was also reported to regulate the transcription-coupled repair of the 8-oxoguanine (8-oxoG) DNA lesion in HCC1937 cells [73]. The 8-oxoG is a well-known DNA lesion that can be caused by oxidative stress [74]. Subsequent studies have repeatedly reported that the loss of BRCA1 function results in the increase of 8-oxoG and overexpression of wild type BRCA1 increases the repair of 8-oxoG DNA lesion [75–78]. The increase of 8-oxoG DNA lesion is linked to transcriptional regulation of the base excision repair (BER) pathway by BRCA1 [76].

Besides the function of BRCA1 in BER to remove the 8-oxoG DNA lesion, varied reports suggest that BRCA1 regulates oxidative stress itself: loss of BRCA1 increases the cellular ROS and overexpression of BRCA1 suppresses ROS production [77–82]. Overexpression of wild type BRCA1 was reported to suppress both basal and the H₂O₂-mediated ROS production determined by carboxy-DCF fluorescence in MCF7 cells. Consistent with this, the siRNA-based BRCA1 knockdown (BRCA1-KD) enhanced both basal and the H₂O₂-mediated ROS production in one normal-like (MCF10A) and two breast cancer (MCF7 and T47D) cell lines [77]. Under these conditions, wild type BRCA1 reduced protein nitration and the H₂O₂-induced 8-oxoG formation in a prostate cancer cell line DU145. Again the 8-oxoG formation was increased in BRCA1-KD DU145 cells [77].

BRCA1-KD also enhanced the ROS accumulation in a normal-like breast epithelial cell, MCF10A in the absence or presence of benzo[a]pyrene (B[a]P) [78,79]. Under these conditions, BRCA1-KD also increased 8-oxoG formation in MCF10A cells [78]. ROS induced by multiple environmental factors such as paraquat, 2,3,7,8-tetrachlorodibenzodioxin (TCDD), 7,12-dimethylbenz[a]anthracene (DMBA), 2-hydroxyestradiol (2OHE2) and 4-hydroxyestradiol (4OHE2) were also increased in BRCA1-KD MCF10A cells in the presence of B[a]P [79]. It had also been reported that overexpression of wild type BRCA1 in HCC1937 cells reduces the generation of H₂O₂ during co-culture with fibroblasts [80]. The inhibition of H₂O₂ generation was also confirmed in an isogenic

ovarian cancer cell model UWB1.289, which lacks a wild type *BRCA1* gene and UWB1.289+BRCA1, which expresses wild type BRCA1 by stable transfection [81]. More recently, it has been demonstrated that wild type BRCA1 suppresses the H₂O₂-induced ROS production in human aortic smooth muscle cells [82]. Thus, there is substantial evidence of oxidative stress regulation by BRCA1 in the destabilized genome.

2.4. *BRCA1* and *p53*: Genome Stability, Oxidative Stress and Tumor Suppression

The tumor suppressor protein p53 also contributes to the regulation of oxidative stress in response to ROS (reviewed in [83]). BRCA1 (aa 224–550) has been found to interact with the C-terminus of p53 and activate its transcriptional activity at the promoter of p21 or BAX [84–86]. Subsequently, the C-terminus of BRCA1 (aa 1760–1863) has been also reported to bind p53 and is sufficient to activate the p53-dependent transcription from the p21 promoter [87]. Interestingly, BRCA1 stabilizes p53 and induces the p53-dependent transcription of genes that are involved in DNA repair and growth arrest rather than proapoptotic genes, while DNA-damaging agents-stabilized p53 leads to the transcription of multiple genes including apoptotic genes [88]. BRCA1 may also promote the accumulation of p53 in an ARF (alternate reading frame; p14^{ARF})-dependent manner: BRCA1 does not stabilize p53 in p14^{ARF}-deficient A375 cells [89]. In addition, the BRCA1/BARD1 complex can stabilize p53 through phosphorylation of p53 at Ser15 residue by the ataxia telangiectasia mutated (ATM) and the ATM and Rad3-related (ATR) proteins in response to gamma- and UV-irradiation, respectively [90]. Although these results suggest that BRCA1 potentially contributes to the oxidative stress response through the regulation of p53, further experiments are needed to determine the exact role of BRCA1/p53 in this regulation. Interestingly, a recent report demonstrated that ATM kinase negatively regulates ARF by ubiquitin-dependent degradation: ATM-activated protein phosphatase 1 (PP1) antagonized Nek2-dependent phosphorylation of nucleophosmin (NPM) to liberate ARF from NPM and render it susceptible to degradation by the UFL1 E3-ubiquitin ligase [91]. Since stabilization of p53 by BRCA1 is dependent on both ARF and ATM, studies on the potential role of BRCA1 in ATM/ARF pathway may provide an additional insight on the regulation of p53 stability.

3. BRCA1 as an E3 Ubiquitin Ligase

One notable feature of BRCA1 is its E3 ligase activity that is mediated by heterodimerization with BARD1 through its N-terminal RING finger domain [58–61]. The ubiquitin-proteasome system is responsible for much of the regulated proteolysis in cells, and also has non-degradative regulatory functions [92]. Ubiquitin is a highly conserved 76 amino-acid protein that can be reversibly attached to other proteins by three critical enzymes: an ubiquitin-activating enzyme (E1), an ubiquitin-conjugating enzyme (E2), and an ubiquitin ligase (E3). The E3 ligase catalyzes the formation of poly- or mono-ubiquitination usually at a lysine side-chain of proteins [92]. The polyubiquitins are commonly linked through Lys48 of ubiquitin and serve as signals for proteasome-mediated proteolysis [93]. The BRCA1-BARD1 ligase complex, however, catalyzes unconventional polyubiquitins including Lys6-linked chains [94–96]. Although BRCA1 catalyzes polyubiquitination of numerous proteins such as the nucleophosmin/B23 (NPM1) [97], the CtBP-interacting protein (CtIP) [98], the polymerase (RNA) II (DNA directed) polypeptide H (POLR2H, also known as RPB8) [99], the transcription factor

IIE (TFIIE) [100] and BRCA1 itself [95,101], there is no direct evidence that these polyubiquitinations signal degradation of substrate proteins [102,103]. A recent report demonstrated that BRCA1 polyubiquitinates G2/M cell cycle proteins including cyclin B and Cdc25C and targets these proteins to be degraded by the ubiquitin-proteasome pathway in an APC/C-independent manner in breast cancer cells [104]. BRCA1-dependent degradation of these proteins was enhanced following DNA damage, which suggests that BRCA1 targets cell cycle proteins for degradation in response to DNA damage. This may represent a possible mechanism to prevent unscheduled mitotic entry by inhibiting accumulation of these proteins [104].

BRCA1 also catalyzes the monoubiquitination of several substrates including core histones H2A, H2B, H3, H4, and the DNA-damage-responsive histone variants H2AX [60,101], as well as γ -tubulin [105] and the estrogen receptor α (ER α) [106]. Recently, it has been demonstrated that BRCA1 ubiquitinates H2A at satellite repeat DNA regions *in vivo* contributing to silencing of heterochromatin [107]. Thus, BRCA1's E3 ubiquitin ligase activity is an important clue as to how it functions.

3.1. BRCA1 E3 Ligase Activity and Tumor Suppression

Homologous recombination (HR) repair is a vital process employed during stalled DNA replication or repair of DSBs caused by many DNA-damaging agents [108]. Because the BRCA1/BARD1 complex is absolutely required for HR repair, the E3 ligase activity of this complex may play a pivotal role in HR repair process (reviewed in [108–110]). As mentioned earlier, a RING mutation (C61G) of Brca1, which lacks ubiquitin ligase activity, failed to protect from tumor formation in mice with a mutated p53 background [56]. On the contrary, another RING mutation (I26A) that also lacks E3 ligase activity could prevent tumor formation in alternative models [57]. The exact role of the E3 ligase activity of BRCA1 in tumor suppression still remains the subject of debate [24].

Currently the role of the BRCA1/BARD1 ligase complex in oxidative stress is not defined yet. However, given the regulation of the nuclear factor erythroid 2-related factor 2 (NRF2) stability through physical interaction with BRCA1 (see below), it is plausible that this ligase activity contributes the modulation of proteins in NRF2 pathway.

3.2. BRCA1: Oxidative Stress and Ubiquitination

Interestingly, BRCA1 induces p21 expression [84–87,111], which itself exerts the antioxidant effects in response to oxidative stress through direct interaction with NRF2 [112]. NRF2 serves as a master transcription factor that activates transcription of antioxidant enzymes [113–116]. The level of NRF2 is tightly regulated by ubiquitination, which is mediated by the Kelch-like erythroid cell-derived protein with CNC homology-associated protein 1 (KEAP1) [113–116]. By binding NRF2, p21 competes with KEAP1 for NRF2 binding to inhibit KEAP1-mediated NRF2 ubiquitination and degradation [112]. Thus, p21 and indirectly BRCA1 regulate ubiquitination and degradation of NRF2 in response to oxidative stress. Indeed, the NRF2-dependent antioxidant response of p21 was further confirmed in mouse embryonic fibroblast (MEF) cells: overexpression of p21 protected Nrf2^{+/+} MEF cells but not Nrf2^{-/-} MEF cells from the H₂O₂-induced cytotoxicity [112].

More recently, it has been reported that BRCA1 can be physically associated with NRF2 to prevent ubiquitin-dependent degradation by KEAP1 [117]. Treatment with L-buthionine-sulfoximine (BSO)

resulted in ROS accumulation and induced the interaction of overexpressed BRCA1 and NRF2 in 293FT cells [117]. Endogenous Brca1-Nrf2 interaction was also confirmed in a mouse epithelial cell line, COMMA-1D treated with BSO [117]. Mutational analysis of NRF2 revealed that an ETGE motif [118] of NRF2 is necessary to bind BRCA1 [117]. Moreover, reconstitution of wild type BRCA1 in HCC1937 cells restored NRF2 activation [117]. Since Nrf2-deficient mice exhibited a relative increase in the formation of tumor foci in the urethane-induced lung carcinogenesis model [119], loss and/or reduction of NRF2 activity by BRCA1 deficiency may contribute to the carcinogenesis of BRCA1-associated tumors. Hence, studies *in vitro* and *in vivo* demonstrate BRCA1 ability to regulate ubiquitination and degradation of proteins in response to oxidative stress.

4. BRCA1 as a Transcriptional Regulator

Although the BRCA1 protein has a DNA-binding domain [13,14], it is not a sequence-specific DNA-binding protein. Rather, it functions as a transcriptional modulator via physical interaction with a series of transcription factors (Figure 1) and regulates their target gene expression [120]. The potential role of BRCA1 in transcriptional regulation was first demonstrated by a fusion construct of BRCA1 C-terminus to the GAL4 DNA-binding domain (GAL4-BRCA1) [121–123]. The GAL4-BRCA1, containing a highly negatively charged C-terminus of BRCA1, could activate transcription in both yeast and mammalian cells [122]. Subsequently, the presence of certain cofactors was found to determine optimal transcriptional activation by GAL4-BRCA1 fusions [124]. In addition, this acidic C-terminal domain of BRCA1 was reported to induce chromatin remodeling and activation of chromosomal DNA replication when tethered to a cellular replication origin [125].

In addition, BRCA1 was found to be bound and co-purified with the RNA polymerase II (RNA Pol II) holoenzyme complex through an interaction with RNA helicase A (RHA) [126–128]. These findings suggest that BRCA1 is a component of the core transcriptional machinery. Interestingly, cancer-associated point mutations of BRCA1 disrupt the BRCA1-RNA Pol II interaction [126]. RHA and BRCA1 are known to interact with the coactivators p300 and the CREB-binding protein (CBP) [128,129]. BRCA1 is also known to modulate the phosphorylation of the C-terminal domain of RNA Pol II by direct interacting and inhibiting the Cdk-activating kinase (CAK) [130].

4.1. Binding Partners in BRCA1 Transcriptional Regulation

As mentioned, a study using GAL4-BRCA1 substantiated its role in chromatin remodeling [125]. Later, a yeast two-hybrid assay revealed the interaction of BRCA1 with the retinoblastoma (Rb)-associated protein 46 (RbAp46) [131]. RbAp46 is a growth suppressor and a component of the histone modifying and remodeling complexes. RbAp46 inhibits transactivation by a GAL4-BRCA1 fusion protein [131]. Another Rb-associated protein, RbAp48 and Rb itself were also found to interact with the BRCT domain of BRCA1 [8]. In addition, the BRCT domain interacts with proteins implicated in chromatin remodeling including the histone deacetylase (HDAC) 1 and 2 [8] or the B-related factor 1 (BRF1) subunit of the SWI/SNF-related complex [132]. The role of BRCA1 in chromatin remodeling has been shown by additional reports: BRCA1 physically interacts with the paralogous histone acetyltransferases (HATs), p300 and CBP [128,129]. These interactions enhance transcriptional activity of BRCA1 [129,132].

ER α , a member of the steroid hormone receptor superfamily, is activated by estrogen and plays important roles in normal development and breast tumorigenesis [133]. There are many reports demonstrating that most ER α -associated proteins participate in chromatin remodeling or in the recruitment of co-factors at the level of transcription initiation [120]. BRCA1 interacts with ER α and represses ER α -mediated transcriptional activity either in an estradiol (E2)-dependent [134–136] or -independent manner [137]. The ligand-dependent repression of ER α by BRCA1 is postulated to occur by binding of the N-terminus of BRCA1 (aa 1–302) to the C-terminal activation function (AF-2) domain of ER α [134,135]. BRCA1 may regulate the availability of p300 or cyclin D1 to compete for the AF-2 domain of ER α [138,139]. Overexpression of BRCA1 was reported to inhibit the induction of >90% of estrogen-inducible genes [140]. Given that ligand-independent repression of ER α transcription by BRCA1 was reversed by an HDAC inhibitor, trichostatin A (TSA), it is possible that HDAC activity may be involved in this regulation [137].

BRCA1 has been also known to be a transcriptional corepressor by interacting with the oncogenic transcription factor, c-Myc. BRCA1 interacts with the helix-loop-helix region of c-Myc through its two N-terminal regions (aa 175–303 and 443–511) and inhibits the transcriptional activity of c-Myc in the CDC25A promoter [141]. In addition, BRCA1 reverses the transforming phenotype of embryonic fibroblasts by the activation of c-Myc and Ras [141]. BRCA1 was also found to interact with the N-Myc-interacting protein (Nmi) through two distinct regions (aa 298–693 and 1301–1863) [142]. The formation of a ternary complex Nmi-c-Myc-BRCA1 is thought to inhibit c-Myc-induced transcription of the human telomerase reverse transcriptase (hTERT) promoter [142]. More recently, a number of genes have been identified as downregulated targets of BRCA1 through its interaction with c-Myc [143].

BRCA1 was also found to interact with CtIP [8–12]. The CtIP is an interacting partner of the transcriptional repressor CtBP (carboxyl-terminal binding protein) and is implicated in the repression of gene transcription [144]. The C-terminal region (aa 1602–1863) of BRCA1 was identified as a binding region to CtIP [8–12]. One proposed model is that phosphorylation of BRCA1 and CtIP by ATM following DNA damage results in dissociation of BRCA1 and CtIP and BRCA1-mediated upregulation of the growth arrest and DNA-damage-inducible gene 45 α (GADD45 α) protein [145]. However, dissociation of BRCA1-CtIP complex by DNA damage remains controversial [12,146].

The zinc finger and BRCA1-interacting protein with a KRAB domain 1 (ZBRK1) is a sequence-specific transcriptional corepressor, which binds to the GADD45 α intron 3 as a complex with BRCA1 [147]. The C-terminal repression domain of ZBRK1 functions in a BRCA1-, HDAC-, and promoter-specific manner, while the N-terminal KRAB repression domain of ZBRK1 exhibits no BRCA1 dependency and broad promoter specificity [148]. Additionally, the angiopoietin-1 (ANG1) and the high mobility group AT-hook 2 (HMGA2) have been identified as targets of the ZBRK1/CtIP/BRCA1 repressor complex [149,150]. The derepression of transcriptional repression of these genes by removing the ZBRK1/CtIP/BRCA1 repressor complex has been reported to enhance mammary tumor growth or tumorigenesis [149,150]. BRCA1 also induces GADD45 α transcription through interaction with transcription factors such as the octamer-binding transcription factor-1 (OCT-1) and the nuclear transcription factor-Y α (NF-YA) [151].

BRCA1 has been found to induce p21 in response to interferon gamma (IFN γ) through its interaction with the signal transducer and activator of transcription 1 (STAT1) [111]. Many, not all, of the IFN γ -responsive genes were identified as targets of BRCA1 and the induction of some of them was

synergistically enhanced by IFN γ [152]. Among them, the interferon regulator factor 7 (IRF7) is a key molecule in the interferon signaling in response to viral infection [153]. BRCA1 was also found to bind p65/RelA subunit of the nuclear factor- κ B (NF- κ B) and stimulate the tumor necrosis factor- α (TNF- α) and the interleukin-1 β (IL-1 β)-induced transcription of NF- κ B-target gene promoters [154]. It has been also reported that the interaction of BRCA1 and NF- κ B is necessary to efficiently activate NF- κ B activity after camptothecin treatment [155]. NF- κ B was also known to constitute a complex with BRCA1-CtIP to regulate DNA DSB repair [156]. These results suggest another role of BRCA1 in the immune response to viral infection [120] as well as tumor immune surveillance [136].

It has been also reported that BRCA1 binds and stabilizes the hypoxia-inducible factor-1 α (HIF-1 α) and activates its target gene vascular endothelial growth factor (VEGF) [157]. Under hypoxic condition, BRCA1 was recruited to the VEGF promoter along with HIF-1 α . Consistent with this, siRNA-based BRCA1-KD reduced the VEGF secretion in cultured cells following hypoxia [157]. BRCA1 was also known to repress estrogen-induced VEGF expression and secretion through interaction with ER α [158].

Expression and activation of the insulin-like growth factor-1 receptor (IGF-1R) is commonly detected in various human cancers [133,159–161]. The activated IGF-1R has been reported to be associated with poor survival of breast cancer patients [161]. Overexpression of wild type BRCA1 suppressed the IGF-1R promoter activity in various cell lines and reduced the endogenous levels of IGF-1R mRNA [162]. The N-terminal region of BRCA1 (aa 260–802) was reported to bind Sp1 transcription factor that strongly stimulates IGF-1R transcription [163]. Electrophoretic mobility shift assays (EMSAs) demonstrated that BRCA1-Sp1 interaction reduced the DNA-binding of Sp1 in a dose-dependent manner [163]. More recently, it has been reported that BRCA1 negatively regulates IGF-1 expression through an estrogen responsive element-like (EREL) site of IGF-1 promoter in an estradiol-dependent manner [164].

BRCA1 also modulates transcriptional activity of diverse transcription factors including the activating transcription factor 1 (ATF1) [165], the four and a half LIM only protein 2 (FHL2) [166], and so on. The significance of these regulations in the tumor suppressor function of BRCA1 remains to be determined. In the long list of BRCA1-interacting transcription factors (Table 1), several proteins, such as AhR/ARNT and NRF2, actively regulate the transcription of various genes in the oxidative stress response. The interactions of BRCA1 with these transcription factors in response to oxidative stress are discussed in following section.

Table 1. Transcription factors that bind to BRCA1.

Transcription factor	Region in BRCA1 (aa)	Assays	References
AhR	1241–1530	Co-IP/GST	[167]
AR (androgen receptor)	758–1064 & 1314–1863	GST/M2H	[168]
ARNT	131–757	Co-IP/GST/GAL4	[169]
ATF1	1–304	Co-IP/GST/Y2H/GAL4	[165]
BRF1	N/D	AP/GST	[132]
CBP	1–300 & 1314–1863	GST	[129]
CtIP	1602–1863	Co-IP/GST/Y2H	[10]
E2F1	N/D	GST	[170]
E2F4	N/D	GST	[170]

Table 1. Cont.

Transcription factor	Region in BRCA1 (aa)	Assays	References
ER α	1–300	Co-IP/GST	[135]
FHL2	1756–1852	GST/GAL4/Y2H	[166]
HDAC1	1553–1863	GST	[8]
HDAC2	1553–1863	GST	[8]
HIF-1 α	N/D	Co-IP	[157]
JunB	1343–1440	Co-IP/GST/GAL4/Y2H	[171]
JunD	AD1	Co-IP/GST/GAL4/Y2H	[171]
c-Jun	AD1	Co-IP/GST/GAL4	[171]
LMO4	1528–1863	Co-IP/GST/Y2H	[172]
c-Myc	175–303 & 443–511	Co-IP/GST/Y2H	[141]
NF-YA	N/D	Co-IP	[151]
Nmi	298–693 & 1301–1863	Co-IP/GST/Y2H	[142]
NRF2	N/D	Co-IP	[117]
OCT-1	N/D	Co-IP	[151]
p300	1528–1863	Co-IP/GAL4	[129]
p53	224–550 & 1760–1863	Co-IP/GST	[86,87]
RbAp46	1553–1863	Co-IP/GST/GAL4/Y2H	[8,131]
RbAp48	1553–1863	Co-IP/GST	[8]
RelA	1–802	Co-IP/GST	[154]
RHA	1650–1800	GST/GAL4/Y2H	[128]
Sp1	260–802	Co-IP/GST	[163]
STAT1	N/D	Co-IP/GST/M2H	[111]
STAT5A	N/D	Co-IP	[173]
USF2	N/D	Co-IP	[174]
ZBRK1	341–748	Co-IP/GST/Y2H	[147]

Abbreviations, AP: affinity purification; Co-IP: co-immunoprecipitation; GAL4: GAL4 reporter gene assay; GST: GST pull-down; M2H: mammalian two hybrid; N/D: not determined; Y2H: yeast two hybrid.

4.2. BRCA1 Regulation of Oxidative Stress through Transcriptional Activities

In addition to all the functions of BRCA1, it regulates the expression of detoxification enzymes including the UDP-glucuronosyltransferase isoform 1A1 (UGT1A1), UGT1A9, and NRF2 through transcriptional regulation by binding to their promoter regions in response to B[a]P [175]. All these genes contain xenobiotic responsive elements (XREs) in their promoter regions and are known to be regulated by a transcription factor aryl hydrocarbon receptor (AhR) [175]. Induction of NRF2 by AhR was also reported previously [176]. The AhR is a major transcriptional activator in xenobiotic responses. Upon ligand binding, AhR forms a complex with the AhR nuclear translocator (ARNT) to translocate into the nucleus and activate the transcription of its target genes [177]. BRCA1 is known to bind AhR and ARNT to activate the transcription of their target genes in breast cancer cells [167,169]. The N-terminal region of BRCA1 (aa 131–757) may be sufficient to interact with ARNT [169] and the C-terminal region (aa 1241–1530) containing the activation domain 1 (AD1) of BRCA1 has been also found to interact with AhR [167]. Interaction between BRCA1 and AhR was enhanced by xenobiotic TCDD in breast cancer cells [167] and BRCA1 was recruited to the promoter regions of CYP1A1 and

CYP1B1 along with ARNT/AhR following TCDD exposure [169]. It is also reported that BRCA1 is recruited to the promoter regions of UGT1A1, UGT1A9 and NRF2 in human normal-like breast MCF10A cells after B[a]P treatment [175]. Surprisingly, BRCA1-KD reduced the binding of ARNT to XRE regions of these promoters in chromatin immunoprecipitation assays. As expected, wild type BRCA1 further activated the XRE-dependent transcriptional activation in the presence of B[a]P in MCF10A cells. Consistently, BRCA1-KD abolished the B[a]P-induced transcriptional activity of UGT1A1 and NRF2 promoters in MCF10A cells [175].

Systematic comparisons of gene expression profiles have revealed that BRCA1 regulates expression of a wide range of genes in human prostate (DU-145) and breast (MCF7) cancer cells [178,179]. Importantly, overexpression of BRCA1, in either DU-145 or MCF7 cells, up-regulates several NRF2-target genes including NQO1, MGST1/2, GSTA2, G6PC, ME2, *etc.* [179]. Consistently, antioxidant response genes were downregulated in *Brcal*-deficient MEF cell lines [179]. As a transcription factor, NRF2 binds to a specific DNA sequence known as antioxidant response element (ARE) in the promoter regions of its target genes and activates transcription of these genes [180]. Interestingly, reporter gene assays utilized with an ARE-luciferase (Luc) construct demonstrated that overexpression of wild type BRCA1 enhanced NRF2-dependent transcription [179] and BRCA1-KD by siRNA reduced the NRF2-dependent transcription in various cell lines [78,179]. Reporter gene assays with expression of different BRCA1 mutants suggest that the *N*-terminal domain of BRCA1 is necessary and sufficient to stimulate NRF2-dependent transcriptional activation of the ARE-Luc reporter [179]. Overexpression of wild type BRCA1 conferred resistance of DU-145 and *Brcal*-defective MEF cells to oxidative stress-inducing agents (H₂O₂ and paraquat); while BRCA1-KD sensitized DU-145 cells to these agents in MTT cell viability assay [179]. Wild type BRCA1 also attenuated the loss of glutathione (GSH) in response to H₂O₂ in human cancer cell lines, DU-145, LNCaP and MCF7 [179]. Interestingly, a recent study reported that NRF2 induced the transcription of the *BRCA1* gene through binding to an ARE in the *BRCA1* gene promoter [181].


5. BRCA1, an ROS Sensor?

Despite all these results supporting the role of BRCA1 in the oxidative stress response, the mechanism of ROS sensing by BRCA1 still remains to be determined. Since BRCA1 is extensively phosphorylated at multiple residues by various kinases including ATM, ATR, the DNA-dependent protein kinase (DNA-PK), the checkpoint kinase 2 (CHK2), the cyclin-dependent kinases (CDKs), the aurora kinase A (AURK), and AKT (reviewed in [62,182–184]), it is possible that these multiple phosphorylations may activate BRCA1 function in response to ROS. In fact, recent evidence supports that oxidative stress activates diverse protein kinases including ATM, ATR, and DNA-PK ([185–189] and reviewed in [25–27]). In addition, oxidative stress triggers various post-translation modifications of proteins such as methylation, acetylation, sumoylation, nitrosylation, nitration, glutathionylation, and oxidation [190–192]. Thus, it stands to reason that BRCA1 and its associated kinases sense oxidative stress and activate tumor suppression activities. Future studies of BRCA1 modifications by oxidative stress are likely to help our understanding of BRCA1 function in the oxidative stress response and tumor suppression.

6. Conclusions

Regulation of oxidative stress by BRCA1 has been less recognized than its other functional aspects. However, there is considerable mounting evidence suggesting that BRCA1 actively regulates ROS in response to oxidative stress at multiple levels (Figure 2). Since oxidative stress causes carcinogenesis through genomic instability by way of DNA-damage and uncontrolled cell proliferation, regulation of the oxidative stress response by BRCA1 may constitute a wheel of the chariot of the BRCA1 tumor suppression.

Figure 2. BRCA1 in the oxidative stress response.



ROS have the Janus faces in human cancers: uncontrolled ROS accelerate carcinogenesis in normal cells; and increased ROS may directly induce apoptotic death of cancer cells [30,193]. Consistently, antioxidant phytochemicals can be chemopreventive in early stage of carcinogenesis and can protect tumor cells from oxidative injuries [34,193]. Recently, targeting ROS has been revisited as a general therapeutic approach to treat cancers [193–196]. As an example, piperlongumine selectively induces the ROS level and apoptosis in cancer cells, but not in normal cells [197]. In fact, many conventional cancer therapies, either chemotherapeutics or radiotherapy, induce ROS in cancer cells [193–196]. Future studies will be focused on how specifically induce ROS in different types of cancers.

Taken together, the emerging role of BRCA1 as a sensor and regulator of ROS has potentially important implications to prevent and/or treat BRCA1-associated tumors. As shown in our recent study [79], surveillance of environmental factors in the context of the oxidative stress (or ROS production) in BRCA1-defective cells may provide a valuable tool to assess the risks of various environmental factors that can potentially induce BRCA1-associated carcinogenesis. This approach will be more helpful to determine potential carcinogens from the variety of environmental factors in conjunction with *Brc1* mouse models, which spontaneously develop *Brc1*-associated mammary gland tumors [65–68]. In addition, this approach can be also applied to identify potential chemopreventive agents. For example, antioxidant phytochemicals such as sulforaphane or resveratrol can be helpful to prevent breast or ovarian cancers that carrying BRCA1 mutations by removing ROS. On the other hand, increasing

cellular ROS production may be utilized to treat BRCA1-associated cancers. Indeed, BRCA1 mutated cancers are particularly sensitive to genotoxic and oxidative agents including PARP inhibitors and chemotherapeutics [198–201]. Thus, specific targeting of oxidative stress in BRCA1-defective tumors may provide alternative treatment options [202,203].

Acknowledgments

This work was supported by the Susan G. Komen for the Cure (FAS0703858) and by the Lombardi Comprehensive Cancer Center, Georgetown University Medical Center (P30-CA051008). We appreciate BioMedText, Inc./Rashmi Nemade for helpful discussions and editing.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Futreal, P.A.; Liu, Q.; Shattuck-Eidens, D.; Cochran, C.; Harshman, K.; Tavtigian, S.; Bennett, L.M.; Haugen-Strano, A.; Swensen, J.; Miki, Y.; *et al.* BRCA1 mutations in primary breast and ovarian carcinomas. *Science* **1994**, *266*, 120–122.
2. Hall, J.M.; Lee, M.K.; Newman, B.; Morrow, J.E.; Anderson, L.A.; Huey, B.; King, M.C. Linkage of Early-Onset Familial Breast Cancer to Chromosome 17q21. *Science* **1990**, *250*, 1684–1689.
3. Easton, D.F.; Bishop, D.T.; Ford, D.; Crockford, G.P. Genetic linkage analysis in familial breast and ovarian cancer: Results from 214 families. The breast cancer linkage consortium. *Am. J. Hum. Genet.* **1993**, *52*, 678–701.
4. Miki, Y.; Swensen, J.; Stattuck-Eidens, D.; Futreal, P.A.; Harshman, K.; Tavtigian, S.; Liu, Q.; Cochran, C.; Bennett, L.M.; Ding, W.; *et al.* A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science* **1994**, *266*, 66–71.
5. Smith, T.M.; Lee, M.K.; Szabo, C.I.; Jerome, N.; McEuen, M.; Taylor, M.; Hood, L.; King, M.C. Complete genomic sequence and analysis of 117 kb of human DNA containing the gene BRCA1. *Genome Res.* **1996**, *6*, 1029–1049.
6. Wu, L.C.; Wang, Z.W.; Tsan, J.T.; Spillman, M.A.; Phung, A.; Xu, X.L.; Yang, M.C.; Hwang, L.Y.; Bowcock, A.M.; Baer, R. Identification of a RING protein that can interact *in vivo* with the BRCA1 gene product. *Nat. Genet.* **1996**, *14*, 430–440.
7. Jensen, D.E.; Proctor, M.; Marquis, S.T.; Gardner, H.P.; Ha, S.I.; Chodosh, L.A.; Ishov, A.M.; Tommerup, N.; Vissing, H.; Sekido, Y.; *et al.* BAP1: A novel ubiquitin hydrolase which binds to the BRCA1 RING finger and enhances BRCA1-mediated cell growth suppression. *Oncogene* **1998**, *16*, 1097–1112.
8. Yarden, R.I.; Brody, L.C. BRCA1 interacts with components of the histone deacetylase complex. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 4983–4988.

9. Yu, X.; Wu, L.C.; Bowcock, A.M.; Aronheim, A.; Baer, R. The C-terminal (BRCT) domains of BRCA1 interact *in vivo* with CtIP, a protein implicated in the CtBP pathway of transcriptional repression. *J. Biol. Chem.* **1998**, *273*, 25388–25392.
10. Li, S.; Chen, P.L.; Subramanian, T.; Chinnadurai, G.; Tomlinson, G.; Osborne, C.K.; Sharp, Z.D.; Lee, W.H. Binding of CtIP to the BRCT repeats of BRCA1 involved in the transcription regulation of p21 is disrupted upon DNA damage. *J. Biol. Chem.* **1999**, *274*, 11334–11338.
11. Wong, A.K.; Ormonde, P.A.; Pero, R.; Chen, Y.; Lian, L.; Salada, G.; Berry, S.; Lawrence, Q.; Dayananth, P.; Ha, P.; *et al.* Characterization of a carboxy-terminal BRCA1 interacting protein. *Oncogene* **1998**, *17*, 2279–2285.
12. Yu, X.; Baer, R. Nuclear localization and cell cycle-specific expression of CtIP, a protein that associates with the BRCA1 tumor suppressor. *J. Biol. Chem.* **2000**, *275*, 18541–18549.
13. Paull, T.T.; Cortez, D.; Bowers, B.; Elledge, S.J.; Gellert, M. Direct DNA binding by Brca1. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 6086–6091.
14. Simons, A.M.; Horwitz, A.A.; Starita, L.M.; Griffin, K.; Williams, R.S.; Glover, J.N.; Parvin, J.D. BRCA1 DNA-binding activity is stimulated by BARD1. *Cancer Res.* **2006**, *66*, 2012–2018.
15. Cortez, D.; Wang, Y.; Qin, J.; Elledge, S.J. Requirement of ATM-dependent phosphorylation of brca1 in the DNA damage response to double-strand breaks. *Science* **1999**, *286*, 1162–1166.
16. Chen, Y.; Chen, C.F.; Riley, D.J.; Allred, D.C.; Chen, P.L.; von Hoff, D.; Osborne, C.K.; Lee, W.H. Aberrant subcellular localization of BRCA1 in breast cancer. *Science* **1995**, *270*, 789–791.
17. Chen, C.F.; Li, S.; Chen, Y.; Chen, P.L.; Sharp, Z.D.; Lee, W.H. The nuclear localization sequences of the BRCA1 protein interact with the importin- α subunit of the nuclear transport signal receptor. *J. Biol. Chem.* **1996**, *271*, 32863–32868.
18. Rodríguez, J.A.; Henderson, B.R. Identification of a functional nuclear export sequence in BRCA1. *J. Biol. Chem.* **2000**, *275*, 38589–38596.
19. Fabbro, M.; Rodríguez, J.A.; Baer, R.; Henderson, B.R. BARD1 induces BRCA1 intranuclear foci formation by increasing RING-dependent BRCA1 nuclear import and inhibiting BRCA1 nuclear export. *J. Biol. Chem.* **2002**, *277*, 21315–21324.
20. Feng, Z.; Kachnic, L.; Zhang, J.; Powell, S.N.; Xia, F. DNA damage induces p53-dependent BRCA1 nuclear export. *J. Biol. Chem.* **2004**, *279*, 28574–28584.
21. Bennett, L.M.; Haugen-Strano, A.; Cochran, C.; Brownlee, H.A.; Fiedorek, F.T., Jr.; Wiseman, R.W. Isolation of the mouse homologue of BRCA1 and genetic mapping to mouse chromosome 11. *Genomics* **1995**, *29*, 576–581.
22. Jin, H.; Selfe, J.; Whitehouse, C.; Morris, J.R.; Solomon, E.; Roberts, R.G. Structural evolution of the BRCA1 genomics region in primates. *Genomics* **2004**, *84*, 1071–1082.
23. Rosen, E.M.; Fan, S.; Pestell, R.G.; Goldberg, I.D. The BRCA1 gene in breast cancer. *J. Cell. Physiol.* **2003**, *196*, 19–41.
24. Silver, D.P.; Livingston, D.M. Mechanism of BRCA1 tumor suppression. *Cancer Discov.* **2012**, *2*, 679–684.
25. Finkel, T. Signal transduction by reactive oxygen species. *J. Cell Biol.* **2011**, *194*, 7–15.
26. Ma, Q. Transcriptional responses to oxidative stress: Pathological and toxicological implications. *Pharmacol. Ther.* **2010**, *125*, 376–393.

27. Ray, P.D.; Huang, B.W.; Tsuji, Y. Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cell. Signal.* **2012**, *24*, 981–990.
28. Corcoran, A.; Cotter, T.G. Redox regulation of protein kinases. *FEBS J.* **2013**, *280*, 1944–1965.
29. Jennings, P.; Limonciel, A.; Felice, L.; Leonard, M.O. An overview of transcriptional regulation in response to toxicological insult. *Arch. Toxicol.* **2013**, *87*, 49–72.
30. Acharya, A.; Das, I.; Chandhok, D.; Saha, T. Redox regulation in cancer: A double-edged sword with therapeutic potential. *Oxid. Med. Cell. Logev.* **2010**, *3*, 23–34.
31. Kryston, T.B.; Georgiev, A.B.; Pissis, P.; Georgakila, A.G. Role of oxidative stress and DNA damage in human carcinogenesis. *Mutat. Res.* **2011**, *711*, 193–201.
32. Caputo, F.; Vegliante, R.; Ghibelli, L. Redox modulation of the DNA damage response. *Biochem. Pharmacol.* **2012**, *84*, 1292–1306.
33. Storr, S.J.; Woolston, C.M.; Zhang, Y.; Martin, S.G. Redox environment, free radical, and oxidative DNA damage. *Antioxid. Redox Signal.* **2013**, *18*, 2399–2408.
34. Saeidnia, S.; Abdollahi, M. Antioxidants: Friends or foe in prevention or treatment of cancer: The debate of the century. *Toxicol. Appl. Pharmacol.* **2013**, *271*, 49–63.
35. Kakehashi, A.; Wei, M.; Fukushima, S.; Wanibuchi, H. Oxidative stress in the carcinogenicity of chemical carcinogens. *Cancers* **2013**, *5*, 1332–1354.
36. Bossy-Wetzell, E.; Schwarzenbacher, R.; Lipton, S.A. Molecular pathways to neurodegeneration. *Nat. Med.* **2004**, *10*, S2–S9.
37. Balaban, R.S.; Nemoto, S.; Finkel, T. Mitochondria, oxidants, and aging. *Cell* **2005**, *120*, 483–495.
38. Jimenez-Del-Rio, M.; Velez-Pardo, C. The bad, the good, and the ugly about oxidative stress. *Oxid. Med. Cell. Logev.* **2012**, doi:10.1155/2012/163913.
39. King, M.C.; Marks, J.H.; Mandell, J.B. Breast and ovarian cancer risk due to inherited mutations in BRCA1 and BRCA2. *Science* **2003**, *302*, 643–646.
40. Wilson, C.A.; Ramos, L.; Villaseñor, M.R.; Anders, K.H.; Press, M.F.; Clarke, K.; Karlan, B.; Chen, J.J.; Scully, R.; Livingston, D.; *et al.* Localization of human BRCA1 and its loss in high-grade, non-inherited breast carcinomas. *Nat. Genet.* **1999**, *21*, 236–240.
41. Esteller, M.; Silva, J.M.; Dominguez, G.; Bonilla, F.; Matisa-Guiu, X.; Lerma, E.; Bussaglia, E.; Prat, J.; Harkes, I.C.; Repasky, E.A.; *et al.* Promoter hypermethylation and BRCA1 inactivation in sporadic breast and ovarian tumors. *J. Natl. Cancer Inst.* **2000**, *95*, 564–569.
42. Thompson, D.; Easton, D.; Breast Cancer Linkage Consortium. Variation in BRCA1 cancer risks by mutation position. *Cancer Epidemiol. Biomark. Prev.* **2002**, *11*, 329–336.
43. Leongamornlert, D.; Mahmud, N.; Tymrakiewicz, M.; Saunders, E.; Dadaev, T.; Castro, E.; Goh, C.; Govindasami, K.; Guy, M.; O'Brien, L.; *et al.* Germline BRCA1 mutations increase prostate cancer risk. *Br. J. Cancer* **2012**, *106*, 1697–1701.
44. Li, D.; Kumaraswamy, E.; Harlan-Williams, L.M.; Jensen, R.A. The role of BRCA1 and BRCA2 in prostate cancer. *Front. Biosci.* **2013**, *18*, 1145–1159.
45. Johannsson, O.; Ostermeyer, E.A.; Håkansson, S.; Friedman, L.S.; Johannsson, U.; Sellberg, G.; Brøndum-Nielsen, K.; Sele, V.; Olsson, H.; King, M.C.; *et al.* Founding BRCA1 mutations in hereditary breast and ovarian cancer in southern Sweden. *Am. J. Hum. Genet.* **1996**, *58*, 441–450.

46. Lal, G.; Liu, G.; Schmocker, B.; Kaurah, P.; Ozcelik, H.; Narod, S.A.; Redston, M.; Gallinger, S. Inherited predisposition to pancreatic adenocarcinoma: Role of family history and germ-line p16, BRCA1, and BRCA2 mutations. *Cancer Res.* **2000**, *60*, 409–416.
47. Thompson, D.; Easton, D.F.; Breast Cancer Linkage Consortium. Cancer incidence in BRCA1 mutation carriers. *J. Natl. Cancer Inst.* **2002**, *94*, 1358–1365.
48. Lynch, H.T.; Deters, C.A.; Snyder, C.L.; Lynch, J.F.; Villeneuve, P.; Silberstein, J.; Martin, H.; Narod, S.A.; Brand, R.E. BRCA1 and pancreatic cancer: Pedigree findings and their causal relationships. *Cancer Genet. Cytogenet.* **2005**, *158*, 119–125.
49. Dagan, E. Predominant Ashkenazi BRCA1/2 mutations in families with pancreatic cancer. *Genet. Test.* **2008**, *12*, 267–271.
50. Al-Sukhni, W.; Rothenmund, H.; Borgida, A.E.; Zogopoulos, G.; O’Shea, A.M.; Pollett, A.; Gallinger, S. Germline BRCA1 mutations predispose to pancreatic adenocarcinoma. *Hum. Genet.* **2008**, *124*, 271–278.
51. Kim, D.H.; Crawford, B.; Ziegler, J.; Beattie, M.S. Prevalence and characteristic cancer in families with BRCA1 and BRCA2 mutations. *Fam. Cancer* **2009**, *8*, 153–158.
52. Stadler, Z.K.; Salo-Mullen, E.; Patil, S.M.; Pietanza, M.C.; Vijai, J.; Saloustros, E.; Hansen, N.A.; Kauff, N.D.; Kurtz, R.C.; Kelsen, D.P.; *et al.* Prevalence of BRCA1 and BRCA2 mutations in Ashkenazi Jewish families with breast and pancreatic cancer. *Cancer* **2012**, *118*, 493–499.
53. Igbal, J.; Ragone, A.; Lubinski, J.; Lynch, H.T.; Moller, P.; Ghadirian, P.; Foulkes, W.D.; Armel, S.; Eisen, A.; Neuhausen, S.L.; *et al.* The incidence of pancreatic cancer in BRCA1 and BRCA2 mutation carriers. *Br. J. Cancer* **2012**, *107*, 2005–2009.
54. Brodie, S.G.; Deng, C.X. BRCA1-associated tumorigenesis: What have we learned from knockout mice? *Trends Genet.* **2001**, *17*, S18–S22.
55. Deng, C.X.; Brodie, S.G. Knockout mouse models and mammary tumorigenesis. *Semin. Cancer Biol.* **2001**, *11*, 387–394.
56. Drost, R.; Bouwman, P.; Rottenberg, S.; Boon, U.; Schut, E.; Klarenbeek, S.; Klijin, C.; van der Heijden, I.; van der Gulden, H.; Wientjens, E.; *et al.* BRCA1 RING function is essential for tumor suppression but dispensable for therapy resistance. *Cancer Cell* **2011**, *20*, 797–809.
57. Shakya, R.; Reid, L.J.; Reczek, C.R.; Cole, F.; Egli, D.; Lin, C.S.; deRoos, D.G.; Hirsch, S.; Ravi, K.; Hicks, J.B.; *et al.* BRCA1 tumor suppression depends on BRCT phosphoprotein binding, but not its E3 ligase activity. *Science* **2011**, *334*, 525–528.
58. Hashizume, R.; Fukuda, M.; Maeda, I.; Nishikawa, H.; Oyake, D.; Yabuki, Y.; Ogata, H.; Ohta, T. The RING heterodimer BRCA1-BARD1 is a ubiquitin ligase inactivated by a breast cancer-derived mutation. *J. Biol. Chem.* **2001**, *276*, 14537–14540.
59. Ruffner, H.; Joazeiro, C.A.; Hemmati, D.; Hunter, T.; Verma, I.M. Cancer-predisposing mutations within the RING domain of BRCA1: Loss of ubiquitin protein ligase activity and protection from radiation hypersensitivity. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 5134–5139.
60. Mallery, D.L.; Vandenberg, C.J.; Hiom, K. Activation of the E3 ligase function of the BRCA1/BARD complex by polyubiquitin chains. *EMBO J.* **2002**, *21*, 6755–6762.
61. Baer, R.; Ludwig, T. The BRCA1/BARD1 heterodimer, a tumor suppressor complex with ubiquitin E3 ligase activity. *Curr. Opin. Genet. Dev.* **2002**, *12*, 86–91.

62. Lee, J.S.; Chung, J.H. Diverse functions of BRCA1 in the DNA damage response. *Expert Rev. Mol. Med.* **2001**, *2001*, 1–11.
63. Yun, M.H.; Hiom, K. Understanding the functions of BRCA1 in the DNA-damage response. *Biochem. Soc. Trans.* **2009**, *37*, 597–604.
64. Dunnick, J.K.; Elwell, M.R.; Hugg, J.; Barrett, J.C. Chemically induced mammary gland cancer in the National Toxicology Program's carcinogenesis bioassay. *Carcinogenesis* **1995**, *16*, 173–179.
65. Xu, X.; Qiao, W.; Linke, S.P.; Cao, L.; Li, W.M.; Furth, P.A.; Harris, C.C.; Deng, C.X. Genetic interactions between tumor suppressors Brca1 and p53 in apoptosis, cell cycle and tumorigenesis. *Nat. Genet.* **2001**, *28*, 266–271.
66. Bachelier, R.; Xu, X.; Wang, X.; Li, W.; Naramura, M.; Gu, H.; Deng, C.X. Normal lymphocyte development and thymic lymphoma formation in Brca1 exon-11-deficient mice. *Oncogene* **2003**, *22*, 528–537.
67. Cao, L.; Li, W.; Kim, S.; Brodie, S.G.; Deng, C.X. Senescence, aging, and malignant transformation mediated by p53 in mice lacking the Brca1 full-length isoform. *Genes Dev.* **2003**, *17*, 201–213.
68. Cao, L.; Kim, S.; Xiao, C.; Wang, R.H.; Coumoul, X.; Wang, X.; Li, W.M.; Xu, X.L.; de Soto, J.A.; Takai, H.; *et al.* ATM-Chk2-p53 activation prevents tumorigenesis at an expense of organ homeostasis upon Brca1 deficiency. *EMBO J.* **2006**, *25*, 2167–2177.
69. Cao, L.; Xu, X.; Cao, L.L.; Wang, R.H.; Coumoul, X.; Kim, S.S.; Deng, C.X. Absence of full-length Brca1 sensitizes mice to oxidative stress and carcinogen-induced tumorigenesis in the esophagus and forestomach. *Carcinogenesis* **2007**, *28*, 1401–1407.
70. Francisco, D.C.; Peddi, P.; Hair, J.M.; Flood, B.A.; Cecil, A.M.; Kalogerinis, P.T.; Sigounas, G.; Georgakilas, A.G. Induction and processing of complex DNA damage in human breast cancer cells MCF-7 and nonmalignant MCF-10A cells. *Free Radic. Biol. Med.* **2008**, *44*, 558–569.
71. Hair, J.M.; Terzoudi, G.I.; Hatzi, V.I.; Lehouckey, K.A.; Srivastava, D.; Wang, W.; Pantelias, G.E.; Georgakilas, A.G. BRCA1 role in the mitigation of radiotoxicity and chromosomal instability through repair of clustered DNA lesions. *Chem. Biol. Interact.* **2010**, *188*, 350–358.
72. Wang, Y.; Cortez, D.; Yazdi, P.; Neff, N.; Elledge, S.J.; Qin, J. BASC, a super complex of BRCA1-associated proteins involved in the recognition and repair of aberrant DNA structures. *Genes Dev.* **2000**, *14*, 927–939.
73. Le Page, F.; Randrianarison, V.; Marot, D.; Cabannes, J.; Perricaudet, M.; Feunteun, J.; Sarasin, A. BRCA1 and BRCA2 are necessary for the transcription-coupled repair of the oxidative 8-oxoguanine lesion in human cells. *Cancer Res.* **2000**, *60*, 5548–5552.
74. Cadet, J.; Berger, M.; Douki, T.; Ravanat, J.L. Oxidative damage to DNA: Formation, measurement and biological significance. *Rev. Physiol. Biochem. Pharmacol.* **1997**, *131*, 1–87.
75. Dziaman, T.; Huzarski, T.; Gackowski, D.; Rozalski, R.; Siomek, A.; Szpila, A.; Jalanta, G.; Lubinski, J.; Olinski, R. Elevated level of 8-oxi-7,8-dihydro-2'-deoxyguanosine in leukocytes of BRCA1 mutation carriers compared to healthy controls. *Int. J. Cancer* **2009**, *125*, 2209–2213.
76. Saha, T.; Rih, J.K.; Roy, R.; Ballal, R.; Rosen, E.M. Transcriptional regulation of the base excision repair pathway by BRCA1. *J. Biol. Chem.* **2010**, *285*, 19092–19105.
77. Saha, T.; Rih, J.K.; Rosen, E.M. BRCA1 down-regulates cellular levels of reactive oxygen species. *FEBS Lett.* **2009**, *583*, 1535–1543.

78. Kang, H.J.; Hong, Y.B.; Kim, H.J.; Wang, A.; Bae, I. Bioactive food components prevent carcinogenic stress via Nrf2 activation in BRCA1 deficient breast epithelial cells. *Toxicol. Lett.* **2012**, *209*, 154–160.
79. Kang, H.J.; Hong, Y.B.; Yi, Y.W.; Cho, C.H.; Wang, A.; Bae, I. The correlation between BRCA1 defect and environmental factors in the risk of breast cancer. *J. Toxicol. Sci.* **2013**, *38*, 355–361.
80. Martinez-Outschoorn, U.E.; Balliet, R.; Lin, Z.; Whitaker-Menez, D.; Birbe, R.C.; Bombonati, A.; Pavlides, S.; Lamb, R.; Sneddon, S.; Howell, A.; *et al.* BRCA1 mutations drive oxidative stress and glycolysis in the tumor microenvironment: Implications for breast cancer prevention with antioxidant therapies. *Cell Cycle* **2012**, *11*, 4402–4413.
81. Martinez-Outschoorn, U.E.; Balliet, R.M.; Lin, Z.; Whitaker-Menezes, D.; Howell, A.; Sotgia, F.; Lisanti, M.P. Hereditary ovarian cancer and two-compartment tumor metabolism: Epithelial loss of BRCA1 induces hydrogen peroxide production, driving oxidative stress and NFκB activation in the tumor stroma. *Cell Cycle* **2012**, *11*, 1–15.
82. Lovren, F.; Pan, Y.; Quan, A.; Singh, K.K.; Gupta, N.; Brezden-Masley, C.; Teoh, H.; Wheatcroft, M.D.; Al-Omaran, M.; Verma, S. BRCA1 shields vascular smooth muscle cells from oxidative stress. *J. Thorac. Cardiovasc. Surg.* **2013**, doi:10.1016/j.jtcvs.2013.09.060.
83. Vurusaner, B.; Poli, G.; Basaga, H. Tumor suppressor genes and ROS: Complex networks of interaction. *Free Radic. Biol. Med.* **2012**, *52*, 7–18.
84. Somasundaram, K.; Zhang, H.; Zeng, Y.W.; Houvras, Y.; Peng, Y.; Zhang, H.; Wu, G.S.; Licht, J.D.; Wever, B.L.; El-Deiry, W.S. Tumour-suppressor BRCA1 requires the CDK-inhibitor p21^{WAF1/CIP1}. *Nature* **1997**, *389*, 187–190.
85. Ouchi, T.; Monteiro, A.N.; August, A.; Aaronson, S.A.; Hanafusa, H. BRCA1 regulates p53-dependent gene expression. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 2302–2306.
86. Zhang, H.; Somasundaram, K.; Peng, Y.; Tian, H.; Zhang, H.; Bi, D.; Weber, B.L.; El-Deiry, W.S. BRCA1 physically associated with p53 and stimulates its transcriptional activity. *Oncogene* **1998**, *16*, 1713–1721.
87. Chai, Y.L.; Cui, J.; Shao, N.; Shyam, E.; Reddy, P.; Rao, V.N. The second BRCT domain of BRCA1 protein interacts with p53 and stimulates transcription from the p21^{WAF1/CIP1} promoter. *Oncogene* **1999**, *18*, 263–268.
88. MacLachlan, T.K.; Takimoto, R.; El-Deiry, W.S. BRCA1 directs a selective p53-dependent transcriptional response towards growth arrest and DNA repair targets. *Mol. Cell. Biol.* **2002**, *22*, 4280–4292.
89. Somasundaram, K.; MacLachlan, T.K.; Burns, T.F.; Sgagias, M.; Cowan, K.H.; Weber, B.L.; El-Deiry, W.S. BRCA1 signals ARF-dependent stabilization and coactivation of p53. *Oncogene* **1999**, *18*, 6605–6614.
90. Fabbro, M.; Savage, K.; Hobson, K.; Deans, A.J.; Powell, S.N.; McArthur, G.A.; Khanna, K.K. BRCA1-BARD1 complexes are required for p53^{Ser-15} phosphorylation and a G₁/S arrest following ionizing radiation-induced DNA damage. *J. Biol. Chem.* **2004**, *279*, 31251–31258.

91. Velimezi, G.; Liontos, M.; Vougas, K.; Roumeliotis, T.; Bartkova, J.; Sideridou, M.; Dereli-Oz, A.; Kocylowski, M.; Pateras, I.S.; Evangelou, K.; *et al.* Functional interplay between the DNA-damage-response kinase ATM and ARF tumour suppressor protein in human cancer. *Nat. Cell Biol.* **2013**, *15*, 967–977.
92. Bedford, L.; Lowe, J.; Dick, L.R.; Mayer, R.J.; Brownell, J.E. Ubiquitin-like protein conjugation and the ubiquitin-proteasome system as drug targets. *Nat. Rev. Drug Discov.* **2011**, *10*, 29–46.
93. Chau, V.; Tobias, J.W.; Bachmair, A.; Marriott, D.; Ecker, D.J.; Gonda, D.K.; Varshavsky, A. A multiubiquitin chain is confined to specific lysine in a targeted short-lived proteins. *Science* **1989**, *243*, 1576–1583.
94. Wu-Baer, F.; Lagrazon, K.; Yuan, W.; Baer, R. The BRCA1/BARD1 heterodimer assembles polyubiquitin chains through an unconventional linkage involving lysine residue K6 of ubiquitin. *J. Biol. Chem.* **2003**, *278*, 34743–34746.
95. Nishikawa, H.; Ooka, S.; Sato, K.; Arima, K.; Okamoto, J.; Klevit, R.E.; Fukuda, M.; Ohta, T. Mass spectrometric and mutational analyses reveal Lys-6-linked polyubiquitin chains catalyzed by BRCA1-BARD1 ubiquitin ligase. *J. Biol. Chem.* **2004**, *279*, 3916–3924.
96. Morris, J.R.; Solomon, E. BRCA1:BARD1 induces the formation of conjugated ubiquitin structures, dependent on K6 of ubiquitin, in cells during DNA replication and repair. *Hum. Mol. Genet.* **2004**, *13*, 807–817.
97. Sato, K.; Hayami, R.; Wu, W.; Nishikawa, T.; Nishikawa, H.; Okuda, Y.; Ogata, H.; Fukuda, M.; Ohta, T. Nucleophosmin/B23 is a candidate substrate for the BRCA1-BARD1 ubiquitin ligase. *J. Biol. Chem.* **2004**, *279*, 30919–30922.
98. Yu, X.; Fu, S.; Lai, M.; Baer, R.; Chen, J. BRCA1 ubiquitinates its phosphorylation-dependent binding partner CtIP. *Genes Dev.* **2006**, *20*, 1721–1726.
99. Wu, W.; Nishikawa, H.; Hayami, R.; Sato, K.; Honda, A.; Aratani, S.; Nakajima, T.; Fukuda, M.; Ohta, T. BRCA1 ubiquitinates RPB8 in response to DNA damage. *Cancer Res.* **2007**, *67*, 951–958.
100. Horwitz, A.A.; Affar, E.B.; Heine, G.F.; Shi, Y.; Parvin, J.D. A mechanism for transcriptional repression dependent on the BRCA1 E3 ubiquitin ligase. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 6614–6619.
101. Chen, A.; Kleiman, F.E.; Manley, J.L.; Ouchi, T.; Pan, Z.Q. Autoubiquitination of the BRCA1-BARD1 RING ubiquitin ligase. *J. Biol. Chem.* **2002**, *277*, 22085–22092.
102. Wu, W.; Koike, A.; Takeshita, T.; Ohta, T. The ubiquitin E3 ligase activity of BRCA1 and its biological functions. *Cell Division* **2008**, *3*, 1.
103. Roy, R.; Chun, J.; Powell, S.M. BRCA1 and BRCA2: Different roles in a common pathway of genome protection. *Nat. Rev. Cancer* **2012**, *12*, 68–78.
104. Shabbeer, S.; Omer, D.; Berneman, D.; Weitzman, O.; Alpaugh, A.; Pietraszkiewicz, A.; Metsuyanin, S.; Shinskaya, A.; Papa, M.Z.; Yarden, R.I. BRCA1 targets G2/M cell cycle proteins for ubiquitination and proteasomal degradation. *Oncogene* **2013**, *32*, 5005–5016.
105. Starita, L.M.; Machida, Y.; Sankaran, S.; Elias, J.E.; Griffin, K.; Schlegel, B.P.; Gygi, S.P.; Parvin, J.D. BRCA1-dependent ubiquitination of γ -tubulin regulates centrosome number. *Mol. Cell. Biol.* **2004**, *24*, 8457–8466.
106. Eakin, C.M.; Maccoss, M.J.; Finney, F.L.; Klevit, R.E. Estrogen receptor α is a putative substrate for the BRCA1 ubiquitin ligase. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 5794–5799.

107. Zhu, Q.; Pao, G.M.; Huynh, A.M.; Suh, H.; Tonnu, N.; Nederlof, P.M.; Gage, F.H.; Verma, I.M. BRCA1 tumor suppression occurs via heterochromatin-mediated silencing. *Nature* **2011**, *477*, 179–184.
108. Ohta, T.; Sato, K.; Wu, W. The BRCA1 ubiquitin ligase and homologous recombination repair. *FEBS Lett.* **2011**, *585*, 2836–2844.
109. Caestecker, K.W.; van de Walle, G.R. The role of BRCA1 in DNA double-strand repair: Past and present. *Exp. Cell Res.* **2013**, *319*, 575–587.
110. Rosen, E.M. BRCA1 in the DNA damage response and at telomeres. *Front. Genet.* **2013**, *4*, 85.
111. Ouchi, T.; Lee, S.W.; Ouchi, M.; Aaronson, S.A.; Horvath, C.M. Collaboration of signal transducer and activator of transcription 1 (STAT1) and BRCA1 in differential regulation of IFN- γ target genes. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 5208–5213.
112. Chen, W.; Sun, Z.; Wang, X.J.; Jiang, T.; Huang, Z.; Fang, D.; Zhang, D.D. Direct interaction between Nrf2 and p21^{Cip1/WAF1} upregulates the Nrf2-mediated antioxidant response. *Mol. Cell* **2009**, *34*, 663–673.
113. Mitshishi, Y.; Motohashi, H.; Yamamoto, M. The Keap1-Nrf2 system in cancers: Stress response and anabolic metabolism. *Front. Oncol.* **2012**, *2*, 200.
114. Bryan, H.K.; Olayanju, A.; Goldring, C.E.; Park, B.K. The Nrf2 cell defense pathway: Keap1-dependent and -independent mechanisms of regulation. *Biochem. Pharmacol.* **2013**, *85*, 705–717.
115. Ma, Q. Role of Nrf2 in oxidative stress and toxicity. *Annu. Rev. Pharmacol. Toxicol.* **2013**, *53*, 401–426.
116. Niture, S.K.; Khatri, R.; Jaiswal, A.K. Regulation of Nrf2—an update. *Free Radic. Biol. Med.* **2014**, *66*, 36–44.
117. Gorrini, C.; Baniasadi, P.S.; Harris, I.S.; Silvester, J.; Inoue, S.; Snow, B.; Joshi, P.A.; Wakeham, A.; Molyneux, S.D.; Martin, B.; *et al.* BRCA1 interacts with Nrf2 to regulate antioxidant signaling and cell survival. *J. Exp. Med.* **2013**, *210*, 1529–1544.
118. Li, W.; Kong, A.N. Molecular mechanisms of Nrf2-mediated antioxidant response. *Mol. Carcinog.* **2009**, *48*, 91–104.
119. Satoh, H.; Moriguchi, T.; Takai, J.; Ebina, M.; Yamamoto, M. Nrf2 prevents initiation but accelerates progression through the Kras signaling pathway during lung carcinogenesis. *Cancer Res.* **2013**, *73*, 4158–4168.
120. Mullan, P.B.; Quinn, J.E.; Harkin, D.P. The role of BRCA1 in transcriptional regulation and cell cycle control. *Oncogene* **2006**, *25*, 5854–5863.
121. Chapman, M.S.; Verma, I.M. Transcriptional activation by BRCA1. *Nature* **1996**, *382*, 678–679.
122. Monteiro, A.N.; August, A.; Hanafusa, H. Evidence for a transcriptional activation function of BRCA1 C-terminal region. *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 13595–13599.
123. Monteiro, A.N.; August, A.; Hanafusa, H. Common BRCA1 variants and transcriptional activation. *Am. J. Hum. Genet.* **1997**, *61*, 761–762.
124. Haile, D.T.; Parvin, J.D. Activation of transcription *in vitro* by the the BRCA1 carboxyl-terminal domain. *J. Biol. Chem.* **1999**, *274*, 2113–2117.

125. Hu, Y.F.; Hao, Z.L.; Li, R. Chromatin remodeling and activation of chromosomal DNA replication by an acidic transcriptional activation domain from BRCA1. *Genes Dev.* **1999**, *13*, 637–642.
126. Scully, R.; Anderson, S.F.; Chao, D.M.; Wei, W.; Ye, L.; Young, R.A.; Livingston, D.M.; Parvin, J.D. BRCA1 is a component of the RNA polymerase II holoenzyme. *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 5605–5610.
127. Schlegel, B.P.; Green, J.V.; Ladias, J.A.; Parvin, J.D. BRCA1 interaction with RNA polymerase II reveals a role for hRPB2 and hRPB10 α in activated transcription. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 3148–3153.
128. Anderson, S.F.; Schlegel, B.P.; Nakajima, T.; Wolpin, E.S.; Parvin, J.D. BRCA1 protein is linked to the RNA polymerase II holoenzyme complex via RNA helicase A. *Nat. Genet.* **1998**, *19*, 254–256.
129. Pao, G.M.; Janknecht, R.; Ruffner, H.; Hunter, T.; Verma, I.M. CBP/p300 interact with and function as transcriptional coactivators of BRCA1. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 1020–1025.
130. Moisan, A.; Larochelle, C.; Guillemette, B.; Gaudreau, L. BRCA1 can modulate RNA polymerase II carboxy-terminal domain phosphorylation levels. *Mol. Cell. Biol.* **2004**, *24*, 6947–6956.
131. Chen, G.C.; Guan, L.S.; Yu, J.H.; Li, G.C.; Choi Kim, H.R.; Wang, Z.Y. Rb-associated protein 46 (RbAp46) inhibits transcriptional transactivation mediated by BRCA1. *Biochem. Biophys. Res. Commun.* **2001**, *284*, 507–514.
132. Bochar, D.A.; Wang, L.; Beniya, H.; Kinev, A.; Xue, Y.; Lane, W.S.; Wang, W.; Kashanchi, F.; Shiekhhattar, R. BRCA1 is associated with a human SWI/SNF-related complex: Linking chromatin remodeling to breast cancer. *Cell* **2000**, *102*, 257–265.
133. Fagan, D.H.; Yee, D. Crosstalk between IGF1R and estrogen receptor signaling in breast cancer. *J. Mammary Gland Biol. Neoplasia* **2008**, *13*, 423–429.
134. Fan, S.; Wang, J.; Yuan, R.; Ma, Y.; Meng, Q.; Erdos, M.R.; Pestell, R.G.; Yuan, F.; Auborn, K.J.; Goldberg, I.D.; *et al.* BRCA1 inhibition of estrogen receptor signaling in transfected cells. *Science* **1999**, *284*, 1354–1356.
135. Fan, S.; Ma, Y.X.; Wang, C.; Yuan, R.Q.; Meng, Q.; Wang, J.A.; Erdos, M.; Goldberg, I.D.; Webb, P.; Kushner, P.J.; *et al.* Role of direct interaction in BRCA1 inhibition of estrogen receptor activity. *Oncogene* **2001**, *20*, 77–87.
136. Rosen, E.M.; Fan, S.; Ma, Y. BRCA1 regulation of transcription. *Cancer Lett.* **2006**, *236*, 175–185.
137. Zheng, L.; Annab, L.A.; Afshari, C.A.; Lee, W.H.; Boyer, T.G. BRCA1 mediates ligand-independent transcriptional repression of the estrogen receptor. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 9587–9592.
138. Fan, S.; Ma, Y.X.; Wang, C.; Yuan, R.Q.; Meng, Q.; Wang, J.A.; Erdos, M.; Goldberg, I.D.; Webb, P.; Kushner, P.J.; *et al.* p300 modulates the BRCA1 inhibition of estrogen receptor activity. *Cancer Res.* **2002**, *62*, 141–151.
139. Wang, C.; Fan, S.; Li, Z.; Fu, M.; Rao, M.; Ma, Y.; Lisanti, M.P.; Albanese, C.; Katzenellenbogen, B.S.; Kushner, P.J.; *et al.* Cyclin D1 antagonizes BRCA1 repression of estrogen receptor α activity. *Cancer Res.* **2005**, *65*, 6557–6567.
140. Xu, J.; Fan, S.; Rosen, E.M. Regulation of the estrogen-inducible gene expression profile by the breast cancer susceptibility gene BRCA1. *Endocrinology* **2005**, *146*, 2031–2047.

141. Wang, Q.; Zhang, H.; Kajino, K.; Greene, M.I. BRCA1 binds c-Myc and inhibits its transcriptional and transforming activity in cells. *Oncogene* **1998**, *17*, 1939–1948.
142. Li, H.; Lee, T.H.; Avraham, H. A novel tricomplex of BRCA1, Nmi, and c-Myc inhibits c-Myc-induced human telomerase reverse transcriptase gene (hTERT) promoter activity in breast cancer. *J. Biol. Chem.* **2002**, *277*, 20965–20973.
143. Kennedy, R.D.; Gorski, J.J.; Quinn, J.E.; Stewart, G.E.; James, C.R.; Moore, S.; Mulligan, K.; Emberley, E.D.; Lioe, T.F.; Morrison, P.J.; *et al.* BRCA1 and c-Myc associate to transcriptionally repress psoriasin, a DNA damage-inducible gene. *Cancer Res.* **2005**, *65*, 10265–10272.
144. Schaeper, U.; Subramanian, T.; Lim, L.; Boyd, J.M.; Chinnadurai, G. Interaction between a cellular protein that binds to the C-terminal region of adenovirus E1A (CtBP) and a novel cellular protein is disrupted by E1A through a conserved PLDLS motif. *J. Biol. Chem.* **1998**, *273*, 8549–8552.
145. Li, S.; Ting, N.S.; Zheng, L.; Chen, P.L.; Ziv, Y.; Shiloh, Y.; Lee, E.Y.; Lee, W.H. Functional link of BRCA1 and ataxia telangiectasia gene product in DNA damage response. *Nature* **2000**, *406*, 210–215.
146. Wu-Baer, F.; Baer, R. Effect of DNA damage on a BRCA1 complex. *Nature* **2001**, *414*, 36.
147. Zheng, L.; Pan, H.; Li, S.; Flesken-Nikitin, A.; Chen, P.L.; Boyer, T.G.; Lee, W.H. Sequence-specific transcriptional corepressor function for BRCA1 through a novel zinc finger protein, ZBRK1. *Mol. Cell* **2000**, *6*, 757–768.
148. Tan, W.; Zheng, L.; Lee, W.H.; Boyer, T.G. Functional dissection of transcription factor ZBRK1 reveals zinc fingers with dual roles in DNA-binding and BRCA1-dependent transcriptional repression. *J. Biol. Chem.* **2004**, *279*, 6576–6587.
149. Furuta, S.; Wang, J.M.; Wei, S.; Jeng, Y.M.; Jian, X.; Gu, B.; Chen, P.L.; Lee, E.Y.; Lee, W.H. Removal of BRCA1/CtIP/ZBRK1 repressor complex on ANG1 promoter leads to accelerated mammary tumor growth contributed by prominent vasculature. *Cancer Cell* **2006**, *10*, 13–24.
150. Ahmed, K.M.; Tsai, C.Y.; Lee, W.H. Derepression of HMGA2 via removal of ZBRK1/BRCA1/CtIP complex enhances mammary tumorigenesis. *J. Biol. Chem.* **2010**, *285*, 4464–4471.
151. Fan, W.; Jin, S.; Tong, T.; Zhao, H.; Fan, F.; Antinore, M.J.; Rajasekaran, B.; Wu, M.; Zhan, Q. BRCA1 regulates GADD45 through its interaction with the OCT-1 and CAAT motif. *J. Biol. Chem.* **2002**, *277*, 8061–8067.
152. Andrews, H.N.; Mullan, P.B.; McWilliams, S.; Sebelova, S.; Quinn, J.E.; Gilmore, P.M.; McCabe, N.; Pace, A.; Koller, B.; Johnston, P.G.; *et al.* BRCA1 regulates the interferon γ -mediated apoptotic response. *J. Biol. Chem.* **2002**, *277*, 26225–26232.
153. Ning, S.; Pagano, J.S.; Barber, G.N. IRF7: Activation, regulation, modification and function. *Genes Immun.* **2011**, *12*, 399–414.
154. Benezra, M.; Chevallier, N.; Morrison, D.J.; MacLachlan, T.K.; El-Deiry, W.S.; Licht, J.D. BRCA1 augments transcription by the NF- κ B transcription factor by binding to the Rel domain of the p65/RelA subunit. *J. Biol. Chem.* **2003**, *278*, 26333–26341.
155. Habraken, Y.; Joloi, O.; Piette, J. Differential involvement of the hMRE11/hRAD50/NBS1 complex, BRCA1 and MLH1 in NF- κ B activation by camptothecin and X-ray. *Oncogene* **2003**, *22*, 6090–6099.

156. Volcic, M.; Karl, S.; Baumann, B.; Salles, D.; Daniel, P.; Fulda, S.; Wiesmüller, L. NF- κ B regulates DNA double-strand break repair in conjunction with BRCA1-CtIP complexes. *Nucleic Acids Res.* **2012**, *40*, 181–195.
157. Kang, H.J.; Kim, H.J.; Rih, J.K.; Mattson, T.L.; Kim, K.W.; Cho, C.H.; Isaacs, J.S.; Bae, I. BRCA1 plays a role in the hypoxic response by regulating HIF-1 α stability and by modulating vascular endothelial growth factor expression. *J. Biol. Chem.* **2006**, *281*, 13047–13056.
158. Kawai, H.; Li, H.; Chun, P.; Avraham, S.; Avraham, H.K. Direct interaction between BRCA1 and the estrogen receptor regulates vascular endothelial growth factor (VEGF) transcription and secretion in breast cancer cells. *Oncogene* **2002**, *21*, 7730–7739.
159. Pollak, M. Insulin and insulin-like growth factor signaling in neoplasia. *Nat. Rev. Cancer* **2008**, *8*, 915–928.
160. Riedemann, J.; Macaulay, V.M. IGF1R signaling and its inhibition. *Endocr. Relat. Cancer* **2006**, *13*, S33–S43.
161. Law, J.H.; Habibi, G.; Hu, K.; Masoudi, H.; Wang, M.Y.; Stratford, A.L.; Park, E.; Gee, J.M.; Finlay, P.; Jones, H.E.; *et al.* Phosphorylated insulin-like growth factor-I/insulin receptor is present in all breast cancer subtypes and is related to poor survival. *Cancer Res.* **2008**, *68*, 10238–10246.
162. Maor, S.B.; Abramovitch, S.; Erdos, M.R.; Brody, L.C.; Werner, H. BRCA1 suppresses insulin-like growth factor-I receptor promoter activity: Potential interaction between BRCA1 and Sp1. *Mol. Genet. Metab.* **2000**, *69*, 130–136.
163. Abramovitch, S.; Glaser, T.; Ouchi, T.; Werner, H. BRCA1-Sp1 interactions in transcriptional regulation of the IGF-1R gene. *FEBS Lett.* **2003**, *541*, 149–154.
164. Kang, H.J.; Yi, Y.W.; Kim, H.J.; Hong, Y.B.; Seong, Y.S.; Bae, I. BRCA1 negatively regulates IGF-1 expression through an estrogen-responsive element-like site. *Cell Death Dis.* **2012**, *3*, e336.
165. Houvras, Y.; Benezra, M.; Zhang, H.; Manfredi, J.J.; Weber, B.L.; Licht, J.D. BRCA1 physically and functionally interact with ATF1. *J. Biol. Chem.* **2000**, *275*, 36230–36237.
166. Yan, J.; Zhu, J.; Zhong, H.; Lu, Q.; Huang, C.; Ye, Q. BRCA1 interacts with FHL2 and enhances FHL2 transactivation function. *FEBS Lett.* **2003**, *553*, 183–189.
167. Kang, H.J.; Kim, H.J.; Cho, C.H.; Hu, Y.; Li, R.; Bae, I. BRCA1 transcriptional activity is enhanced by interactions between its AD1 domain and AhR. *Cancer Chemother. Pharmacol.* **2008**, *62*, 965–975.
168. Yeh, S.; Hu, Y.C.; Rahman, M.; Lin, H.K.; Hsu, C.L.; Ting, H.J.; Kang, H.Y.; Chang, C. Increase of androgen-induced cell death and androgen receptor transactivation by BRCA1 in prostate cancer. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 11256–11261.
169. Kang, H.J.; Kim, H.J.; Kim, S.K.; Barouki, R.; Cho, C.H.; Khanna, K.K.; Rosen, E.M.; Bae, I. BRCA1 modulates xenobiotic stress-inducible gene expression by interacting with ARNT in human breast cancer cells. *J. Biol. Chem.* **2006**, *281*, 14654–14662.
170. Wang, H.; Shao, N.; Ding, Q.M.; Cui, J.; Reddy, E.S.; Rao, V.N. BRCA1 proteins are transported to the nucleus in the absence of serum and splice variants BRCA1a, BRCA1b are tyrosine phosphoproteins that associate with E2F, cyclins and cyclin dependent kinases. *Oncogene* **1997**, *15*, 143–157.

171. Hu, Y.F.; Li, R. JunB potentiates function of BRCA1 activation domain 1 (AD1) through a coiled-coil-mediated interaction. *Genes Dev.* **2002**, *16*, 1509–1517.
172. Sum, E.Y.; Peng, B.; Yu, X.; Chen, J.; Byrne, J.; Lindeman, G.J.; Visvader, J.E. The LIM domain protein LMO4 interacts with the cofactor CtIP and the tumor suppressor BRCA1 and inhibits BRCA1 activity. *J. Biol. Chem.* **2002**, *277*, 7849–7856.
173. Vidarsson, H.; Mikaelssdottir, E.K.; Rafnar, T.; Bertwistle, D.; Ashworth, A.; Eyfjord, J.E.; Valgeirsdottir, S. BRCA1 and BRCA2 bind Stat5a and suppress its transcriptional activity. *FEBS Lett.* **2002**, *532*, 247–252.
174. Cable, P.L.; Wilson, C.A.; Calzone, F.J.; Rauscher, F.J., 3rd.; Scully, R.; Livingston, D.M.; Li, L.; Blackwell, C.B.; Futreal, P.A.; Afshari, C.A. Novel consensus DNA-binding sequence for BRCA1 protein complexes. *Mol. Carcinog.* **2003**, *38*, 85–96.
175. Kang, H.J.; Hong, Y.B.; Kim, H.J.; Rodriguez, O.C.; Nath, R.G.; Tilli, E.M.; Albanese, C.; Chung, F.L.; Kwon, S.H.; Bae, I. Detoxification: A novel function of BRCA1 in tumor suppression? *Toxicol. Sci.* **2011**, *122*, 26–37.
176. Miao, W.; Hu, L.; Scrivens, P.J.; Batist, G. Transcriptional regulation of NF-E2 p45-related factor (NRF2) expression by the aryl hydrocarbon receptor-xenobiotic response element signaling pathway: Direct cross-talk between phase I and II drug-metabolizing enzymes. *J. Biol. Chem.* **2005**, *280*, 20340–20348.
177. Hankinson, O. The aryl hydrocarbon receptor complex. *Annu. Rev. Pharmacol. Toxicol.* **1995**, *35*, 307–340.
178. Bae, I.; Rih, J.K.; Kim, H.J.; Kang, H.J.; Haddad, B.; Kirilyuk, A.; Fan, S.; Avantaggiati, M.L.; Rosen, E.M. BRCA1 regulates gene expression for orderly mitotic progress. *Cell Cycle* **2005**, *4*, 1641–1666.
179. Bae, I.; Fan, S.; Meng, Q.; Rih, J.K.; Kim, H.J.; Kang, H.J.; Xu, J.; Goldberg, I.D.; Jaiswal, A.K.; Rosen, E.M. BRCA1 induces antioxidant gene expression and resistance to oxidative stress. *Cancer Res.* **2004**, *64*, 7893–7909.
180. Nguyen, T.; Sherratt, P.J.; Pickett, C.B. Regulatory mechanisms controlling gene expression mediated by the antioxidant response element. *Annu. Rev. Pharmacol. Toxicol.* **2003**, *43*, 233–260.
181. Wang, Q.; Li, J.; Yang, X.; Sun, H.; Gao, S.; Zhu, H.; Wu, J.; Jin, W. Nrf2 is associated with the regulation of basal transcription activity of the *BRCA1* gene. *Acta Biochim. Biophys. Sin.* **2013**, *45*, 179–187.
182. Yoshida, K.; Miki, Y. Role of BRCA1 and BRCA1 as regulators of DNA repair, transcription, and cell cycle in response to DNA damage. *Cancer Sci.* **2004**, *95*, 866–871.
183. Narod, S.A.; Foulkes, W.D. BRCA1 and BRCA2: 1994 and beyond. *Nat. Rev. Cancer* **2004**, *4*, 665–676.
184. Ouchi, T. BRCA1 phosphorylation: Biological consequence. *Cancer Biol. Ther.* **2006**, *5*, 470–475.
185. Guo, Z.; Kozlov, S.; Lavin, M.F.; Person, M.D.; Paull, T.T. ATM activation by oxidative stress. *Science* **2010**, *330*, 517–521.
186. Guo, Z.; Deshpande, R.; Paull, T.T. ATM activation in the presence of oxidative stress. *Cell Cycle* **2010**, *9*, 4805–4811.
187. Willis, J.; Patel, Y.; Lentz, B.L.; Yan, S. APE2 is required for ATR-Chk1 checkpoint activation in response to oxidative stress. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 10592–10597.

188. Zhang, J.; Kim, J.; Alexander, A.; Cai, S.; Tripathi, D.N.; Dere, R.; Tee, A.R.; Tait-Mulder, J.; Nardo, A.D.; Han, J.M.; Kwiatkowski, E.; *et al.* A tuberous sclerosis complex signalling node at the peroxisome regulates mTORC1 and autophagy in response to ROS. *Nat. Cell Biol.* **2013**, *15*, 1186–1196.
189. Chen, B.P.C.; Li, M.; Asaithamby, A. New insights into the roles of ATM and DNA-PKcs in the cellular response to oxidative stress. *Cancer Lett.* **2012**, *327*, 103–110.
190. Jorgenson, T.C.; Zhong, W.; Oberley, T.D. Redox imbalance and biochemical changes in cancer. *Cancer Res.* **2013**, *73*, 6118–6123.
191. Miki, H.; Funato, Y. Regulation of intracellular signaling through cysteine oxidation by reactive oxygen species. *J. Biochem.* **2012**, *151*, 255–261.
192. Paulsen, C.E.; Carroll, K. Cysteine-mediated redox signaling: Chemistry, biology, and tools for discovery. *Chem. Rev.* **2013**, *113*, 4633–4679.
193. Watson, J. Oxidants, antioxidants and the current incurability of metastatic cancers. *Open Biol.* **2013**, doi:10.1098/rsob.120144.
194. Lau, A.T.Y.; Wang, Y.; Chiu, J.F. Reactive oxygen species: Current knowledge and applications in cancer research and therapeutic. *J. Cell. Biochem.* **2008**, *104*, 657–667.
195. Chibber, S.; Farhan, M.; Hassan, I.; Naseem, I. Novel aspect of chemophototherapy in treatment of cancer. *Tumor Biol.* **2012**, *33*, 701–706.
196. Nogueira, V.; Hay, N. Molecular pathway: Reactive oxygen species homeostasis in cancer cells and implications for cancer therapy. *Clin. Cancer Res.* **2013**, *19*, 4309–4314.
197. Raj, L.; Ide, T.; Gurkar, A.U.; Foley, M.; Schenone, M.; Li, X.; Tolliday, N.J.; Golub, T.R.; Carr, S.A.; Shamji, A.F.; *et al.* Selective killing of cancer cells by small molecule targeting the stress responses to ROS. *Nature* **2011**, *475*, 231–234.
198. Tassone, P.; Tagliaferri, P.; Perricelli, A.; Blotta, S.; Quaresima, B.; Martelli, M.L.; Goel, A.; Barbieri, V.; Costanzo, F.; Boland, C.R.; *et al.* BRCA1 expression modulates chemosensitivity of BRCA1-defective HCC1937 human breast cancer cells. *Br. J. Cancer* **2003**, *88*, 1285–1291.
199. Santarosa, M.; del Col, L.; Tonin, E.; Caragnano, A.; Viel, A.; Maestro, R. Premature senescence is a major response to DNA cross-linking agents in BRCA1-defective cells: Implication for tailored treatments of BRCA1 mutation carriers. *Mol. Cancer Ther.* **2009**, *8*, 844–854.
200. Fasano, J.; Muggia, F. Breast cancer arising in a BRCA-mutated background: Therapeutic implications from an animal model and drug development. *Ann. Oncol.* **2009**, *20*, 609–614.
201. Johnson, N.; Johnson, S.F.; Yao, W.; Li, Y.C.; Choi, Y.E.; Bernhardt, A.J.; Wang, Y.; Capelletti, M.; Sarosiek, K.A.; Moreau, L.A.; *et al.* Stabilization of mutant BRCA1 protein confers PARP inhibitor and platinum resistance. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 17041–17046.
202. Gorrini, C.; Harris, I.S.; Mak, T.W. Modulation of oxidative stress as an anticancer strategy. *Nat. Rev. Drug Discov.* **2013**, *12*, 931–947.
203. Parri, M.; Chiarugi, P. Redox molecular machines involved in tumor progression. *Antioxid. Redox Signal.* **2013**, *19*, 1828–1845.