

Aerobic Oxidation of Benzyl Alcohol on a Strontium-Based Gold Material: Remarkable Intrinsic Basicity and Reusable Catalyst

Karla Patrícia R. Castro ¹, Marco Aurélio S. Garcia ¹, Wiury C. de Abreu ^{1,2}, Samuel Anderson A. de Sousa ¹, Carla Verônica R. de Moura ¹, Jean Cláudio S. Costa ¹ and Edmilson M. de Moura ^{2,*}

¹ Chemistry Department, Federal University of Piauí, Teresina 64049-550, PI, Brazil; karlaribeirocastro@gmail.com (K.P.R.C.); marcoasuller@gmail.com (M.A.S.G.); wiury.abreu@ifma.edu.br (W.C.d.A.); samuelsousa@ufpi.edu.br (S.A.A.d.S.); carla@ufpi.edu.br (C.V.R.d.M.); jean-cla@iq.usp.br (J.C.S.C.)

² Federal Institute of Maranhão, Buriticupu 65393-000, MA, Brazil

* Correspondence: mmoura@ufpi.edu.br; Tel.: +55-869-9432-4080




Fig. S1. XPS pattern of the Au/Sr_{sup} catalyst considering the presence of O, Sr, Au, Cl and Na for the at.% calculation.




Fig. S2. XPS pattern of the Au/Sr_{sup} catalyst disregarding the presence of Cl and Na for the at.% calculation.