



Article

Comparison between Chemical Fertilization and Integrated Nutrient Management: Yield, Quality, N, and P Contents in *Dendranthema grandiflorum* (Ramat.) Kitam. Cultivars

Beniamino Leoni, Danilo Loconsole , Giuseppe Cristiano  and Barbara De Lucia 

Department of Agro-Environmental and Territorial Science (DISAAT), University of Bari, "Aldo Moro", Italy Via Amendola 165/A, 70120 Bari, Italy; beniamino.leoni@uniba.it (B.L.); giuseppe.cristiano@uniba.it (G.C.); barbara.delucia@uniba.it (B.D.L.)

* Correspondence: danilo.loconsole@hotmail.it; Tel.: +39-080-544-3039

Received: 27 February 2019; Accepted: 17 April 2019; Published: 19 April 2019



Abstract: To assess the effects of a new integrated nutrient management protocol on yield and cut stem quality, root morphology, N accumulation, nitrogen utilization efficiency (NUE), and P content in tissue, a biennial (2011 and 2012) chrysanthemum cut flower cultivation was carried out. In both years, two nutrition management (CNM: conventional NM and INM: integrated NM) treatments and two *Dendranthema grandiflorum* (Ramat.) Kitamura cultivar ("White CV₁" and "Yellow CV₂") treatments were compared. The treatments were arranged in a split-plot design with three replicates. CNM was fertilized using a recommended dose fertilization of mineral NPK; INM treatment was fertilized using a half dose (50%) of CNM plus a combined usage of N organic fertilizer, seaweed extract (*Ascophyllum nodosum*), and microorganism consortium (*Glomus* sp. and *Bacillus* sp.). Yield at harvest (+19%), number of leaves (+33%), leaf area (+46%), number of flower heads (+27%), and total aboveground dry weight (+40%) were significantly increased by the INM application compared to the control. In terms of the root system, the increase was evident in terms of length (+174%), volume (+167%), projected area (+166%), and surface area (+165%), tips (+175%), forks (+285%), and crossings (+464%). The greatest N accumulation, in both years, was registered by INM treatment at harvest: +94% in 2011 and +55% in 2012. Differences in the NM were evident in the NUE, which was highest in CNM (on average 162) compared to INM (on average 142). In both years the P content in above-ground chrysanthemum tissues was in the order of head > leaves > stems, which was maintained in both INM and CNM treatments. A higher yield (138 stems m⁻²) was obtained in "CV₂ Yellow" compared to "CV₁ White" (120 stems m⁻²). Based on our findings, applying INM to chrysanthemum improves yield, cut flower quality, and plant nutrient uptake, in an agro-environmentally sustainable way. A basic economic analysis on fertilizers, cost gross production, and takings difference obtained, was carried out.

Keywords: N organic fertilizer; seaweed extract; mycorrhizal inoculants; phosphate-solubilizing microorganisms; biofertilizers; microorganism consortium

1. Introduction

Fertilization is essential for optimizing crop productivity [1]. Mineral fertilizers, particularly nitrogen (N) and phosphorus (P), are important for plant nutrition [2,3]. However, when used in overly large doses they are also a potential source of environmental pollution [4–6]. Nutrient overapplication has introduced major challenges in terms of soil infertility [7], N and P runoff [8,9], environmental degradation [10], and climate change [11,12].

Today there is an increasing need for a balanced fertilization strategy, minimizing the use of mineral fertilizers to enhance both crop production and quality and nutrient uptake under low input conditions [13]. Mineral fertilizers can be replaced by organic fertilizers [14], plant biostimulants [15], and beneficial microbial inoculants [16].

Possible interventions in conservation agriculture include the combined use of inorganic and organic fertilizers, as well as biostimulants and biofertilizers in order to increase a balanced nutrient supply [17]. Integrated nutrition management (INM) focusing on the optimization of the biological potential improves fertilizer input efficiency, reduces environmental risks, and increases crop productivity, through root/rhizosphere management [18].

In terms of biostimulants, seaweed extracts are used in sustainable agriculture in order to increase growth, quality, and shelf life [19–21]. Many studies have demonstrated the positive effects of seaweed extracts on a wide range of crops, including cereals [22], ornamental and flowering plants [23], vegetables [24], and field crops [25].

Biofertilizers are also an important alternative source of plant nutrients and are key components of integrated nutrient management in crop production. The use of microbial inoculants with P solubilizing activities in soils is an environmental-friendly alternative to further applications of chemical-based P fertilizers [26,27]. Various studies have examined the potential of different bacterial species to solubilize inorganic phosphate compounds. *Bacillus* spp., and in particular *B. subtilis* and *B. megaterium*, may provide the available forms of P to plants, thus considerably improving plant growth performance [28–31].

Other microbial inoculants, such as arbuscular mycorrhiza fungi (AMF), increase the P availability through the expansion of the root surface area by extraradical hyphae formation [32,33].

The various benefits of AMF include increased growth and nutrient uptake (especially N, P, and K) and crop yields [34–38]. The AMF also produce a heat-stable protein called glomalin, which is a glycoprotein that enhances soil aggregation and helps in soil carbon sequestration. Together, glomalin and mycorrhizal hyphae lead to a stable soil structure.

The combined use of N organic fertilizers, biostimulants, and biofertilizers is therefore a new approach that has not been widely investigated in ornamentals, which entails developing many efficient formulations with low mineral inputs, with positive impacts on crops and environment.

Chrysanthemum (*Dendranthema grandiflorum* (Ramat.) Kitamura) is a commercial cut flower, belonging to the *Asteraceae* family, with nearly 200 cultivars. It is one of the top ten elite cut flowers globally, due to its different shapes, dazzling colors, varying sizes, and excellent vase life. In Italy, where our research was carried out, there is a considerable demand in both domestic and export markets.

Extracts of the plants (stems and flowers) have many potential medicinal properties, including anti-HIV, antibacterial, and antimycotic [39]. N, P, and K play a vital role in the production of good quality flowers. N is essential for the creation of biomass as well as for the biosynthesis of enzymes in chrysanthemum leaves [40].

The N requirements of chrysanthemums are known to be higher during the first seven weeks of growth, and during this time, deficiencies are more difficult to correct than in later stages of development [41]. Chrysanthemums take up N at an even rate from the time of planting until the flower bud differentiation stage where after N uptake decreases [42]. In chrysanthemums, the need for P is significantly lower than that of nitrogen [43]. K requirements are high, and its presence in the plant favorably affects growth and flower color [44].

To the best of our knowledge, there are no available data on how the INM system based on mineral and organic N fertilizers, seaweed extracts, plus a consortium between AMF (arbuscular mycorrhiza fungi) and PSB (Phosphate solubilizing bacteria), affects yield and quality in chrysanthemums.

The goal of this research was to evaluate the effects of an innovative INM compared to conventional nutrient management, in chrysanthemum cut flower cultivation, on: (1) yield and cut stem quality, (2) N concentration, accumulation, and utilization efficiency and P uptake, (3) root architecture, and (4) soil fertility.

2. Materials and Methods

2.1. Experimental Conditions

Two field experiments were carried out in 2011 and 2012, from August to December, at a floricultural farm located in Sannicandro di Bari (southern Italy: 40°59'24" N, 16°47'01"E, 181 m a.s.l.). The local climatic conditions are characterized by hot dry summers and mild rainy autumns and winters, typical of the Mediterranean climate. During the plant growth period under natural photoperiod, the mean air temperature was 17.2 °C and 18.2 °C in 2011 and 2012, respectively; minimum air temperature was 3.7 °C in December 2011 and 5.4 °C in December 2012; maximum air temperature was 32.4 °C in August 2011, and 32.6 °C in August 2012.

Seasonal chrysanthemum cuttings (Minstreel Serie, Straathof Plants BV, The Netherlands), ideal for blooming from November to late December, were obtained from a local commercial propagator, with the following characteristics: stem length, 11.6 cm; number of leaves, 8; leaf area, 81.1 cm²; plant fresh weight, 3.1 g; and plant dry weight, 0.3 g. In both years, plants were transplanted on 6 August into an uncovered tunnel. In the first week of October in both years, the tunnel was covered by ethylene vinyl acetate (EVA) film.

The main soil characteristics (taken from 0 to 25 cm depth) are described in Table 1. Soil pH was determined with a pH meter (P9991, Hanna Instruments, Italy) in a settling suspension on a 60 g sample mixed with 150 mL of deionized water, after shaking for 60 min at room temperature (22 °C). The soil used for our experiment was slightly sub-alkaline (pH = 7.34, near to neutrality) and it was representative of Apulian soils in which chrysanthemum was cultivated with remarkable production results. Chrysanthemum plants generally grow with a pH ranging between 6 and 7.2 [45].

The electric conductivity (EC) was measured on water extract (1:5 v/v) with a conductivity meter (HI 4321, Hanna Instruments, Italy). Soil organic carbon (SOC) was determined by wet oxidation. Based on USDA classification, experimental soil was classified as clay loam soil. Experimental soil was moderately provided with organic matter and CEC was also classified as moderate [46].

The total Kjeldahl N (TKN) was measured using 1 g samples of both growing media and plant tissues using the Kjeldahl method after 96% H₂SO₄ hot digestion. Total phosphorus was determined (P) by the colorimetric molybdovanadate phosphoric acid method. Exchangeable K, Ca, and Mg were determined using 0.2 g of dry sample (105 °C for 24 h) after acid digestion in a microwave oven (CEM Mars Xpress, Cologno al Serio, IT). Substrate digests were filtered, diluted, and analyzed by atomic absorption spectrometry (Perkin-Elmer Analyst 200, Waltham, MA, USA). The analyses were carried out in triplicate.

The soil was sandy clay with a slightly alkaline pH of 7.3 (IUSS), EC of 1.77 dS m⁻¹, and moderately high CEC (cation exchange capacity) of 23.8 Meq 100 g⁻¹.

Table 1. Initial soil physico-chemical characteristics (mean ± standard error). Data are the means of three samples.

Parameter	Value
pH (soil:H ₂ O ratio 1:2.5)	7.34 ± 0.2
Electric Conductivity (EC) (soil:H ₂ O ratio 1:5) (dS m ⁻¹)	1.77 ± 0.08
Cation exchange capacity (CEC) (Meq100g ⁻¹)	23.8
Sand (%)	52 ± 3
Silt (%)	16 ± 2
Clay (%)	32 ± 4
Total C (g kg ⁻¹)	12.54 ± 1.2
Organic matter (g kg ⁻¹)	21.61 ± 1.9
T Kjeldahl-N (g kg ⁻¹)	1.15 ± 0.13
P (mg kg ⁻¹)	71.25 ± 0.9
Available K (mg kg ⁻¹)	579 ± 10.1
Available Ca (mg kg ⁻¹)	2160 ± 22
Available Mg (mg kg ⁻¹)	495 ± 31

2.2. Treatments and Experimental Design

In both years, four treatments in total consisting of two nutrition management (NM) and two *Dendranthema grandiflorum* (Ramat.) Kitamura cultivar (CV) were compared as follows:

1. Conventional NM (CNM or control) and integrated NM (INM);
2. “White CV₁” and “Yellow CV₂”.

Treatments were carried out using a split-plot design with three replicates, with NM as the main plot and CV as the subplot. The surface of each experimental plot measured 2.2 m².

CNM treatment was applied through a fertigation system using a recommended dose of mineral NPK: 17 g m⁻² N, 16 g m⁻² P₂O₅, and 17 g m⁻² K₂O plus microelements, starting one week after transplanting, every week, for 12 weeks, the last one during the second week of November (flower bud differentiation).

INM treatment was applied by fertigation at a half dose (50%) of CNM plus a mixture of an N organic fertilizer, seaweed extract and microorganism consortium as shown in Table 2, starting from transplantation. Commercial products were applied at the manufacturer’s recommended rates.

NPK doses added with INM fertilization were the following: 11.8 g m⁻² N, 8 g m⁻² P₂O₅, and 12 g m⁻² K₂O. N organic fertilizer added to the mineral NPK dose mentioned above, was derived by hydrolyzed animal epithelium, beet molasses extract, and brown seaweed extract.

In the second year, the same treatments were repeated.

In both years, the growing density was 34 plants m⁻².

Table 2. Combined use of N organic fertilizer, seaweed extract, and microorganism consortium applied in two experiments (2011 and 2012).

Type and Commercial Product (*)	Content	Total Rate (g/100m ²)	Weeks of Applications (n) (**)
N organic fertilizer (Euroflorid)	N = 5% w/w (N = 0.5 g/m ²)	75	I, II, III, IV, V
N organic fertilizer (Ammiostim-bio)	N = 6% w/w (N = 0.9 g/m ²)	25	VI, VII, VIII
Seaweed extract (Euroalg)	<i>Ascophyllum nodosum</i> (L.) Le Jol. 32% w/w, N = 1.5% w/w (N = 0.9 g/m ²), K ₂ O = 5.0% w/w	58	I, II, III, IV, V, VI, VII, VIII, IX, X, XI, XII
Microorganism consortium (Micotric L)	<i>Glomus mosseae</i> and <i>G. intraradices</i> (2 spores g ⁻¹) and <i>Bacillus megaterium</i> var. <i>Phosphaticum</i> 6 × 10 ⁷ CFUs g ⁻¹	90	I, II, III, IV
Microorganism consortium (Europlus)	<i>Glomus mosseae</i> (2 spores g ⁻¹) + <i>Trichoderma viride</i> and <i>Bacillus megaterium</i> var. <i>Phosphaticum</i> 6 × 10 ⁷ CFUs g ⁻¹	162	I, II, III, IV

(*) by Eurovix SpA, Entratico (BG), Italy; (**) fertigation from transplant (week 1) to flower bud differentiation (week 12).

During the experiments, all field management procedures (e.g., irrigation and pest control) were the same among treatments. The irrigation system was a micro drip; each drip line was placed between two plants rows with an emitter (pressure compensating) discharge rate of 2.0 L h⁻¹. Except for nutrition, production was carried out using the grower’s standard practices. Cut flowers were harvested when 50% of flower heads had opened.

Morpho-biometric measurements were carried out at the Department of Agro-environmental and Territorial Sciences (DISAAT), University of Bari, Italy. Plants were sampled for aboveground and ground biomass and N and P content (%) at 55, 93, and 131 DAT (days after transplant) in both growing periods.

The growth and yield observations were recorded on twelve randomly selected plants from each treatment.

In both years, at harvest (second ten days of December), the soil was washed from roots, and aboveground plants were divided into stems, leaves, and flowers, which were oven dried at 70 °C until they reached a constant mass to measure the respective dry weights.

At flower harvest, the measurements involved: yield (secondary branches = stems m⁻²), stem length (cm), inflorescence (n and diameter, cm), leaves (n), and leaf area (cm²), Chlorophyll SPAD

(Single-photon avalanche diode) index (Minolta Chlorophyll Meter SPAD-502), dry and fresh weight of leaves, stems, inflorescences, roots, and whole plants. In order to perform root morphology analysis, only in the first year, roots were spread out, washed, and then scanned at 300 dpi on an HP DeskScan II scanner (HEWLETT PACKARD C6261A, Palo Alto, CA, USA). Root analysis was performed using the WinRHIZO® image analysis system (V 4.1c Régent Instruments, Quebec, Canada); measurements involved total root length, average root diameter, projected and surface area, tips, forks, and crossings.

The total Kjeldahl N (TKN) content was measured, both in the first and second years, using 1 g samples of foliar and radical tissues, using the Kjeldahl method after 96% H₂SO₄ hot digestion. On the other hand, the P-Olsen measurement was only used during the first year.

Nitrogen utilization efficiency (NUE) was estimated by the ratio of dry biomass to plant N accumulation at harvest.

2.3. Economic Analysis

A basic economic analysis about fertilizer costs (for CNM and INM), gross sealable production, and profit raised was developed.

2.4. Statistical Analysis

The data were analyzed by three-way ANOVA using CoStat-Statistics Software. Treatment means were separated with Student–Newman–Keuls (SNK) ($p \leq 0.05$).

3. Results

The overall aims of this research were to evaluate the effects of an innovative INM compared to CNM, in a biennial chrysanthemum cut flower cultivation, on (i) yield and cut stem quality; (ii) root morphology; and (iii) N accumulation, NUE, and P content in plant tissue.

The main effect of NM was found to be highly significant for most of the parameters investigated.

Yield at harvest, as determined by the harvestable number of cut stems per plant (Table 3), increased significantly in INM (140 stems m⁻², +19%) compared to those under CNM (118 stems m⁻²).

Genotype influenced marketable yield: CV₂ registered the highest value (138 stems m⁻²), surpassing that of CV₁ by 15% (on average 120 stems m⁻²).

Concerning the Y factor, the yields were not different (133 stems m⁻² on average).

Table 3. Main effects of nutrient management, cultivar on yield, stem height, leaf number, leaf area, chlorophyll index, and number of flower heads in chrysanthemum plants over the two years of application.

Treatments	Yield (no. stems m ⁻²)	Stem Height (cm)	Leaves (no./plant)	Leaf Area (cm ⁻²)	Chl. Index (SPAD)	Flower Heads (no. stems ⁻¹)
Nutrition Management (NM)						
Conventional NM (CNM)	118a	103	60b	2064b	44.8	6.6b
Integrated NM (INM)	140b	106	80a	3017a	46.9	8.4a
Significance	*	ns	*	**	ns	**
Cultivar (CV)						
CV 1	120b	105	64	2416	45.1	8.2
CV 2	138a	104	76	2665	46.6	6.9
Significance	*	ns	ns	ns	ns	ns
Year						
2011	133	116a	82	2795	46.6	8.1
2012	133	92b	59	2486	45.1	7.3
Significance	ns	*	*	ns	ns	ns
Interaction						
NM × CV	*	ns	ns	ns	ns	*
NM × Year	ns	ns	ns	ns	ns	ns
CV × Year	ns	ns	ns	ns	ns	ns
NM × CV × year	ns	ns	ns	ns	ns	ns

Different letters within each column indicate significant differences according to SNK test ($p \leq 0.05$). NS not significant * $p < 0.05$ and ** 0.01, indicate level of significance.

Table 3 also shows the influence of the treatments on the commercial quality parameters of the cut stems at harvest. The stem height is an important parameter that is used for the classification of the stems for marketing and sales, and in fact, customers often prefer flowers with a longer stem. Stem height was not found to be significant between both NM and CV treatments; however, it showed significant differences among Y: in 2012 it was 20% lower (92 cm) than 2011 (116 cm).

Regarding the number of leaves per plant, the INM treatment led to an increase of 33% (80 leaves/stem) compared to CNM (60 leaves/stem); in 2012 the number of leaves (59) showed an average decrease of 28% (82 leaves) compared to 2011.

The INM treatment also showed a significant increase of 46% (3017 cm²) in the leaf area value compared to CNM (2064 cm²).

The chlorophyll index SPAD was not significant in any of the treatments.

The number of flower heads per stem was highest (8.4) with an increase of +27% when plants were treated with INM, compared to CNM (6.6). No differences were found between the cultivars and years.

Concerning the leaves, stems, flower heads, and aboveground dry weight, Table 4 shows the statistically significant differences in favor of INM compared to CNM. Leaf values showed a 38% increase, stem value a 37% increase, and flower heads a 55% increase, which were reflected in the increase of aboveground dry weight (+40%). No difference was found between the cultivars.

Table 4 also shows that 2011 had the highest aboveground dry weight value, which decreased to 25% during 2012.

Table 4. Main effects of nutrient management, cultivar on dry weight of various organs, and above-plant on chrysanthemum over the two years of application at harvest time.

Treatments	Dry Weight (g)			
	Leaves	Stem	Heads	Above-plant
Nutrition Management (NM)				
CNM	8.30 b	14.61 b	5.65 b	28.56 b
INM	11.10 a	20.05 a	8.75 a	39.90 a
Significance	**	**	**	**
Cultivar (CV)				
CV 1	9.40	17.10	8.05	34.55
CV 2	10.00	16.60	7.95	34.55
Significance	ns	ns	*	ns
Year				
2011	10.80	19.80	8.70 a	39.30 a
2012	8.60	14.40	6.50 b	29.5 b
Significance	*	**	*	**
Interaction				
NM × CV	ns	ns	ns	ns
NM × Year	*	ns	*	*
CV × Year	ns	ns	ns	ns
NM × CV × year	ns	ns	ns	ns

Different letters within each column indicate significant differences according to SNK test ($p \leq 0.05$). NS not significant * $p < 0.05$ and ** 0.01, indicate level of significance.

In 2011 the root morphology (Table 5) was evaluated. Parameter values for the plants under INM treatment were higher than CNM as follows: root length (+174%), area projection (+166%), surface area (165%), root volume (+167%), tips (+175%), forks (+285%), and crossings (+464%).

Regarding the CV, the best performing root system was White (CV₁) compared to Yellow (CV₂): root length (+63%), area projection (+37%), surface area (+38%), root volume (+19%), tips (+100%), forks (+109%), and crossings (+197%).

Table 5. Main effects of nutrient protocol management and cultivar on total root length (TRL), area projection (AP), surface area (SA), root volume (RV), root tips (RT), root forks (RF), and root crossings (RC) at 2011 harvest period in chrysanthemum plants.

Treatments	TRL (cm)	AP (cm ²)	SA (cm ²)	RV (cm ³)	RT (no.)	RF (no.)	RC (no.)
Nutrition Management (NM)							
CNM	382.1 b	21.2 b	66.7 b	0.9 b	1264.8 b	948.4 b	34.7 b
INM	1049.2 a	56.3 a	177.0 a	2.4 a	3486.6 a	3655.4 a	195.6 a
Significance	**	**	**	**	**	**	*
Cultivar (CV)							
CV 1	975.0 a	48.7 a	153.0 a	1.9 a	3535.2 a	3501.8 a	208.2 a
CV 2	597.4 b	35.5 b	110.8 b	1.6 b	1768.3 b	1676.0 b	70.0 b
Significance	*	*	*	*	**	**	**
Interaction							
NM × CV	*	*	*	*	**	**	**

Different letters within each column indicate significant differences according to SNK test ($p \leq 0.05$). NS not significant * $p < 0.05$ and **0.01, indicate level of significance.

Concerning the plant N accumulation (g m⁻²) at every DAT in both years (Table 6), the maximum value was obtained under INM, which was the result of a simultaneous increase in dry weight (Table 4). The highest N accumulation, in both years, was at harvest (131 DAT), in 2011 with an increase of 94%, and in 2012 with an increase of 55%. No significant difference was found between the CVs, except for the flower head value at 131 DAT in both years.

Table 6. Main effects of nutrient management and cultivar on N accumulation (g m⁻²) at three different days after transplant (DAT) in chrysanthemum plants over the two years of application.

Treatments	DAT		
	55	93	131
First Year			
Nutrition Management (NM)			
CNM	4.33 b	5.47 b	6.20 b
INM	6.28 a	9.13 a	10.39 a
Significance	*	*	*
Cultivar (CV)			
CV 1	5.55	7.20	8.99
CV 2	5.61	7.68	8.48
Significance	ns	ns	ns
Interaction			
NM × CV	ns	ns	ns
Second Year			
Nutrition Management (NM)			
CNM	2.67	5.05	6.37 b
INM	3.31	6.12	9.90 a
Significance	*	*	**
Cultivar (CV)			
CV 1	2.93	5.42	8.47 a
CV 2	3.03	5.72	7.57 b
Significance	NS	NS	*
Interaction			
NM × CV	ns	ns	ns

Different letters within each column indicate significant differences according to the SNK test ($p \leq 0.05$). NS not significant * $p < 0.05$ and **0.01, indicate level of significance.

Table 7 shows that in both years CNM treatment statistically influenced N accumulation (gm^{-2}) in all plant epigeal organs and on all sample dates. In the first year, the highest N accumulation was observed compared to CNM in the leaves (+48%) at 55 DAT, stems at 93 DAT (+85%), and flower buds (+79%) at 131 DAT. Regarding INM, in the second year, the highest value was recorded in leaves (+28%) at 55 DAT, stems (+46%), and flower buds (+117%) at 131 DAT. In both years the CVs did not influence N accumulation.

Table 7. Main effects of nutrient management and cultivar on N accumulation (g m^{-2}) in different organs at three different DAT in chrysanthemum plants over the two years of application.

Treatments	DAT						
	55		93		131		
	Leaves	Stems	Leaves	Stems	Leaves	Heads	Stems
First Year							
Nutrition Management (NM)							
CNM	3.05	1.28	3.31	2.16	2.83	1.87	1.50
INM	4.52	1.76	5.14	3.99	4.50	3.34	2.55
Significance	**	*	**	**	**	**	**
Cultivar (CV)							
CV 1	4.09	1.46	4.08	3.13	3.52	3.48	2.00
CV 2	4.04	1.57	4.53	3.26	3.75	2.73	2.00
Significance	ns	ns	ns	ns	ns	*	ns
Interaction NM \times CV							
Second Year							
Nutrition Management (NM)							
CNM	1.84	0.83	3.07	1.88	2.75	1.28	1.75
INM	2.35	0.96	3.57	2.55	3.30	2.78	2.57
Significance	*	*	*	*	*	**	*
Cultivar (CV)							
CV 1	2.07	0.87	3.23	2.19	3.09	1.71	2.54
CV 2	2.11	0.91	3.52	2.20	2.95	1.97	1.77
Significance	ns	ns	ns	ns	ns	ns	*
Interaction NM \times CV							

Different letters within each column indicate significant differences according to SNK test ($p \leq 0.05$). NS not significant * $p < 0.05$ and **0.01, indicate level of significance.

Figure 1 shows that in both years the N utilization efficiency (NUE) value was highest in CNM (on average 162) compared to INM (on average 142); no significant difference was found between the CVs.

In both years, the P content (%), at harvest, in above-ground vegetative tissues (leaves, stems, and heads) of INM plants was higher than those of CNM plants (Table 8). In the first year, the increase in INM compared to CNM was 11% in the leaves, 20% in the stems, and 21% in the flower heads. In the second year, the increase in P content in the leaves under INM was similar to that recorded in the first year (12%), while it was lower for stems (+12%) and flower heads (+14%). In both years the P content in above-ground vegetative tissues were in the order of head > leaves > stems, which was maintained in both INM and CNM treatments.




Figure 1. Main effects of nutrient management and cultivar on N utilization efficiency (NUE) in first (A) and second (B) year in chrysanthemum plants at harvest time (* indicates the level of significance at $p < 0.05$).

Table 8. Main effects of nutrient management and cultivar on phosphorus content (%) in different organs at harvest in chrysanthemum plants over the two years of application.

TMTS	Leaves	Stem	Flower Heads
First Year			
Nutrition Management (NM)			
CNM	0.19 b	0.15 b	0.33 b
INM	0.21 a	0.18 a	0.40 a
Significance	*	*	*
Cultivar (CV)			
CV 1	0.20	0.16	0.32
CV 2	0.21	0.17	0.31
Significance	ns	ns	ns
Interaction			
NM × CV	ns	ns	ns
Second Year			
Nutrition Management (NM)			
CNM	0.17 b	0.16 b	0.29 b
INM	0.19 a	0.18 a	0.33 a
Significance	*	*	*
Cultivar (CV)			
CV 1	0.18	0.19	0.31
CV 2	0.18	0.18	0.30
Significance	ns	ns	ns
Interaction			
NM × CV	ns	ns	ns

Different letters within each column indicate significant differences according to SNK test ($p \leq 0.05$). NS not significant * $p < 0.05$ and **0.01, indicate level of significance.

Concerning an economic point of view, the increased yield obtained with INM (140 stems m^{-2}) compared with CNM (118 stems m^{-2}) led consequently to an increase of gross production of exactly €50,600.00 (takings difference). This amount was much greater than the cost increases needed for INM compared to CNM (Table 9).

Table 9. Basic economic analysis of fertilizers cost, gross production, and takings difference obtained.

	Fertilizer Cost	Yield	Gross Production	Takings Difference
	(€ ha ⁻¹)	(stems m ⁻²)	(€ ha ⁻¹) *	(€)
CNM	1945.00	118	271,400.00	
INM	3144.00	140	322,000.00	+50,600.00

* Chrysanthemum price calculated at 2012–2013: 0.23 €/stem (ISMEA/2012–2013).

4. Discussion

In our study, mineral nutrient management (CNM) and integrated nutrient management (INM) were compared in chrysanthemum cultivation. The INM protocol, which combined the application of half the rate of CNM and seaweed extract, organic and biofertilizer (AMF + PSB), improved yield, cut stem quality traits, root morphology, as well as N accumulation and P content in tissues. Based on other research about INM practices [47–49], this protocol seems to be suitable in order to obtain advantages on profits and sustainability. Our aim was to verify a new mixture in order to reduce mineral fertilizer application, making chrysanthemum cultivation more sustainable, as well as highly profitable.

Compared to CNM, the INM protocol led to a significantly higher yield in terms of the number of secondary branches per m^{-2} (Table 3). This could be attributed to a better nutrient translocation in the plant, which led to the production of a greater number of axillary buds and therefore of secondary axes,

in line with Kale et al. [50] in *Salvia* and Nethra [51] in the Chinese aster. In other studies regarding biostimulant applications, yield also increased in seaweed treated plants influenced by cytokinin content, which enhances nutrient mobilization in plant organs [52].

Regarding cut stem quality traits (Table 3), our results are in agreement with Verma et al. [53], who applied an INM on chrysanthemum CV Roja. The treatment that consisted of *Azospirillum*, PSB, vermicompost, and 50% of recommended mineral NPK recorded the highest plant height, number of branches, and flowers per plant. Similar results were reported in *Crossandra* [54] and *Dahlia* [55]. The combination of biofertilizers with the recommended NPK dose yielded a higher flower production in *Limonium* [56] and *Calendula* [57].

In our study, a higher leaf area value was found in chrysanthemum plants under INM. According to De Lucia and Vecchiatti [58], the application of seaweed extract (*A. nodosum*) in *Lilium* CV Brindisi, greatly affected these parameters (12.3 cm² of treated plants, compared with 10.3 cm² of untreated plants). This was potentially due to the direct effect of the biostimulant containing betaine. The nutrient concentration present in both the N organic fertilizer and seaweed extract biostimulant cannot on its own explain the positive response as an increase in aerial organ dry weight (Table 4). In fact cytokinins have a considerable influence on nutrient mobilization in vegetative and reproductive organs [59].

Microbial inoculants are also good supplement with half the recommended mineral dose of fertilizer. Wu et al. [60] reported that *G. mossae* plus *B. megaterium* on maize increased plant growth and NPK assimilation. As regards the effect of applying INM, the chrysanthemum root development exhibited a remarkable increase in all parameters compared to CNM (Table 5). The root growth promoting activity has been observed in snapdragon, when a biostimulant was applied [61]. Previous research has shown that the brown seaweed extract, rich in auxin, improved lateral root formation when applied to mung bean [62]. A study carried out by Mancuso et al. [63] on potted *Vitis vinifera* under seaweed extracts, showed an increase in the total volume of the root system.

Concerning the nutrient uptake, Biswas et al. [64] and Adesemoye et al. [65] showed that PGPR (plant growth-promoting rhizobacteria) also influences this parameter through a more pronounced development of the root surface area. The INM seems to encourage a better uptake of mineral nutrients by plants, which results in a higher number of branches as well as leaf area, and more flowers [66].

The N uptake by chrysanthemum plants may be enhanced by the use of biofertilizers, possibly because they stimulate better root architecture or due to the influence of growth hormones contained in seaweed extracts. These substances can increase the ability of nutrient absorption as well as enzymatic activity, in agreement with Kumari et al. [67].

The N accumulation (g m⁻²) value in the INM treatment could be caused by the better availability and uptake of nutrients facilitated by the application of both mineral and organic fertilizers, biostimulants, and biofertilizers (Tables 6 and 7). Mahadik et al. [68] showed that the increase in N and P uptake by chrysanthemum plants was the highest with the application of *Azotobacter* plus PSB, 50% of RDF (Recommended Dose of Fertilizers) (100:100:100 kg ha⁻¹ NPK), and 10 t ha⁻¹ of vermicompost. Regarding the P content in plant tissue (Table 8), our findings are validated by similar results found in a number of earlier studies on bacteria.

Shirmardi et al. [69] reported that PGPR solubilizes the inorganic phosphate and produces IAA, thus improving plant growth by increasing P-uptake from the soil and its transport to plant shoots. A significant increase in sunflower growth parameters, including plant P content, was found in inoculated plants after inoculation with *Bacillus* sp., possibly due to the P-solubilizing, IAA-synthesizing, and root-colonizing of these strains [70], which increase nutrient uptake.

Richardson (1994) and Rodríguez and Fraga [71] studied the influence of several soil bacteria on the supply of P to plants as a consequence of their capacity for inorganic or organic P solubilization and, therefore, for improving plant growth performance. In addition, in a 1994 study, Garbaye [72] postulated that some PSB behave like mycorrhizal helper bacteria with a synergistic interaction.

Compared to the non-treated control, the combined application of mycorrhizal fungus and rhizobacteria significantly increased growth parameters, i.e., total fresh weight, aerial dry weight,

shoot length, and leaf area, in bananas. The leaf mineral content, i.e., N, P, and K, also increased significantly following the combined application of both microorganisms [73].

Finally, integrated nutrient management practices could be viable for sustainable floriculture on a commercial and profitable scale. Our data on the economics of chrysanthemum flowers are in agreement with those Verma, who showed that the cost of fertilizer can be saved with inoculation of both *Azospirillum* and PSB, obtaining higher flower yield compared to CNM. Angadi too, carried out a study that shed light on the combination of *Azospirillum*, PSB, 50% vermicompost, and 1/2 recommended NPK dose, giving the maximum net returns per euro invested.

5. Conclusions

The quality and quantity of fertilizers are the key factor affecting the growth, yield, and quality of cut flowers. Since chrysanthemum is an energy-intensive ornamental crop with a very high input of fertilizers, several experiments have been aimed at using alternative methods, reducing mineral fertilizers, and in particular the INM.

Our results shows that the INM protocol, 50% mineral RDF with N organic fertilizer plus biostimulant (seaweed extract) plus biofertilizer (microbial consortium of *Glomus* sp. and *Bacillus* sp.), is effective in enhancing yield, quality, root morphology, and nutrient uptake compared to RDF. This indicates the possibility of the sustainable, eco-friendly cultivation of chrysanthemum. In order to discern the influence of each component of INM mixture on yield and quality traits, future research is needed.

Author Contributions: B.L.: substantial contribution to the experimental work and data collection; D.L.: bibliographic research and support for drafting; G.C.: support with data analysis, interpretation, and critical revision of the manuscript; B.D.L.: experimental design and coordination of the work, interpretation of the data, and critical revision of the manuscript.

Funding: This research was funded by Chemicaverde Flor (Province of Bari).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Yousaf, M.; Li, J.; Lu, J.; Ren, T.; Cong, R.; Fahad, S.; Li, X. Effects of fertilization on crop production and nutrient-supplying capacity under rice-oilseed rape rotation system. *Sci. Rep.* **2017**, *7*, 1270. [[CrossRef](#)] [[PubMed](#)]
2. Stellacci, A.M.; Cristiano, G.; Rubino, P.; De Lucia, B.; Cazzato, E. Nitrogen uptake, nitrogen partitioning and N-use efficiency of container-grown Holm oak (*Quercus ilex* L.) under different nitrogen levels and fertilizer sources. *J. Food Agric. Environ.* **2013**, *11*, 990–994.
3. Mantovani, J.R.; Silveira, L.G.D.; Landgraf, P.R.C.; Santos, A.R.D.; Costa, B.D.S. Phosphorus rates and use of cattle manure in potted gerbera cultivation. *Ornam. Hortic.* **2017**, *23*, 412–418. [[CrossRef](#)]
4. Vitousek, P.M.; Aber, J.D.; Howarth, R.W.; Likens, G.E.; Matson, P.A.; Schindler, D.W.; Schlesinger, W.H.; Tilman, D.G. Human alteration of the global nitrogen cycle: Sources and consequences. *Ecol. Appl.* **1997**, *7*, 737–750. [[CrossRef](#)]
5. Tilman, D. The greening of the green revolution. *Nature* **1998**, *396*, 211–212. [[CrossRef](#)]
6. Berg, M.; Köpke, U.; Haas, G. Nitrate leaching: Comparing conventional, integrated and organic agricultural production systems. *Intern. Assoc. Hydrol. Sci.* **2002**, *273*, 131–136.
7. Chen, J.H. The combined use of chemical and organic fertilizers and/or biofertilizer for crop growth and soil fertility. In Proceedings of the International Workshop on Sustained Management of the Soil-Rhizosphere System for Efficient Crop Production and Fertilizer Use, Bangkok, Thailand, 16–20 October 2006; Volume 16, p. 20.
8. Hansen, B.; Thorling, L.; Schullehner, J.; Termansen, M.; Dalgaard, T. Groundwater nitrate response to sustainable nitrogen management. *Sci. Rep.* **2017**, *7*, 8566. [[CrossRef](#)] [[PubMed](#)]

9. Shu, J.; Wu, H.; Chen, M.; Peng, H.; Li, B.; Liu, R.; Liu, Z.; Wang, B.; Huang, T.; Hu, Z. Fractional removal of manganese and ammonia nitrogen from electrolytic metal manganese residue leachate using carbonate and struvite precipitation. *Water Res.* **2019**, *153*, 229–238. [[CrossRef](#)]
10. Torbert, H.A.; King, K.W.; Harmel, R.D. Impact of Soil Amendments on Reducing Phosphorus Losses from Runoff in Sod. *J. Environ. Qual.* **2005**, *34*, 1415. [[CrossRef](#)] [[PubMed](#)]
11. Bouwman, L.; Goldewijk, K.K.; Van Der Hoek, K.W.; Beusen, A.H.W.; Van Vuuren, D.P.; Willems, J.; Rufino, M.C.; Stehfest, E. Exploring global changes in nitrogen and phosphorus cycles in agriculture induced by livestock production over the 1900–2050 period. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 20882–20887. [[CrossRef](#)]
12. Zhang, X.; Davidson, E.A.; Mauzerall, D.L.; Searchinger, T.D.; Dumas, P.; Shen, Y. Managing nitrogen for sustainable development. *Nature* **2015**, *528*, 51–59. [[CrossRef](#)]
13. Murillo-Amador, B.; Morales-Prado, L.E.; Troyo-Diéguez, E.; Córdoba-Matson, M.V.; Hernández-Montiel, L.G.; Rueda-Puente, E.O.; Nieto-Garibay, A. Changing environmental conditions and applying organic fertilizers in *Origanum vulgare* L. *Front. Plant Sci.* **2015**, *6*, 549. [[CrossRef](#)]
14. Szparaga, A.; Kocira, S. Generalized logistic functions in modelling emergence of *Brassica napus* L. *PLoS ONE* **2018**, *13*, e0201980. [[CrossRef](#)] [[PubMed](#)]
15. Alori, E.T.; Babalola, O.O. Microbial Inoculants for Improving Crop Quality and Human Health in Africa. *Front. Microbiol.* **2018**, *9*, 2213. [[CrossRef](#)] [[PubMed](#)]
16. Fiorentino, N.; Ventorino, V.; Woo, S.L.; Pepe, O.; De Rosa, A.; Gioia, L.; Romano, I.; Lombardi, N.; Napolitano, M.; Colla, G.; et al. Trichoderma-based biostimulants modulate rhizosphere microbial populations and improve N uptake efficiency, yield, and nutritional quality of leafy vegetables. *Front. Plant Sci.* **2018**, *9*, 743. [[CrossRef](#)] [[PubMed](#)]
17. Wani, S.A.; Chand, S.; Wani, M.A.; Ramzan, M.; Hakeem, K.R. Azotobacter chroococcum—A Potential Biofertilizer in Agriculture: An Overview. In *Soil Science: Agricultural and Environmental Perspectives*; Springer International Publishing: Cham, Switzerland, 2016; pp. 333–348, ISBN 9783319344515.
18. Gruhn, P.; Goletti, F.; Yudelman, M. Integrated nutrient management, soil fertility, and sustainable agriculture: Current issues and future challenges. *Food Agric. Environ. Discuss. Pap.* **2000**, *32*.
19. Ertani, A.; Francioso, O.; Tinti, A.; Schiavon, M.; Pizzeghello, D.; Nardi, S. Evaluation of Seaweed Extracts From *Laminaria* and *Ascophyllum nodosum* spp. as Biostimulants in *Zea mays* L. Using a Combination of Chemical, Biochemical and Morphological Approaches. *Front. Plant Sci.* **2018**, *9*, 428. [[CrossRef](#)]
20. De Pascale, S.; Rouphael, Y.; Colla, G. Plant biostimulants: Innovative tool for enhancing plant nutrition in organic farming. *Eur. J. Hortic. Sci.* **2018**, *82*, 277–285. [[CrossRef](#)]
21. Chiaiese, P.; Corrado, G.; Colla, G.; Kyriacou, M.C.; Rouphael, Y. Renewable Sources of Plant Biostimulation: Microalgae as a Sustainable Means to Improve Crop Performance. *Front. Plant Sci.* **2018**, *9*, 1782. [[CrossRef](#)]
22. Nelson, W.R.; van Staden, J. Effect of seaweed concentrate on the growth of wheat. *South Afr. J. Sci.* **1986**, *82*, 199–200.
23. Abetz, P.; Young, C.L. The Effect of Seaweed Extract Sprays Derived from *Ascophyllum nodosum* on Lettuce and Cauliflower Crops. *Bot. Mar.* **1983**, *26*, 487–492. [[CrossRef](#)]
24. Crouch, I.J.; Van Staden, J. Effect of seaweed concentrate on the establishment and yield of greenhouse tomato plants. *J. Appl. Phycol.* **1992**, *4*, 291–296. [[CrossRef](#)]
25. Burchett, S.; Fuller, M.P.; Jellings, A.J. Application of seaweed extract improves winter hardiness of winter barley cv Igri. In Proceedings of the Society for Experimental Biology, Annual Meeting, The York University, Toronto, ON, Canada, 22–27 March 1998; Springer: Berlin/Heidelberg, Germany, 1998.
26. Zaidi, A.; Khan, M.S.; Ahemad, M.; Oves, M.; Wani, P.A. Recent Advances in Plant Growth Promotion by Phosphate-Solubilizing Microbes. In *Microbial Strategies for Crop Improvement*; Springer: Berlin/Heidelberg, Germany, 2009; pp. 23–50, ISBN 9783642019791.
27. Rafi, M.M.; Krishnaveni, M.S.; Charyulu, P.B.B.N. Phosphate-Solubilizing Microorganisms and Their Emerging Role in Sustainable Agriculture. In *Recent Developments in Applied Microbiology and Biochemistry*; Buddolla, V., Ed.; Academic Press: Dordrecht, The Netherlands, 2019; Chapter 17; pp. 223–233, ISBN 9780128163283. [[CrossRef](#)]
28. Canbolat, M.Y.; Bilen, S.; Çakmakçı, R.; Şahin, F.; Aydın, A. Effect of plant growth-promoting bacteria and soil compaction on barley seedling growth, nutrient uptake, soil properties and rhizosphere microflora. *Biol. Fertil. Soils* **2006**, *42*, 350–357. [[CrossRef](#)]

29. Ipsilantis, I.; Karamesouti, M.; Gasparatos, D. Beneficial Microorganisms for the Management of Soil Phosphorus. In *Sustainable Agriculture Reviews*; Lichtfouse, E., Ed.; Springer: Cham, Switzerland, 2018; Volume 32.
30. Rajkumar, M.; Nagendran, R.; Lee, K.J.; Lee, W.H.; Kim, S.Z. Influence of plant growth promoting bacteria and Cr6+ on the growth of Indian mustard. *Chemosphere* **2006**, *62*, 741–748. [[CrossRef](#)] [[PubMed](#)]
31. Yasmin, S.; Rahman Bakar, M.A.; Malik, K.A.; Hafeez, F.Y. Isolation, characterization and beneficial effects of rice associated plant growth promoting bacteria from Zanzibar soils. *J. Basic Microbiol.* **2004**, *44*, 241–252. [[CrossRef](#)]
32. Stewart, L.I.; Hamel, C.; Hogue, R.; Moutoglis, P. Response of strawberry to inoculation with arbuscular mycorrhizal fungi under very high soil phosphorus conditions. *Mycorrhiza* **2005**, *15*, 612–619. [[CrossRef](#)]
33. Li, H.; Smith, S.E.; Holloway, R.E.; Zhu, Y.; Smith, F.A. Arbuscular mycorrhizal fungi contribute to phosphorus uptake by wheat grown in a phosphorus-fixing soil even in the absence of positive growth responses. *New Phytol.* **2006**, *172*, 536–543. [[CrossRef](#)]
34. Estaun, V.; Camprubi, A.; Joner, E.J. Selecting arbuscular mycorrhizal fungi for field application. In *Mycorrhizal Technology in Agriculture*; Gianinazzi, S., Schuepp, H., Barea, J.M., Haselwandter, K., Eds.; Birkhauser Verlag: Berlin, Germany, 2002; pp. 249–259.
35. Wang, C.; Li, X.; Zhou, J.; Wang, G.; Dong, Y. Effects of arbuscular mycorrhizal fungi on growth and yield of cucumber plants. *Commun. Soil Sci. Plant Anal* **2008**, *39*, 499–509. [[CrossRef](#)]
36. Kapulnik, Y.; Tsror (Lahkim), L.; Zipori, I.; Hazanovsky, M.; Wininger, S.; Dag, A. Effect of AMF application on growth, productivity and susceptibility to Verticillium wilt of olives grown under desert conditions. *Symbiosis* **2010**, *52*, 103–111. [[CrossRef](#)]
37. Ortas, I. Effect of mycorrhiza application on plant growth and nutrient concentration in cucumber production under field conditions. *Span. J. Agric. Res.* **2010**, *8*, 116–122. [[CrossRef](#)]
38. Mwangi, A.M.K.; Kahangi, E.M.; Ateka, E.; Onguso, J.; Mukhongo, R.W.; Mwangi, E.K.; Jefwa, J.M. Growth effects of microorganisms based commercial products inoculated to tissue cultured banana cultivated in three different soils in Kenya. *Appl. Soil Ecol.* **2013**, *64*, 152–162. [[CrossRef](#)]
39. Lim, T.K. Chrysanthemum morifolium. In *Edible Medicinal and Non-Medicinal Plants*; Springer Netherlands: Dordrecht, The Netherlands, 2014; pp. 250–269.
40. Liu, D.; Liu, W.; Zhu, D.; Geng, M.; Zhou, W.; Yang, T. Nitrogen effects on total flavonoids, chlorogenic acid, and antioxidant activity of the medicinal plant *Chrysanthemum morifolium*. *J. Plant Nutr. Soil Sci.* **2010**, *173*, 268–274. [[CrossRef](#)]
41. Lunt, O.R.; Kofranek, A.M. Nitrogen and potassium nutrition of chrysanthemum. *Proc. Am. Soc. Hort. Sci* **1958**, *72*, 487–497.
42. Yoon, H.S.; Goto, T.; Kageyama, Y. Developing a Nitrogen Application Curve for Spray Chrysanthemums grown in Hydroponic System and its Practical Use in NFT System. *Engei Gakkai zasshi* **2000**, *69*, 416–422. [[CrossRef](#)]
43. Li, Y.; Jiang, Y.; Li, Z.; Zhao, L. A study of phosphorus uptake kinetic characteristic of four terrestrial economic plants. In *Proceedings of the 2009 Beijing International Environmental Technology Conference*, Beijing, China, 16–19 October 2009; pp. 140–145.
44. Vanek, V.; Balik, J.; Cerny, J.; Pavlik, M.; Pavlikova, D.; Tlustos, P.; Valtera, J. The Nutrition of Horticultural Plants. Academic Press: Prague, Czech Republic, 2012.
45. Tesi, R. Colture Protette, Ortoflorovivaismo. *Colt. Protette* **1987**.
46. Ceccon, P.; Fagnano, M.; Grignani, C.; Monti, M.; Orlandini, S. (Eds.) *Agronomia*; SES: Napoli, Italy, 2017.
47. Nammidevi, M.; Shalini, B.; Bhongle, S.A.; Khiratkar, S.D. Effect of bio-inoculants with graded doses of NPK on flowering, yield attributes and economics of annual chrysanthemum. *J. Soils Crop.* **2008**, *18*, 217–220.
48. Deshmukh, P.G.; Khiratkar, S.D.; Badge, S.A.; Bhongle, S.A. Effect of bio-inoculants with graded doses of NPK on growth and yield of gaillardia. *J. Soils Crop.* **2008**, *18*, 212–216.
49. Sunitha, H.M.; Hunje, R.; Vyakaranahal, B.S.; Bablad, H.B. Effect of plant population and integrated nutrient management on growth, seed yield and quality of African marigold (*Tagetes erecta* L.). *Karnataka J. Agric. Sci* **2010**, *23*, 783–786.
50. Kale, R.D.; Bano, K.; Sreenivasa, M.N.; Bagyaraj, D.J. Influence of worms cast on the grown and mycorrhizal colonization of two ornamental plants. *South Indian Hortic.* **1987**, *35*, 433–437.

51. Nethra, N.N. Effect PH, Organic and Inorganic Fertilizer on Growth, Yield and Post Harvest Life of China Aster (*Callistephus Chinensis* (L.) Nexx.). Ph.D. Dissertation, University of Agricultural Sciences, GKVK, Karnataka, India, 1996.
52. Gersani, M.; Kende, H. Studies on cytokinin-stimulated translocation in isolated bean leaves. *J. Plant Growth Regul.* **1982**, *1*, 161–171.
53. Verma, S.K.; Angadi, S.G.; Patil, V.S.; Mokashi, A.N.; Mathad, J.C.; Mummigatti, U.V. Growth, yield and quality of chrysanthemum (*Chrysanthemum morifolium* Ramat.) cv. Raja as influenced by integrated nutrient management. *Karn J. Agric. Sci.* **2011**, *24*, 681–683.
54. Narsimha Raju, S.; Haripriya, K. Integrated nutrient management in cossandra (*Crossandra infundibuliformis* L.) Cv. Dindigul local. *South Indian Hortic.* **2001**, *49*, 318–322.
55. Warade, A.P.; Golliwar, V.J.; Chopde, N.; Lanje, P.W.; Thakre, S.A. Effect of organic manures and bio-fertilizers on growth, flowering and yield of dahlia. *J. Soils Crop.* **2007**, *17*, 354–357.
56. Gajathri, H.N.; Jayaprased, K.V.; Narayanaswamy, P. Response of bio-fertilizers and their combined application with different levels of inorganic fertilizers in statice (*Limonium caspia*). *J. Ornam. Hortic.* **2004**, *7*, 70–74.
57. Shashikanth, G. Effect of Different Sources of Nutrients on Growth and Yield of Tall Marigold. Master's Thesis, University of Agricultural Sciences, Dharwad, India, 2005.
58. De Lucia, B.; Vecchietti, L. Type of bio-stimulant and application method effects on stem quality and root system growth in L.A. Lily. *Eur. J. Hortic. Sci.* **2012**, *77*, 10.
59. Khan, W.; Rayirath, U.P.; Subramanian, S.; Jithesh, M.N.; Rayorath, P.; Hodges, D.M.; Critchley, A.T.; Craigie, J.S.; Norrie, J.; Prithiviraj, B. Seaweed Extracts as Biostimulants of Plant Growth and Development. *J. Plant Growth Regul.* **2009**, *28*, 386–399. [[CrossRef](#)]
60. Wu, S.C.; Cao, Z.H.; Li, Z.G.; Cheung, K.C.; Wong, M.H. Effects of biofertilizer containing N-fixer, P and K solubilizers and AM fungi on maize growth: A greenhouse trial. *Geoderma* **2005**, *125*, 155–166. [[CrossRef](#)]
61. Cristiano, G.; Pallozzi, E.; Conversa, G.; Tufarelli, V.; De Lucia, B. Effects of an Animal-Derived Biostimulant on the Growth and Physiological Parameters of Potted Snapdragon (*Antirrhinum majus* L.). *Front. Plant Sci.* **2018**, *9*, 861. [[CrossRef](#)]
62. Crouch, I.J.; Van Staden, J. Evidence for rooting factors in a seaweed concentrate prepared from *Ecklonia maxima*. *J. Plant Physiol.* **1991**, *137*, 319–322. [[CrossRef](#)]
63. Mancuso, S.; Azzarello, E.; Mugnai, S.; Briand, X. Marine bioactive substances (IPA extract) improve foliar ion uptake and water stress tolerance in potted *Vitis vinifera* plants. *Adv. Hortic. Sci.* **2006**, *20*, 1000–1006.
64. Biswas, J.C.; Ladha, J.K.; Dazzo, F.B. Rhizobia Inoculation Improves Nutrient Uptake and Growth of Lowland Rice. *Soil Sci. Soc. Am. J.* **2000**, *64*, 1644. [[CrossRef](#)]
65. Adesemoye, A.O.; Torbert, H.A.; Kloepper, J.W. Enhanced plant nutrient use efficiency with PGPR and AMF in an integrated nutrient management system. *Can. J. Microbiol.* **2008**, *54*, 876–886. [[CrossRef](#)] [[PubMed](#)]
66. Patanwar, M.; Sharma, G.; Banjare, C.; Chandravanshi, D.; Sahu, E. Growth and development of chrysanthemum (*Dendranthema grandiflora* Tzvelev) as influenced by integrated nutrient management. *Ecscan* **2014**, *6*, 459–462.
67. Kumari, A.; Goyal, R.K.; Choudhary, M.; Sindhu, S.S. Effect of different nitrogen levels and biofertilizers on growth, yield and nutrient content of chrysanthemum. *Ann. Agric. Res. New Ser.* **2014**, *35*, 156–163.
68. Mahadik, M.K.; Dalal, S.R.; Patil, S.W.U. Dry matter production and nutrients composition of chrysanthemum in relation to integrated nutrient management. *J. Pharmacogn. Phytochem.* **2017**, *6*, 1879–1882.
69. Shirmardi, M.; Savaghebi, G.R.; Khavazi, K.; Akbarzadeh, A.; Farahbakhsh, M.; Rejali, F.; Sadat, A. Effect of Microbial Inoculants on Uptake of Nutrient Elements in Two Cultivars of Sunflower (*Helianthus annuus* L.) in Saline Soils. *Notulae Scientia Biologicae* **2010**, *2*, 57. [[CrossRef](#)]
70. Shahid, M.; Hameed, S.; Tariq, M.; Zafar, M.; Ali, A.; Ahmad, N. Characterization of mineral phosphate-solubilizing bacteria for enhanced sunflower growth and yield-attributing traits. *Ann. Microbiol.* **2015**, *65*, 1525–1536. [[CrossRef](#)]
71. Rodríguez, H.; Fraga, R. Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol. Adv.* **1999**, *17*, 319–339. [[CrossRef](#)]

72. Garbaye, J. Tansley Review No. 76 Helper bacteria: A new dimension to the mycorrhizal symbiosis. *New Phytol.* **1994**, *128*, 197–210. [[CrossRef](#)]
73. Rodríguez-Romero, A.S.; Guerra, M.S.P.; Del Carmen Jaizme-Vega, M. Effect of arbuscular mycorrhizal fungi and rhizobacteria on banana growth and nutrition. *Agron. Sustain. Dev.* **2005**, *25*, 395–399. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).