

Review

## Role of Cell Division Autoantigen-1 (CDA1) in Cell Proliferation and Fibrosis

Ban-Hock Toh <sup>1,\*</sup>, Yugang Tu <sup>2</sup>, Zemin Cao <sup>2</sup>, Mark E. Cooper <sup>2</sup> and Zhonglin Chai <sup>2</sup>

<sup>1</sup> Autoimmunity Laboratory, Centre for Inflammatory Diseases, Department of Medicine, Faculty of Medicine, Nursing and Health Sciences, Monash University, Clayton, Victoria 3168, Australia

<sup>2</sup> Diabetes and Metabolism Division, Baker IDI Heart and Diabetes Institute, 75 Commercial Rd, Melbourne, Victoria 3004, Australia; E-Mails: Yugang.Tu@bakeridi.edu.au (Y.T.); Zemin.Cao@bakeridi.edu.au (Z.C.); Mark.Cooper@bakeridi.edu.au (M.E.C.); Zhonglin.Chai@bakeridi.edu.au (Z.C.)

\* Author to whom correspondence should be addressed; E-Mail: Ban-Hock.Toth@monash.edu; Tel.: +61-3-95940713; Fax: +61-3-95940713.

Received: 5 August 2010; in revised form: 3 September 2010 / Accepted: 17 September 2010 /

Published: 30 September 2010

---

**Abstract:** Cell Division Autoantigen-1 (CDA-1) was discovered following screening a human expression library with serum from a patient with Discoid Lupus Erythematosus. CDA1, encoded by TSPYL2 on the X chromosome, shares anti-proliferative and pro-fibrotic properties with TGF- $\beta$ . It inhibits cell growth through p53, pERK1/2 and p21-mediated pathways and is implicated in tumorigenesis and the DNA damage response. Its pro-fibrotic property is mediated through cross-talk with TGF- $\beta$  that results in upregulation of extracellular matrix proteins. The latter properties have identified a key role for CDA1 in diabetes associated atherosclerosis. These dual properties place CDA1 as an attractive molecular target for treating tumors and vascular fibrosis including atherosclerosis and other vascular disorders associated with enhanced TGF- $\beta$  action and tissue scarring.

**Keywords:** TGF- $\beta$ ; tumorigenesis; DNA damage; atherosclerosis; diabetes mellitus

---

## 1. Introduction

Anti-nuclear autoantibodies (ANA) are useful diagnostic markers of systemic autoimmune diseases [1]. As ANAs typically react with highly conserved epitopes of biologically important nuclear antigens, they have also been useful for the molecular cloning and functional characterization of their cognate nuclear autoantigens. Thus, ANAs played pivotal roles in the molecular characterization of the extractable nuclear antigen “Sm” as components of the spliceosome and of proliferating cell nuclear antigen, “PCNA”, as a molecule implicated in cell proliferation.

Discoid lupus erythematosus (DLE) is primarily a cutaneous subset of systemic lupus erythematosus associated with low titer ANAs in up to 40% of patients [2]. We found that serum from a patient with DLE contains not only autoantibodies to phosphoepitopes of mitotic chromosomes [3] but also to a nuclear phosphoprotein that we named CDA1 (Cell Division Autoantigen 1) [4].

## 2. Molecular Cloning of CDA1 Using an Autoimmune DLE Serum

We cloned CDA1 by using the DLE serum to screen a human testis cDNA expression library. Its open reading frame comprises a cDNA of 2,079 base pairs encoding 693 amino acids with predicted molecular mass of 79,430 daltons [4]. Localization of CDA1 to the nucleus was based on the following observations [4]: Its predicted amino acid sequence contains four putative nuclear localization signals (NLSs); it localized to the nuclear but not cytoplasmic fraction of HeLa cells by immunoblotting; Myc-tagged or enhanced green fluorescent protein-tagged CDA1; and its N-terminal segment containing its four NLSs localized to the nucleus and nucleolus of transfected HeLa cells. Nuclear localization of CDA1 is supported by the study of Ueki *et al.* [5]. They reported that a partial cDNA clone (GenBank<sup>TM</sup> accession number [AB015345](#)) encoding aa 208–693 of CDA1 that included its third and fourth NLS localized to the nucleus of COS-7 cells. Thus the third and fourth NLS are sufficient to target CDA1 to the nucleus while the role of the first and second remains unknown.


## 3. CDA1 in Man and Mouse

CDA1 is encoded by TSPYL2 (TSPY-like 2, a homolog of TSPY on the X chromosome), a member of the testis-specific protein Y-encoded, TSPY-like/SET/nucleosome assembly protein-1 superfamily [6]. CDA1 has also been described as cutaneous T-cell lymphoma-associated antigen se20-4 [7], DENTT (Differentially Expressed Nucleolar TGF-beta1 Target) [8], and NP79 (a 79Kda nuclear protein encoded by a gene expressed in samples of normal hearts and hearts with common congenital defects) [9]. A mouse homologue of CDA1 was named CASK-interacting nucleosome assembly protein (CINAP). It was reported as a co-transcription factor in neurons with CASK binding and nucleosome assembly protein activity [10].

## 4. Domain Structure of CDA1

CDA1 has 693 amino acid residues with three structural domains (Figure 1): An N-terminal proline-rich domain, a central basic domain, and a C-terminal acidic domain. The four NLSs reside in the N-terminal proline-rich domain and central basic domain.

**Figure 1.** Domain structure of CDA1 showing proline-rich domain (Pr), basic and acidic domains with four nuclear localization signals (bars) and two consensus phosphorylation sites (flags).



## 5. CDA1 Requires its C-Terminal Domain for Anti-Proliferative Activity

Endogenous CDA1 levels are low in resting serum starved HeLa cells and dramatically elevated in serum stimulated cells. These observations suggest a role for CDA1 in cell growth. We explored its role in cell growth in stable HeLa cell lines transfected with Myc-tagged CDA1 in which CDA1 levels were regulated by a tetracyclin-off promoter [5]. Maximal expression of CDA1 transgene observed after 3–5 days of culture without tetracyclin was followed by dramatic arrest of cell growth and cell density after 4–5 days of culture. DNA synthesis assessed by BrdUrd uptake showed a gradual decline in the first three days followed by a virtual complete arrest by days 4–5. Incremental increases in CDA1 expression regulated by decreasing doxycycline concentrations, was accompanied by a corresponding incremental decrease of HeLa colony numbers, cell growth, cell density, and BrdUrd uptake. These observations indicate the level of CDA1 transgene expression, regulated cell growth and DNA synthesis. The inhibitory effects of CDA1 were not accompanied by a change in cell viability assessed by trypan blue dye exclusion, nor in cell cycle profiles assessed by flow cytometry. The results suggest that CDA1 may exert inhibitory effects at multiple stages of the cell cycle, acting as a negative regulator of cell cycle progression. The suggestion is consistent with the elevated levels of endogenous CDA1 in G<sub>1</sub>, S, and M phases of the cell cycle.

The acidic C-terminal tail of CDA1 is required for its inhibitory effects on cell growth and DNA synthesis as stable HeLa cell transfectants lacking this domain did not arrest cell growth or DNA synthesis. Thus transfectants with its N-terminal domain only, comprising of nucleotides 132–1,487, failed to affect cell growth [5]. The acidic C-terminal tail of CDA1 shares ~40% identity and 68% similarity with the acidic C-terminal tail of the leukemia associated protein SET [5]. The central region of CDA1 shares the same level of identity and similarity with most of the remainder of SET. SET is a 32-kDa ubiquitous nuclear protein identified fused to CAN in acute undifferentiated leukemia [11]. CAN has also been found fused with DEK in a subtype of acute myeloid leukemia [12], suggesting that CAN is an oncogene activated by fusion with SET or DEK. The only common motif of SET and DEK is their acidic domain. SET is a potent and specific inhibitor of protein phosphatase 2A [13].

The action of CDA1 in inhibiting cell growth is supported by observations that CDA1/DENTT inhibited the growth of human and mouse tumors and cell lines [14] and that its levels are upregulated in growth-arrested Jurkat T cells [15]. That CDA1 also regulates the growth and development of *Toxoplasma gondii* [16] suggests its growth inhibitory activity is evolutionarily conserved, a finding consistent with other nuclear autoantigens [1]. The ability of CDA1 to inhibit cell growth also required

phosphorylation of its two consensus Cdk phosphorylation sites. A CDA1 double mutant with both phosphorylation sites mutated to alanine (S20A and T340A), disabled its capacity to inhibit cell growth and failed to inhibit DNA synthesis in S-phase cells [4]. These observations suggest that in addition to its expression levels, phosphorylation of these two sites by cyclin/CDKs play important roles in regulating its role in cell proliferation. Cyclin D/Cdk4, cyclin A/Cdk2, and cyclin B/Cdk1 phosphorylated CDA1 *in vitro* at either or both phosphorylation sites, whereas cyclin E/CDK2 did not. Phospho-amino acid analysis confirmed phosphorylation of both serine and threonine sites by cyclin A/Cdk2 [4]. Our studies suggest that CDA1 may be differentially phosphorylated throughout the cell cycle on serine 20 and threonine 340 to regulate cell proliferation.

## 6. Proline-Rich N Terminal Domain of CDA1

CDA1 contains an N-terminal Pr domain not present in SET or other related proteins. Pr domains bind to Src Homology 3 (SH3) domains of intracellular and membrane-associated proteins and kinases [17,18] involved in signal transduction. The amino acid sequence motif of Pr domains for SH3 domain binding is PXXP with neighboring residues forming patterns specific for individual SH3 domain of different proteins [19]. CDA1 has three regions containing PXXP motifs, although no specific patterns are found for binding to known SH3 domains. The region also contains stretches of nine and five P residues. CDA1 may interact, through its Pr domain, with unique cognate protein(s), the identification of which may provide more information on its function.

## 7. CDA1 Regulates p21 Expression

Cell cycle progression is controlled by kinase activities of Cdks complexed with specific cyclins at certain cell cycle phases. These activities are negatively regulated by Cdk inhibitor proteins (CdkIs) of the Cip/Kip and INK4 families. The two families of Cdk inhibitors inactivate different classes of Cdks [20,21]. The Cip/Kip family member, p21<sup>Waf1/Cip1</sup> (p21) was first identified in a quaternary complex containing D cyclin, Cdk, and PCNA [22]. p21 inhibits cell proliferation and activities of several cyclin-Cdk complexes *in vitro* [23]. The tumor suppressor protein p53 controls transcriptional regulation of the p21 gene by binding to the p53 responsive distal region of the p21 promoter in response to intracellular signals such as DNA damage [24–26]. TGF- $\beta$  also transcriptionally regulates p21 by binding to the TGF- $\beta$  responsive element at the proximal region, which does not necessarily require functional p53 in HaCat cells [27]. The MEK/ERK intracellular signaling pathway upregulates p21 in response to stimulation by several factors including TGF- $\beta$  [28].

Using the Tet-Off HeLa cell line where CDA1 levels can be regulated by doxycycline concentrations [4] we have provided the following compelling data implicating CDA1 in regulating p21 transcription and expression [29]: A progressive time- and dose-dependent increase in CDA1 protein and mRNA levels was accompanied by a corresponding but delayed progressive increase in mRNA and protein expression of p21; a progressive increase followed by a subsequent decrease in CDA1 levels, was also matched by a corresponding rise and fall in p21 levels; and CDA1 overexpression activated the p21 promoter in a luciferase reporter assay.

Inhibiting CDA1 transgene overexpression after 48 h gradually decreased CDA1 protein levels, indicating a relatively long half-life of CDA1 protein in these cells. When CDA1 protein and p21 mRNA

levels dropped to a level similar to that seen endogenously, cell growth arrest was released, allowing cells to proliferate again [29]. This indicates that CDA1-overexpressing cells remained viable, consistent with our earlier report [4]. More importantly, this finding demonstrated that cell growth arrest and p21 induction are both quantitatively dependent on CDA1 expression.

## 8. CDA1 is an Upstream Regulator of p53 Expression

p53 is induced by DNA damage that results in transcription of p21. We found that p53 is also induced by CDA1 overexpression, an effect lost with CDA1 knockdown by siRNA [29]. The observation that CDA1 inactivates the ubiquitin ligase for p53 degradation (MDM2), without affecting p53 mRNA levels, suggests that the increased p53 levels are due to increased protein stability. MDM2's ability to degrade p53 is positively regulated by phosphorylation at S166 by Akt [30]. As CDA1 also inactivates Akt, it suggests a mechanism contributing to MDM2 inactivation [29]. Structural studies of the p21 gene promoter revealed a major p53 responsive element located <2.4 kb upstream of the p21 gene transcription start site [31]. This has been confirmed by global screening of p53 binding sites in the human genome [32]. The p21 promoter with deletion of the distal p53 responsive element (p21(p53-)-Luc) largely lost its basal and almost 100% of its responsive activities to CDA1 overexpression in HeLa cells. P53 knockdown by siRNA attenuated p21 induction by CDA1. The data suggest CDA1 induces p21 gene transcription by p53 binding to its distal responsive element of the p21 gene promoter.

These observations suggest that CDA1 may have a key role as an upstream regulator of p53 in the DNA damage response where p53 mediates cell cycle arrest, or apoptosis [33–39]. To test this hypothesis we treated cells with CPT, a molecule that inhibits DNA topoisomerase I by trapping it in cleavable complexes within transcribed chromosomal regions leading to irreversible DNA breaks [40,41]. CPT treatment indeed increased CDA1 mRNA and protein expression levels and induced p53 protein induction while CDA1 knockdown attenuated CPT-induced p53 induction [29]. This is the first report demonstrating CDA1 as an essential upstream regulator of p53 in the DNA damage response. The exact mechanism whereby CDA1 is up-regulated by DNA damage remains unclear.


## 9. CDA1 Activates ERK1/MAPK Pathways

Mild stimulation of the residual activity of p21(p53-)-Luc by CDA1 overexpression raised the possibility of other responsive elements in the downstream region of the p53 responsive element. MEK/ERK1/2 MAPK pathway may contribute to this residual activity as inhibition of this pathway by two MEK inhibitors, PD98059 and U0126, blocked the transcriptional induction of p21 and p21 promoter activity stimulated by CDA1 [29]. These data are consistent with reports of induction of p21 gene expression by this signaling pathway [42–45]. Thus, it is likely that both p53 and activated ERK1/2 MAPK activate the p21 gene by binding to its promoter. However the target of activated ERK1/2 MAPK on the p21 gene promoter remains unknown. ERK1/2 phosphorylation was induced by CDA1 overexpression suggesting that this may be responsible for activating the ERK1/2 MAPK pathway. A gradual decrease in ERK1/2 phosphorylation while CDA1 is still expressed at a high level probably reflects negative regulation of ERK1/2 by dual-specificity MAP kinase phosphatases [46,47]. Given the nuclear localization of CDA1, its overexpression may modulate upstream kinases such as


MEK, Raf, Ras, and cell surface receptor kinases linked to the MEK/1/2 MAPK pathway. However, it is possible that CDA1 can shuttle between the cytoplasm and nucleus since CDA1, also known as DENTT, has been detected in the cytoplasm, nucleus or both in certain tissues of adult mouse and monkey [48,49].

Thus, CDA1 appears to downregulate p21 expression through upregulation of p53 and phosphorylation of ERK1/2, as schematized in Figure 2.

**Figure 2.** CDA1 inhibits cell growth through upregulation of p21 mediated by p53 and pERK1/2. These activities of CDA1 appear to be mediated by binding of p53 and pERK1/2 to binding sites on the p21 gene promoter as outlined in Figure 3.



**Figure 3.** Schematic representation of p21 binding sites for p53, pERK1/2 and TGF $\beta$  responsive elements.



## 10. CDA1 Inhibits Cyclin-Dependent Kinases Through Upregulation of CDK Inhibitors

P21 expression induced by CDA1 is accompanied by inhibition of the activity of Cdk2 and Cdk1 activity, consistent with its binding to cyclins, Cdks, and the cyclin-Cdk complex [50–54]. The data

are consistent with our report that CDA1 overexpression inhibits DNA synthesis [4], an activity dependent on S-phase Cdks. However, CDA1 overexpression also inhibits the activities of Cdk4 and Cdk6. These activities cannot be attributed solely to induction of p21 expression as p21 plays a positive role in regulating activities of the D type cyclins-Cdk 4/6 [55–58], by promoting formation of active cyclin/Cdk complexes. It seems likely therefore that CDA1 overexpression may have also induced other Cdk inhibitor proteins that, together with p21, are potent inhibitors of G<sub>1</sub>, G<sub>1</sub>/S, S, and G<sub>2</sub>/M-phase Cdks.

## 11. CDA1 is Upregulated in Diabetes-Associated Atherosclerosis

Diabetes mellitus aggravates atherosclerosis-based cardiovascular disease [59]. Excess accumulation of extracellular matrix (ECM) leading to vascular fibrosis may contribute to increased plaque size, leading to luminal narrowing [60–63].

Streptozotocin induced diabetic *ApoE*<sup>−/−</sup> mouse is a well characterized animal model of diabetes accelerated atherosclerosis. These mice develop increased vascular ECM accumulation associated with rapid development of atherosclerotic plaques similar to human atherosclerosis [62–66].

Using this model, we found that after 10 weeks of diabetes mRNA levels of the CDA1-encoding gene *Tspyl2* and profibrotic growth factors *Tgf-β* and *Ctgf* increased 1.5 to 2-fold in aortas of *ApoE*<sup>−/−</sup> mice.[67]. At 20 weeks of diabetes, expression of these genes increased by more than 5-fold (*Tspyl2*), 10-fold (*Tgf-β*) and 15-fold (*Ctgf*), compared with age matched non-diabetic animals. These findings are consistent with our previous reports [62,63]. Our observation suggests that CDA1 is induced at an earlier stage, before the disease becomes manifest, consistent with a role for CDA1 in the development and progression of diabetes associated atherosclerosis in these mice.

TGF-β is the major growth factor implicated in vascular matrix accumulation via the SMAD signaling pathway [27,68–70]. TGF-β, together with angiotensin II, AGE, glucose and mechanical stretch are enhanced in the diabetic vasculature. These agents likely stimulate ECM accumulation and vascular remodeling through TGF-β-dependent and -independent signaling pathways. Targeting downstream signaling of TGF-β could therefore be beneficial in reducing ECM accumulation. Indeed, blockade of the renin-angiotensin system with drugs such as ACE inhibitors reduced fibrosis and attenuated diabetes associated atherosclerosis, partly via TGF-β-dependent pathways [62]. However, targeting TGF-β mediated ECM accumulation in the atherosclerotic plaque at a later stage remains controversial, as it may decrease the fibrous cap and increase the risk of plaque rupture [28,71].

## 12. CDA1 and TGFβ Cross-Talk

The MAPK pathway and transcriptional upregulation of p21<sup>Waf1/Cip1</sup> are both regulated by TGF-β [28,71–73]. Cross-talk between CDA1 and TGF-β is supported by the observation that CDA1 is upregulated by TGF-β and that CDA1 increased luciferase activities of TGF-β reporter constructs in some lung cancer cell lines [8,48,49]. These findings implicate CDA1 in TGF-β signaling and its biological functions. The pro-fibrotic activity of TGF-β has been suggested to play a central role in the promotion of diabetic complications, such as nephropathy and macrovascular diseases [72,74].

We carried out studies in vascular smooth muscle cells (VSMCs) to address the hypothesis that an increase in CDA1-encoding gene *Tspyl2* expression in diabetic atherosclerosis enhances the level of


TGF- $\beta$  that promotes ECM accumulation, TGF- $\beta$  treatment rapidly increased CDA1 protein levels followed by SMAD3 phosphorylation. CDA1 protein levels gradually decreased for 1 to 2 h after peaking, rising again at 4 to 5 h. The first increase in CDA1 is apparently a result of exogenous TGF- $\beta$  stimulation. The second peak in CDA1 at 4 to 5 h may reflect an autocrine effect of endogenous TGF- $\beta$ , consistent with a previous observation that the protein level of CDA1 is increased by TGF- $\beta$  treatment at 4 and 8 h in a human lung cell line

Overproduction of the CDA1-encoding gene *Tspyl2* in VSMCs increased expression of *Ctgf* and ECM genes, including collagen I, III and IV, and fibronectin. Conversely, siRNA-mediated knockdown of *Tspyl2* markedly abrogated TGF- $\beta$ -dependent expression of *Ctgf* and ECM genes. Our data clearly show that CDA1 promotes the pro-fibrotic effects of TGF- $\beta$  in the development of atherosclerosis in diabetes.

TGF- $\beta$  mediates its effect on vascular fibrosis [75] and diabetic nephropathy [76,77] through SMAD3. Knockdown of *Tspyl2*, the gene encoding CDA1, also attenuated TGF- $\beta$ -stimulated SMAD3 phosphorylation and the stimulatory effect of TGF- $\beta$  on its target genes associated with ECM production, including *Ctgf*, collagen I, III and IV, and fibronectin. The observations support the potential efficacy of targeting CDA1 to inhibit or reduce ECM production arising from enhanced TGF- $\beta$  action coordinated by elevated CDA1 production. Targeting TGF- $\beta$  or TGF- $\beta$  receptors result in deleterious effects due to inhibition of its other biological effects. In contrast, targeting CDA1 would most likely block only the pro-fibrotic action of TGF- $\beta$ .

TGF- $\beta$  receptor type I has dual specificity kinase activities that phosphorylate R-SMAD proteins including SMAD3 [78,79] and activate the ERK MAPK pathway. This is consistent with a role for CDA1, not only in enhancing TGF- $\beta$ -stimulated SMAD3 activity, but also in activating ERK MAPK, a pathway implicated in diabetic complications [80]. As crosstalk between SMAD and MAPK pathways plays an important role in fibrosis in the diabetic vasculature and kidney [81], we suggest that the cross-talk is mediated by CDA1 acting through TGF- $\beta$  receptor 1 (Figure 4).

**Figure 4.** Schematic representation of the promotion of fibrosis arising from cross-talk between CDA1 and TGF- $\beta$ .





### 13. Conclusions

CDA1 shares anti-proliferative and pro-fibrotic properties with TGF- $\beta$ . Our findings suggest that CDA1 plays a key role in these TGF $\beta$ -dependent activities. These properties of CDA1 implicate CDA1 as a negative regulator of tumorigenesis, and as a key player in p53-dependent DNA damage responses and fibrotic diseases. Targeting CDA1 may therefore be effective in the control of tumor growth and in attenuating the profibrotic action of TGF- $\beta$ . Therefore, CDA1 appears to be an attractive molecular target for treating neoplasia and vascular fibrosis including atherosclerosis and other vascular disorders associated with enhanced TGF- $\beta$  action and tissue scarring as commonly seen in diabetes.

### References and Notes

1. Tan, E.M. Antinuclear antibodies: Diagnostic markers for autoimmune diseases and probes for cell biology. *Adv. Immunol.* **1989**, *44*, 93–151.
2. Callen, J.P. Chronic cutaneous lupus erythematosus. Clinical, laboratory, therapeutic, and prognostic examination of 62 patients. *Arch. Dermatol.* **1982**, *118*, 412–416.
3. Hirota, T.; Lipp, J.J.; Toh, B.H.; Peters, J.M. Histone H3 serine 10 phosphorylation by Aurora B causes HP1 dissociation from heterochromatin. *Nature* **2005**, *438*, 1176–1180.
4. Chai, Z.; Sarcevic, B.; Mawson, A.; Toh, B.H. SET-related cell division autoantigen-1 (CDA1) arrests cell growth. *J. Biol. Chem.* **2001**, *276*, 33665–33674.
5. Ueki, N.; Oda, T.; Kondo, M.; Yano, K.; Noguchi, T.; Muramatsu, M. Selection system for genes encoding nuclear-targeted proteins. *Nat. Biotechnol.* **1998**, *16*, 1338–1342.
6. HGNC: 24358; GeneID: 64061; Available online: [http://www.genenames.org/data/hgnc\\_data.php?hgnc\\_id=24358](http://www.genenames.org/data/hgnc_data.php?hgnc_id=24358) (accessed on 1 September 2010).
7. Eichmüller, S.; Usener, D.; Dummer, R.; Stein, A.; Thiel, D.; Schadendorf, D. Serological detection of cutaneous T-cell lymphoma-associated antigens. *Proc. Natl. Acad. Sci. U. S. A.* **2001**, *98*, 629–634.
8. Ozbun, L.L.; You, L.; Kiang, S.; Angdisen, J.; Martinez, A.; Jakowlew, S.B. Identification of differentially expressed nucleolar TGF-beta1 target (DENTT) in human lung cancer cells that is a new member of the TSPY/SET/NAP-1 superfamily. *Genomics* **2001**, *73*, 179–193.
9. Sun, G.; Yuen C.S.; Yuan, Y.; Wang C.K.; Qiu, G.; Sun, K.; Ping L.M. Isolation of differentially expressed genes in human heart tissues. *Biochim. Biophys. Acta* **2002**, *1588*, 241–246.
10. Wang, G.S.; Hong, C.J.; Yen, T.Y.; Huang, H.Y.; Ou, Y.; Huang, T.N.; Jung, W.G.; Kuo, T.Y.; Sheng, M.; Wang, T.F.; Hsueh, Y.P. Transcriptional modification by a CASK-interacting nucleosome assembly protein. *Neuron* **2004**, *42*, 113–128.
11. von Lindern, M.; van Baal, S.; Wiegant, J.; Raap, A.; Hagemeijer, A.; Grosveld, G. Can, a putative oncogene associated with myeloid leukemogenesis, may be activated by fusion of its 3' half to different genes: Characterization of the set gene. *Mol. Cell Biol.* **1992**, *12*, 3346–3355.
12. von Lindern, M.; Fornerod, M.; Soekarman, N.; van Baal, S.; Jaegle, M.; Hagemeijer, A.; Bootsma, D.; Grosveld, G. Translocation t(6;9) in acute non-lymphocytic leukaemia results in the formation of a DEK-CAN fusion gene. *Baillieres Clin. Haematol.* **1992**, *5*, 857–879.

13. Li, M.; Makkinje, A.; Damuni, Z. The myeloid leukemia-associated protein SET is a potent inhibitor of protein phosphatase 2A. *J. Biol. Chem.* **1996**, *271*, 11059–11062.
14. Kandalaf, L.E.; Zudaire, E.; Portal-Nunez, S.; Cuttitta, F.; Jakowlew, S.B. Differentially Expressed Nucleolar TGF- $\beta$ 1 Target (DENTT) exhibits an inhibitory role on tumorigenesis. *Carcinog.* **2008**, *29*, 1282–1289.
15. Ito, T.; Tsukumo, S.; Suzuki, N.; Motohashi, H.; Yamamoto, M.; Fujii-Kuriyama, Y.; Mimura, J.; Lin, T. M.; Peterson, R.E.; Tohyama, C.; Nohara, K. A Constitutively Active Arylhydrocarbon Receptor Induces Growth Inhibition of Jurkat T Cells through Changes in the Expression of Genes Related to Apoptosis and Cell Cycle Arrest. *J. Biol. Chem.* **2004**, *279*, 25204–25210.
16. Radke, J.R.; Donald, R.G.; Eibs, A.; Jerome, M.E.; Behnke, M.S.; Liberator, P.; White, M.W. Changes in the expression of human cell division autoantigen-1 influence *Toxoplasma gondii* growth and development. *PLoS Pathog.* **2006**, *2*, e105.
17. Morton, C.J.; Campbell, I.D. SH3 domains. Molecular ‘Velcro’. *Curr. Biol.* **1994**, *4*, 615–617.
18. Birge, R.B.; Knudsen, B.S.; Besser, D.; Hanafusa, H. SH2 and SH3-containing adaptor proteins: Redundant or independent mediators of intracellular signal transduction. *Genes Cells* **1996**, *1*, 595–613.
19. Kaneko, T.; Li, L.; Li, S.S. The SH3 domain—a family of versatile peptide- and protein-recognition module. *Front. Biosci.* **2008**, *13*, 4938–4952.
20. Sherr, C.J.; Roberts, J.M. CDK inhibitors: Positive and negative regulators of G1-phase progression. *Genes Dev.* **1999**, *13*, 1501–1512.
21. Xiong, Y.; Hannon, G.J.; Zhang, H.; Casso, D.; Kobayashi, R.; Beach, D. p21 is a universal inhibitor of cyclin kinases. *Nature* **1993**, *366*, 701–704.
22. Xiong, Y.; Zhang, H.; Beach, D. D type cyclins associate with multiple protein kinases and the DNA replication and repair factor PCNA. *Cell* **1992**, *71*, 505–514.
23. Harper, J.W.; Adami, G.R.; Wei, N.; Keyomarsi, K.; Elledge, S.J. The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell* **1993**, *75*, 805–816.
24. Dulic, V.; Kaufmann, W.K.; Wilson, S.J.; Tlsty, T.D.; Lees, E.; Harper, J.W.; Elledge, S.J.; Reed, S.I. p53-dependent inhibition of cyclin-dependent kinase activities in human fibroblasts during radiation-induced G1 arrest. *Cell* **1994**, *76*, 1013–1023.
25. Waga, S.; Hannon, G.J.; Beach, D.; Stillman, B. The p21 inhibitor of cyclin-dependent kinases controls DNA replication by interaction with PCNA. *Nature* **1994**, *369*, 574–578.
26. Peter, M.; Herskowitz, I. Joining the complex: Cyclin-dependent kinase inhibitory proteins and the cell cycle. *Cell* **1994**, *79*, 181–184.
27. Datto, M.B.; Yu, Y.; Wang, X.F. Functional Analysis of the Transforming Growth Factor beta Responsive Elements in the WAF1/Cip1/p21 Promoter. *J. Biol. Chem.* **1995**, *270*, 28623–28628.
28. Kim, Y.K.; Bae, G.U.; Kang, J.K.; Park, J.W.; Lee, E.K.; Lee, H.Y.; Choi, W.S.; Lee, H.W.; Han, J.W. Cooperation of H<sub>2</sub>O<sub>2</sub>-mediated ERK activation with Smad pathway in TGF- $\beta$ 1 induction of p21WAF1/Cip1. *Cell. Signalling* **2006**, *18*, 236–243.
29. Tu, Y.; Wu, W.; Wu, T.; Cao, Z.; Wilkins, R.; Toh, B.H.; Cooper, M.; Chai, Z. Antiproliferative autoantigen CDA1 transcriptionally upregulates p21Waf1/Cip1 by activating p53 and MEK/ERK1/2 MAPK pathways. *J. Biol. Chem.* **2007**, *282*, 11722–11731.

30. Zhou, B.P.; Liao, Y.; Xia, W.; Zou, Y.; Spohn, B.; Hung, M.C. HER-2/neu induces p53 ubiquitination via Akt-mediated MDM2 phosphorylation. *Nat. Cell Biol.* **2001**, *3*, 973–982.
31. el-Deiry, W.S.; Harper, J. W.; O'Connor, P.M.; Velculescu, V.E.; Canman, C.E.; Jackman, J.; Pietenpol, J.A.; Burrell, M.; Hill, D.E.; Wang, Y.; *et al.* WAF1/CIP1 is induced in p53-mediated G1 arrest and apoptosis. *Cancer Res.* **1994**, *54*, 1169–1174.
32. Wei, C.L.; Wu, Q.; Vega, V.B.; Chiu, K.P.; Ng, P.; Zhang, T.; Shahab, A.; Yong, H.C.; Fu, Y.; Weng, Z.; Liu, J.; Zhao, X.D.; Chew, J.L.; Lee, Y.L.; Kuznetsov, V.A.; Sung, W.K.; Miller, L.D.; Lim, B.; Liu, E.T.; Yu, Q.; Ng, H.H.; Ruan, Y. A global map of p53 transcription-factor binding sites in the human genome. *Cell* **2006**, *124*, 207–219.
33. Lowe, S.W.; Schmitt, E.M.; Smith, S.W.; Osborne, B.A.; Jacks, T. p53 is required for radiation-induced apoptosis in mouse thymocytes. *Nature* **1993**, *362*, 847–849.
34. Lozano, G.; Elledge, S.J. p53 sends nucleotides to repair DNA. *Nature* **2000**, *404*, 24–25.
35. Tanaka, H.; Arakawa, H.; Yamaguchi, T.; Shiraishi, K.; Fukuda, S.; Matsui, K.; Takei, Y.; Nakamura, Y. A ribonucleotide reductase gene involved in a p53-dependent cell-cycle checkpoint for DNA damage. *Nature* **2000**, *404*, 42–49.
36. Caspari, T. How to activate p53. *Curr. Biol.* **2000**, *10*, R315–317.
37. Colman, M.S.; Afshari, C.A.; Barrett, J.C. Regulation of p53 stability and activity in response to genotoxic stress. *Mutat. Res.* **2000**, *462*, 179–188.
38. Zhou, B.B.; Elledge, S.J. The DNA damage response: putting checkpoints in perspective. *Nature* **2000**, *408*, 433–439.
39. Clarke, A.R.; Purdie, C.A.; Harrison, D.J.; Morris, R.G.; Bird, C.C.; Hooper, M.L.; Wyllie, A.H. Thymocyte apoptosis induced by p53-dependent and independent pathways. *Nature* **1993**, *362*, 849–852.
40. Hertzberg, R.P.; Busby, R.W.; Caranfa, M.J.; Holden, K.G.; Johnson, R.K.; Hecht, S.M.; Kingsbury, W.D. Irreversible trapping of the DNA-topoisomerase I covalent complex. Affinity labeling of the camptothecin binding site. *J. Biol. Chem.* **1990**, *265*, 19287–19295.
41. Gupta, R.S.; Gupta, R.; Eng, B.; Lock, R.B.; Ross, W.E.; Hertzberg, R.P.; Caranfa, M.J.; Johnson, R.K. Camptothecin-resistant mutants of Chinese hamster ovary cells containing a resistant form of topoisomerase I. *Cancer Res.* **1988**, *48*, 6404–6410.
42. Hu, P.P.; Shen, X.; Huang, D.; Liu, Y.; Counter, C.; Wang, X.F. The MEK pathway is required for stimulation of p21(WAF1/CIP1) by transforming growth factor-beta. *J. Biol. Chem.* **1999**, *274*, 35381–35387.
43. Kivinen, L.; Laiho, M. Ras- and mitogen-activated protein kinase kinase-dependent and -independent pathways in p21Cip1/Waf1 induction by fibroblast growth factor-2, platelet-derived growth factor, and transforming growth factor-beta1. *Cell Growth Differ.* **1999**, *10*, 621–628.
44. Lee, B.; Moon, S.K. Ras/ERK signaling pathway mediates activation of the p21WAF1 gene promoter in vascular smooth muscle cells by platelet-derived growth factor. *Arch. Biochem. Biophys.* **2005**, *443*, 113–119.
45. Lee, Y.; Mantel, C.; Anzai, N.; Braun, S.E.; Broxmeyer, H.E. Transcriptional and ERK1/2-dependent synergistic upregulation of p21(cip1/waf1) associated with steel factor synergy in MO7e. *Biochem. Biophys. Res. Commun.* **2001**, *280*, 675–683.

46. Camps, M.; Nichols, A.; Arkinstall, S. Dual specificity phosphatases: A gene family for control of MAP kinase function. *Faseb. J.* **2000**, *14*, 6–16.
47. Dickinson, R.J.; Keyse, S.M. Diverse physiological functions for dual-specificity MAP kinase phosphatases. *J. Cell Sci.* **2006**, *119*, 4607–4615.
48. Ozbun, L.L.; Martinez, A.; Jakowlew, S.B. Differentially expressed nucleolar TGF-beta1 target (DENTT) shows tissue-specific nuclear and cytoplasmic localization and increases TGF-beta1-responsive transcription in primates. *Biochim. Biophys. Acta* **2005**, *1728*, 163–180.
49. Martinez, A.; Ozbun, L.L.; Angdisen, J.; Jakowlew, S.B. Expression of differentially expressed nucleolar transforming growth factor-beta1 target (DENTT) in adult mouse tissues. *Dev. Dyn.* **2002**, *224*, 186–199.
50. Lin, J.; Reichner, C.; Wu, X.; Levine, A.J. Analysis of wild-type and mutant p21WAF-1 gene activities. *Mol. Cell Biol.* **1996**, *16*, 1786–1793.
51. Russo, A.A.; Jeffrey, P.D.; Patten, A.K.; Massague, J.; Pavletich, N.P. Crystal structure of the p27Kip1 cyclin-dependent-kinase inhibitor bound to the cyclin A-Cdk2 complex. *Nature* **1996**, *382*, 325–331.
52. Chen, J.; Jackson, P.K.; Kirschner, M.W.; Dutta, A. Separate domains of p21 involved in the inhibition of Cdk kinase and PCNA. *Nature* **1995**, *374*, 386–388.
53. Chen, J.; Saha, P.; Kornbluth, S.; Dynlacht, B.D.; Dutta, A. Cyclin-binding motifs are essential for the function of p21CIP1. *Mol. Cell Biol.* **1996**, *16*, 4673–4682.
54. Nakanishi, M.; Robetorye, R.S.; Adami, G.R.; Pereira-Smith, O.M.; Smith, J.R. Identification of the active region of the DNA synthesis inhibitory gene p21Sdi1/CIP1/WAF1. *EMBO J.* **1995**, *14*, 555–563.
55. Zhang, H.; Hannon, G.J.; Beach, D. p21-containing cyclin kinases exist in both active and inactive states. *Genes Dev.* **1994**, *8*, 1750–1758.
56. Sherr, C.J.; Roberts, J.M. Living with or without cyclins and cyclin-dependent kinases. *Genes Dev.* **2004**, *18*, 2699–2711.
57. LaBaer, J.; Garrett, M.D.; Stevenson, L.F.; Slingerland, J.M.; Sandhu, C.; Chou, H.S.; Fattaey, A.; Harlow, E. New functional activities for the p21 family of CDK inhibitors. *Genes Dev.* **1997**, *11*, 847–862.
58. Cheng, M.; Olivier, P.; Diehl, J.A.; Fero, M.; Roussel, M.F.; Roberts, J.M.; Sherr, C.J. The p21(Cip1) and p27(Kip1) CDK ‘inhibitors’ are essential activators of cyclin D-dependent kinases in murine fibroblasts. *EMBO J.* **1999**, *18*, 1571–1583.
59. Basson, M. Cardiovascular disease. *Nature* **2008**, *451*, 903.
60. Rumble, J.R.; Cooper, M.E.; Soulis, T.; Cox, A.; Wu, L.; Youssef, S.; Jasik, M.; Jerums, G.; Gilbert, R. E. Vascular hypertrophy in experimental diabetes. Role of advanced glycation end products. *J. Clin. Invest.* **1997**, *99*, 1016–1027.
61. Cao, Z.M.; Hulthen, U.L.; Allen, T.J.; Cooper, M.E. Angiotensin Converting Enzyme Inhibition and Calcium Antagonism Attenuate Streptozotocin-Diabetes-Associated Mesenteric Vascular Hypertrophy Independently of Their Hypotensive Action. *J. Hypertens.* **1998**, *16*, 793–799.

62. Candido, R.; Jandeleit-Dahm, K.A.; Cao, Z.; Nesteroff, S.P.; Burns, W.C.; Twigg, S.M.; Dilley, R.J.; Cooper, M.E.; Allen, T.J. Prevention of accelerated atherosclerosis by angiotensin-converting enzyme inhibition in diabetic apolipoprotein E-deficient mice. *Circulation* **2002**, *106*, 246–253.
63. Candido, R.; Allen, T.J.; Lassila, M.; Cao, Z.; Thallas, V.; Cooper, M.E.; Jandeleit-Dahm, K.A. Irbesartan but not amlodipine suppresses diabetes-associated atherosclerosis. *Circulation* **2004**, *109*, 1536–1542.
64. Hsueh, W.; Abel, E.D.; Breslow, J.L.; Maeda, N.; Davis, R.C.; Fisher, E.A.; Dansky, H.; McClain, D. A.; McIndoe, R.; Wassef, M.K.; Rabadan-Diehl, C.; Goldberg, I.J. Recipes for creating animal models of diabetic cardiovascular disease. *Circ. Res.* **2007**, *100*, 1415–1427.
65. Leask, A.; Abraham, D. J. TGF-beta signaling and the fibrotic response. *FASEB J.* **2004**, *18*, 816–827.
66. Sorescu, D. Smad3 mediates angiotensin II- and TGF-beta1-induced vascular fibrosis: Smad3 thickens the plot. *Circ. Res.* **2006**, *98*, 988–989.
67. Pham, Y.; Tu, Y.; Wu, T.; Allen, T. J.; Calkin, A.C.; Watson, A.M.; Li, J.; Jandeleit-Dahm, K.A.; Toh, B.H.; Cao, Z.; Cooper, M.E.; Chai, Z. Cell division autoantigen 1 plays a profibrotic role by modulating downstream signalling of TGF-beta in a murine diabetic model of atherosclerosis. *Diabetologia* **2010**, *53*, 170–179.
68. Hayden, M.R.; Tyagi, S.C. Arteriogenesis: Angiogenesis within Unstable Atherosclerotic Plaque—Interactions with Extracellular Matrix. *Curr. Interv. Cardiol. Rep.* **2000**, *2*, 218–227.
69. Schmidt, A.; Lorkowski, S.; Seidler, D.; Breithardt, G.; Buddecke, E. TGF-beta1 generates a specific multicomponent extracellular matrix in human coronary SMC. *Eur. J. Clin. Invest.* **2006**, *36*, 473–482.
70. Datto, M.B.; Li, Y.; Panus, J.F.; Howe, D.J.; Xiong, Y.; Wang, X.F. Transforming growth factor beta induces the cyclin-dependent kinase inhibitor p21 through a p53-independent mechanism. *Proc. Nat. Acad. Sci. U. S. A.* **1995**, *92*, 5545–5549.
71. Goldberg, H.J.; Huszar, T.; Mozes, M.M.; Rosivall, L.; Mucsi, I. Overexpression of the type II transforming growth factor-beta receptor inhibits fibroblasts proliferation and activates extracellular signal regulated kinase and c-Jun N-terminal kinase. *Cell Biol. Int.* **2002**, *26*, 165–174.
72. Lee, M.K.; Pardoux, C.; Hall, M.C.; Lee, P.S.; Warburton, D.; Qing, J.; Smith, S.M.; Derynck, R. TGF-beta activates Erk MAP kinase signalling through direct phosphorylation of ShcA. *EMBO J.* **2007**, *26*, 3957–3967.
73. Wang, A.; Ziyadeh, F.N.; Lee, E.Y.; Pyagay, P.E.; Sung, S.H.; Sheardown, S.A.; Laping, N.J.; Chen, S. Interference with TGF-beta signaling by Smad3-knockout in mice limits diabetic glomerulosclerosis without affecting albuminuria. *Am. J. Physiol. Renal. Physiol.* **2007**, *293*, F1657–F1665.
74. Yokoyama, H.; Deckert, T. Central role of TGF-beta in the pathogenesis of diabetic nephropathy and macrovascular complications: A hypothesis. *Diabet. Med.* **1996**, *13*, 313–320.
75. Twigg, S.M.; Cooper, M.E. The time has come to target connective tissue growth factor in diabetic complications. *Diabetologia* **2004**, *47*, 965–968.

76. Wang, W.; Huang, X.R.; Canlas, E.; Oka, K.; Truong, L.D.; Deng, C.; Bhowmick, N.A.; Ju, W.; Bottinger, E.P.; Lan, H.Y. Essential role of Smad3 in angiotensin II-induced vascular fibrosis. *Circ. Res.* **2006**, *98*, 1032–1039.
77. Li, J.H.; Huang, X.R.; Zhu, H.J.; Johnson, R.; Lan, H.Y. Role of TGF-beta signaling in extracellular matrix production under high glucose conditions. *Kidney Int.* **2003**, *63*, 2010–2019.
78. Macias-Silva, M.; Abdollah, S.; Hoodless, P.A.; Pirone, R.; Attisano, L.; Wrana, J.L. MADR2 is a substrate of the TGFbeta receptor and its phosphorylation is required for nuclear accumulation and signaling. *Cell* **1996**, *87*, 1215–1224.
79. Liu, X.; Sun, Y.; Constantinescu, S.N.; Karam, E.; Weinberg, R.A.; Lodish, H.F. Transforming growth factor beta-induced phosphorylation of Smad3 is required for growth inhibition and transcriptional induction in epithelial cells. *Proc. Nat. Acad. Sci. U. S. A.* **1997**, *94*, 10669–10674.
80. Li, J.H.; Wang, W.; Huang, X.R.; Oldfield, M.; Schmidt, A.M.; Cooper, M.E.; Lan, H.Y. Advanced glycation end products induce tubular epithelial-myofibroblast transition through the RAGE-ERK1/2 MAP kinase signaling pathway. *Am. J. Pathol.* **2004**, *164*, 1389–1397.
81. Li, J.H.; Huang, X.R.; Zhu, H.J.; Oldfield, M.; Cooper, M.; Truong, L.D.; Johnson, R.J.; Lan, H.Y. Advanced glycation end products activate Smad signaling via TGF-beta-dependent and independent mechanisms: implications for diabetic renal and vascular disease. *FASEB J.* **2004**, *18*, 176–178.