

**Table S3.** Genome size of *Crepis* species analysed in present study and data retrieved from literature

Species	Genome size (pg/1C ± SD present study)	Genome size (pg/1C ± SD) data taken from literature
<i>Crepis aculeata</i>	2.89 ± 0.04	-
<i>C. acuminata</i> (2x)		13.8 ± 0.4 [1]
<i>C. albida</i>	3.08 ± 0.03	-
<i>C. alpestris</i>	2.99 ± 0.03	-
<i>C. alpina</i>	2.20 ± 0.05	3.00 [2]
<i>C. aspera</i>	2.15 ± 0.02	-
<i>C. atribarba</i> (2x)		13.3 ± 0.7 [1]
<i>C. aurea</i>	1.63 ± 0.10	1.75 ± 0.07 [3]
<i>C. biennis</i>	10.45 ± 0.33	9.51 ± 0.492 [3]
<i>C. bungei</i>		4.05 ± 0.053 [3]
<i>C. capillaris</i>	2.07 ± 0.08	-
<i>C. chrysanthia</i> subsp. <i>chrysanthia</i>		4.64 ± 0.053 [3]
<i>C. conyzifolia</i>	6.08 ± 0.16	-
<i>C. crocea</i> (4x)		9.61 ± 0.053 [3]
<i>C. dioscoridis</i>	4.58 ± 0.13	-
<i>C. foetida</i>	2.03 ± 0.08	1.46 ± 0.034 [3]
<i>C. foetida</i> subsp. <i>rhoaedifolia</i>	2.17 ± 0.02	-
<i>C. jacquinnii</i>	5.12 ± 0.05	-
<i>C. kotschyana</i>	2.92 ± 0.08	-
<i>C. lacera</i>	7.46 ± 0.10	-
<i>C. lapsanoides</i>		5.6 [4]
<i>C. leontodontoides</i>	1.06 ± 0.03	1.14 ± 0.012 [3]
<i>C. mollis</i>	2.53 ± 0.04	-
<i>C. modocensis</i> (2x)		12.2 ± 0.3 [1]
<i>C. multicaulis</i> subsp. <i>congesta</i>		1.78 ± 0.018 [3]
<i>C. nicaeensis</i>	3.17 ± 0.02	-
<i>C. occidentalis</i> (4x)		23.8 ± 1.7 [1]
<i>C. palaestina</i>	7.05 ± 0.19	6.10 [4]
<i>C. paludosa</i>	4.53 ± 0.19	4.53 ± 0.074 [3]
<i>C. pannonica</i>	7.27 ± 0.06	-
<i>C. polymorpha</i>	3.13 ± 0.02	-
<i>C. pontana</i>		6.9 [4]
<i>C. pulchra</i>	5.59 ± 0.12	5.50 [5]
<i>C. pusilla</i>		1.11 ± 0.025 [3]
<i>C. praemorsa</i>		5.3 [4]
<i>C. pyrenaica</i>	3.54 ± 0.05	3.58 ± 0.051 [3]
<i>C. rubra</i>	2.86 ± 0.07	2.90 [4]

<i>C. sancta</i>	1.60 ± 0.02	-
<i>C. setosa</i>	1.67 ± 0.02	1.70 [5]
<i>C. sibirica</i>	6.98 ± 0.04	-
<i>C. succisifolia</i>	2.34 ± 0.05	-
<i>C. syriaca</i>	2.39 ± 0.28	-
<i>C. taraxacifolia</i>	2.47 ± 0.58	-
<i>C. tectorum</i>	3.06 ± 0.05	-
<i>C. vesicaria</i> (1) 4x	2.78 ± 0.09	-
<i>C. vesicaria</i> (3) 2x	2.43 ± 0.03	-
<i>C. viscidula</i> subsp. <i>gieracioides</i>		4.90 [5]
<i>C. zacintha</i>	1.03 ± 0.02	-
<i>Lapsana communis</i>	1.22 ± 0.04	-

## References

- [1] Sears, C.J., Whitton, J., 2016. A reexamination of the North American *Crepis* agamic complex and comparison with the findings of Babcock and Stebbins' classic biosystematic monograph. *American Journal of Botany* 103, 1289-1299.
- [2] Wallace, H., Sparkes, C.A., Maden, M., 1972. Nuclear DNA content of three *Crepis* species. *Heredity* 29, 367-373
- [3] Enke, N, Fuchs, J, Gemeinholzer, B. 2011. Shrinking genomes? Evidence from genome size variation in *Crepis* (Compositae). *Plant Biology* 13: 185-193.
- [4] Bennett, MD, Smith, JB. 1976. Nuclear-DNA Amounts in Angiosperms. *Philosophical Transactions of the Royal Society B-Biological Sciences* 274: 227-274.
- [5] Dimitrova, D, Greilhuber, J. 2000. Karyotype and DNA-content evolution in ten species of *Crepis* (Asteraceae) distributed in Bulgaria. *Botanical Journal of the Linnean Society* 132: 281-297.