

xerophilic species of open land (x); df: 5, 74				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.344	3.060	18.140	<0.001
Coefficients _{HAC}	B	SE	t	p
Shade [%]	-0.101	0.038	-2.675	0.092
Herb [%]	0.090	0.111	0.811	0.420
Grass [%]	-0.067	0.051	-1.318	0.192
Litter [%]	0.010	0.044	0.222	0.825
Open ground [%]	0.082	0.071	1.158	0.251
hygrophilic species of open land (h); df: 5, 74				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.401	2.935	33.792	<0.001
Coefficients _{HAC}	B	SE	t	p
Shade [%]	-0.063	0.029	-2.195	0.013
Herb [%]	-0.184	0.109	-1.691	0.095
Grass [%]	0.012	0.038	0.313	0.756
Litter [%]	-0.119	0.033	-3.580	0.001
Open ground [%]	-0.132	0.025	-5.231	<0.001
eurytopic species of open land (eu); df: 5, 74				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.323	3.000	12.236	<0.001
Coefficients _{HAC}	B	SE	t	p
Shade [%]	-0.079	0.017	-4.529	<0.001
Herb [%]	0.091	0.064	1.416	0.161
Grass [%]	0.071	0.051	1.397	0.167
Litter [%]	0.060	0.041	1.462	0.148
Open ground [%]	0.124	0.030	4.101	<0.001
slightly hygrophilic forest species ((h)f) ¹⁾ ; df: 5, 72				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.413	3.596	18.662	<0.001
Coefficients _{HAC}	B	SE	t	p
Shade [%]	0.146	0.017	8.755	<0.001
Herb [%]	0.002	0.082	0.025	0.981
Grass [%]	-0.101	0.042	-4.424	0.018
Litter [%]	-0.055	0.025	-2.179	0.033
Open ground [%]	-0.142	0.032	-4.385	<0.001
eurytopic forest species (f); df: 5, 74				
Model overview	R ²	SE	F	p
Statistics _{HAC}	0.487	4.042	73.986	<0.001
Coefficients _{HAC}	B	SE	t	p
Shade [%]	0.094	0.030	3.105	0.003
Herb [%]	0.069	0.100	0.692	0.491
Grass [%]	0.167	0.063	2.626	0.011
Litter [%]	0.140	0.050	2.789	0.007
Open ground [%]	0.093	0.050	1.878	0.064

After elimination of the outliers in SRC3 trap1 2014 and SRC3 Trap2 2014. Outlier test +/- 3 SD