

| Species | Author | Ind | ET | HP |
|------------------------------------|-------------------------|------|--------|-----|
| <i>Aculepeira ceropegia</i> | (Walckenaer, 1802) | 1 | (x) | FIE |
| <i>Agroeca proxima</i> | (O. P.-Cambridge, 1871) | 2 | x | OTH |
| <i>Agyneta affinis</i> | (Kulczynski, 1898) | 1 | (h) | OHA |
| <i>Agyneta decora</i> | (O. P.-Cambridge, 1871) | 20 | h | FAL |
| <i>Agyneta rurestris</i> | (C.L. Koch, 1836) | 172 | (x) | FIE |
| <i>Allomengea vidua</i> | (L. Koch, 1879) | 1 | h | WME |
| <i>Alopecosa cuneata</i> | (Clerck, 1757) | 912 | (x) | PAS |
| <i>Alopecosa pulverulenta</i> | (Clerck, 1757) | 496 | (x) | OHA |
| <i>Alopecosa taeniata</i> | (C.L. Koch, 1835) | 1 | (h)f | LMF |
| <i>Amaurobius fenestralis</i> | (Ström, 1768) | 4 | (h)f | MFO |
| <i>Anelasmacephalus cambridgei</i> | (Westwood, 1874) | 31 | (h)f | MFO |
| <i>Anguliphantes angulipalpis</i> | (Westring, 1851) | 1 | (x)f | AFO |
| <i>Anyphaena accentuata</i> | (Walckenaer, 1802) | 7 | (h)f | MFO |
| <i>Araeoncus humilis</i> | (Blackwall, 1841) | 14 | (x) | FIE |
| <i>Araneus diadematus</i> | Clerck, 1757 | 1 | (x)(f) | AFO |
| <i>Araneus quadratus</i> | Clerck, 1757 | 4 | (h) | GRL |
| <i>Araneus triguttatus</i> | (Fabricius, 1793) | 1 | (h)f | MFO |
| <i>Aulonia albimana</i> | (Walckenaer, 1805) | 9 | x | ODH |
| <i>Bathyphantes gracilis</i> | (Blackwall, 1841) | 247 | (x) | FIE |
| <i>Bathyphantes parvulus</i> | (Westring, 1851) | 52 | eu | FAL |
| <i>Centromerita bicolor</i> | (Blackwall, 1833) | 1070 | (x)(f) | PFO |
| <i>Centromerus brevipalpus</i> | (Menge, 1866) | 5 | h(f) | MFO |
| <i>Centromerus dilutus</i> | (O. P.-Cambridge, 1875) | 3 | (h)f | MFO |
| <i>Centromerus incilium</i> | (L. Koch, 1881) | 1 | (h)(f) | PFO |
| <i>Centromerus sylvaticus</i> | (Blackwall, 1841) | 418 | (h)(f) | PFO |
| <i>Ceratinella brevis</i> | (Wider, 1834) | 15 | (h)f | MFO |
| <i>Cheiracanthium erraticum</i> | (Walckenaer, 1802) | 2 | eu | OHA |
| <i>Cicurina cicur</i> | (Fabricius, 1793) | 3 | (h)(f) | PFO |
| <i>Clubiona compta</i> | C.L. Koch, 1839 | 8 | (x)f | AFO |
| <i>Clubiona diversa</i> | O. P.-Cambridge, 1862 | 8 | (h) | OHA |
| <i>Clubiona pallidula</i> | (Clerck, 1757) | 2 | (h)(f) | FOR |
| <i>Clubiona reclusa</i> | O. P.-Cambridge, 1863 | 115 | (x) | FAL |
| <i>Clubiona terrestris</i> | Westring, 1851 | 21 | (x)f | AFO |
| <i>Cnephalocotes obscurus</i> | (Blackwall, 1834) | 35 | (h) | FAL |
| <i>Coelotes terrestris</i> | (Wider, 1834) | 311 | (h)f | MFO |
| <i>Collinsia inerrans</i> | (O. P.-Cambridge, 1885) | 129 | (h) | FIE |
| <i>Cryptachaea riparia</i> | (Blackwall, 1834) | 1 | (x) | FAL |
| <i>Cyclosa conica</i> | (Pallas, 1772) | 3 | (x)f | AFO |
| <i>Diaea dorsata</i> | (Fabricius, 1777) | 1 | (h)f | MFO |
| <i>Dicymbium brevisetosum</i> | Locket, 1962 | 515 | eu | FAL |
| <i>Diplocephalus latifrons</i> | (O. P.-Cambridge, 1863) | 66 | (h)f | MFO |
| <i>Diplocephalus picinus</i> | (Blackwall, 1841) | 677 | (h)f | MFO |
| <i>Diplostyla concolor</i> | (Wider, 1834) | 247 | (h)(f) | PFO |
| <i>Drassodes lapidosus</i> | (Walckenaer, 1802) | 2 | eu | ODH |
| <i>Drassodes pubescens</i> | (Thorell, 1856) | 5 | x | FAL |
| <i>Drassyllus lutetianus</i> | (L. Koch, 1866) | 130 | (h)(f) | FAL |
| <i>Drassyllus pusillus</i> | (C.L. Koch, 1833) | 166 | (x)(f) | OHA |
| <i>Enoplognatha ovata</i> | (Clerck, 1757) | 14 | (x)(f) | PFO |
| <i>Enoplognatha thoracica</i> | (Hahn, 1833) | 45 | (x) | PAS |
| <i>Episinus angulatus</i> | (Blackwall, 1836) | 1 | (h)(f) | FOR |
| <i>Eratigena picta</i> | (Simon, 1870) | 6 | (x)(f) | MFO |
| <i>Erigone atra</i> | Blackwall, 1833 | 2108 | (h) | FIE |
| <i>Erigone dentipalpis</i> | (Wider, 1834) | 1412 | (x) | FIE |

| | | | | |
|---------------------------------|-------------------------|------|--------|-----|
| <i>Erigonella hiemalis</i> | (Blackwall, 1841) | 128 | (h) | FAL |
| <i>Ero furcata</i> | (Villers, 1789) | 3 | (x)(f) | AFO |
| <i>Evarcha arcuata</i> | (Clerck, 1757) | 2 | eu | OHA |
| <i>Evarcha falcata</i> | (Clerck, 1757) | 1 | (x)(f) | AFO |
| <i>Floronia bucculenta</i> | (Clerck, 1757) | 4 | (h)f | MFO |
| <i>Gongyliellum latebricola</i> | (O. P.-Cambridge, 1871) | 3 | (h)f | FOR |
| <i>Gongyliellum vivum</i> | (O. P.-Cambridge, 1875) | 1 | h | BOG |
| <i>Hahnia helveola</i> | Simon, 1875 | 4 | (h)f | MFO |
| <i>Hahnia nava</i> | (Blackwall, 1841) | 3 | (x)(f) | NHP |
| <i>Haplodrassus signifer</i> | (C.L. Koch, 1839) | 38 | x | OTH |
| <i>Haplodrassus silvestris</i> | (Blackwall, 1833) | 14 | (x)f | AFO |
| <i>Histoipona torpida</i> | (C.L. Koch, 1837) | 33 | (h)f | LMF |
| <i>Inermocoelotes inermis</i> | (L. Koch, 1855) | 80 | (h)f | LMF |
| <i>Labulla thoracica</i> | (Wider, 1834) | 1 | (h)f | LMF |
| <i>Lacinius dentiger</i> | (C.L. Koch, 1847) | 134 | (h)f | MFO |
| <i>Lacinius ephippiatus</i> | (C.L. Koch, 1835) | 1833 | (h)f | MFO |
| <i>Larinioides cornutus</i> | (Clerck, 1757) | 2 | h | REE |
| <i>Leiobunum blackwalli</i> | Meade, 1861 | 3 | (h)(f) | MFO |
| <i>Leiobunum rotundum</i> | (Latreille, 1798) | 1 | f | AFO |
| <i>Lepthyphantes minutus</i> | (Blackwall, 1833) | 4 | (x)f | AFO |
| <i>Linyphia hortensis</i> | Sundevall, 1830 | 8 | (h)f | MFO |
| <i>Linyphia triangularis</i> | (Clerck, 1757) | 2 | (x)(f) | NHP |
| <i>Lophopilio palpalis</i> | (Herbst, 1799) | 133 | (h)f | MFO |
| <i>Mangora acalypha</i> | (Walckenaer, 1802) | 4 | eu | NHP |
| <i>Maso sundevalli</i> | (Westring, 1851) | 6 | (h)f | MFO |
| <i>Mermessus trilobatus</i> | (Emerton, 1882) | 56 | eu | NHP |
| <i>Metellina segmentata</i> | (Clerck, 1757) | 4 | (f) | MFO |
| <i>Micaria pulicaria</i> | (Sundevall, 1831) | 12 | eu | OHA |
| <i>Micrargus herbigradus</i> | (Blackwall, 1854) | 10 | (x)f | FOR |
| <i>Microlinyphia pusilla</i> | (Sundevall, 1830) | 4 | (h) | OHA |
| <i>Microneta viaria</i> | (Blackwall, 1841) | 210 | (h)f | MFO |
| <i>Mioxena blanda</i> | (Simon, 1884) | 4 | x | OHA |
| <i>Mitostoma chrysomelas</i> | (Hermann, 1804) | 4 | (h)(f) | FAL |
| <i>Monocephalus fuscipes</i> | (Blackwall, 1836) | 6 | (h)f | LMF |
| <i>Nemastoma lugubre</i> | (O.F. Müller, 1776) | 362 | (h)f | MFO |
| <i>Meom reticulatus</i> | (Blackwall, 1853) | 1 | (h)f | MFO |
| <i>Neottiura bimaculata</i> | (Linnaeus, 1767) | 7 | (h) | OHA |
| <i>Neriere clathrata</i> | (Sundevall, 1830) | 10 | (h)(f) | PFO |
| <i>Oedothorax apicatus</i> | (Blackwall, 1850) | 5532 | (x) | FIE |
| <i>Oedothorax fuscus</i> | (Blackwall, 1834) | 22 | (h) | WME |
| <i>Oedothorax gibbosus</i> | (Blackwall, 1841) | 5 | h | BOG |
| <i>Oedothorax retusus</i> | (Westring, 1851) | 23 | h | PAS |
| <i>Oligolophus tridens</i> | (C.L. Koch, 1836) | 3195 | (h)(f) | MFO |
| <i>Opilio canestrinii</i> | (Thorell, 1876) | 6 | (h)f | MFO |
| <i>Ostearius melanopygius</i> | (O. P.-Cambridge, 1879) | 2 | (x) | PIO |
| <i>Ozyptila praticola</i> | (C.L. Koch, 1837) | 28 | (x)f | MFO |
| <i>Pachygnatha clercki</i> | Sundevall, 1823 | 478 | h | OHA |
| <i>Pachygnatha degeeri</i> | Sundevall, 1830 | 1701 | eu | OHA |
| <i>Pachygnatha listeri</i> | Sundevall, 1830 | 2 | (h)f | MFO |
| <i>Palliduphantes pallidus</i> | (O. P.-Cambridge, 1871) | 21 | (h)f | MFO |
| <i>Pardosa agrestis</i> | (Westring, 1861) | 49 | (x) | FIE |
| <i>Pardosa amentata</i> | (Clerck, 1757) | 4380 | h (f) | PFO |
| <i>Pardosa lugubris</i> | (Walckenaer, 1802) | 266 | (h)(f) | PFO |
| <i>Pardosa nigriceps</i> | (Thorell, 1856) | 2 | x | DGL |

| | | | | |
|---------------------------------|---------------------------|------|--------|-----|
| <i>Pardosa palustris</i> | (Linnaeus, 1758) | 6203 | eu | GRL |
| <i>Pardosa pullata</i> | (Clerck, 1757) | 2359 | (h) | OHA |
| <i>Pardosa saltans</i> | Töpfer-Hofmann, 2000 | 127 | (h)(f) | PFO |
| <i>Paroligolophus agrestis</i> | (Meade, 1855) | 5 | (h)(f) | PFO |
| <i>Pelecopsis parallela</i> | (Wider, 1834) | 1 | (h) | OHA |
| <i>Phalangium opilio</i> | Linnaeus, 1758 | 87 | (x) | FIE |
| <i>Philodromus aureolus</i> | (Clerck, 1757) | 1 | (x)f | AFO |
| <i>Philodromus cespitum</i> | (Walckenaer, 1802) | 1 | (x)f | AFO |
| <i>Pholcomma gibbum</i> | (Westring, 1851) | 1 | (x)(f) | AFO |
| <i>Phrurolithus festivus</i> | (C.L. Koch, 1835) | 1 | (x) | DRY |
| <i>Phrurolithus minimus</i> | C.L. Koch, 1839 | 3 | x | DGL |
| <i>Phylloneta impressa</i> | (L. Koch, 1881) | 1 | (x) | FAL |
| <i>Piratula hygrophila</i> | (Thorell, 1872) | 1 | h(f) | WFO |
| <i>Piratula latitans</i> | (Blackwall, 1841) | 13 | h | OWH |
| <i>Piratula uliginosa</i> | (Thorell, 1856) | 274 | h | FAL |
| <i>Pisaura mirabilis</i> | (Clerck, 1757) | 12 | (h) | OHA |
| <i>Platybunus bucephalus</i> | (C.L. Koch, 1835) | 35 | (h)f | LMF |
| <i>Pocadicnemis juncea</i> | Locket & Millidge, 1953 | 3 | h | OHA |
| <i>Pocadicnemis pumila</i> | (Blackwall, 1841) | 7 | (h)(f) | FOR |
| <i>Porrhomma errans</i> | (Blackwall, 1841) | 70 | (h)(f) | PFO |
| <i>Porrhomma microphthalmum</i> | (O. P.-Cambridge, 1871) | 35 | (x) | FIE |
| <i>Porrhomma oblitum</i> | (O. P.-Cambridge, 1871) | 8 | h | VRB |
| <i>Porrhomma pygmaeum</i> | (Blackwall, 1834) | 1 | h(f) | WFO |
| <i>Pseudophrys erratica</i> | (Walckenaer, 1826) | 1 | (x)f | AFO |
| <i>Rilaena triangularis</i> | (Herbst, 1799) | 800 | (h)(f) | GRL |
| <i>Robertus arundineti</i> | (O. P.-Cambridge, 1871) | 6 | eu | OHA |
| <i>Robertus lividus</i> | (Blackwall, 1836) | 20 | (x)f | MFO |
| <i>Robertus neglectus</i> | (O. P.-Cambridge, 1871) | 51 | (h)f | MFO |
| <i>Segestria senoculata</i> | (Linnaeus, 1758) | 3 | (x)f | AFO |
| <i>Stemonyphantes lineatus</i> | (Linnaeus, 1758) | 6 | (x)f | PFO |
| <i>Styloctetor compar</i> | (Westring, 1861) | 1 | h | WME |
| <i>Talavera aequipes</i> | (O. P.-Cambridge, 1871) | 1 | x | DGL |
| <i>Tallusia experta</i> | (O. P.-Cambridge, 1871) | 31 | h(f) | FAL |
| <i>Tapinocyba insecta</i> | (L. Koch, 1869) | 50 | (h)f | MFO |
| <i>Tapinocyba praecox</i> | (O. P.-Cambridge, 1873) | 2 | x | FAL |
| <i>Tapinopa longidens</i> | (Wider, 1834) | 1 | (x)f | AFO |
| <i>Tenuiphantes cristatus</i> | (Menge, 1866) | 1 | (h)f | MFO |
| <i>Tenuiphantes flavipes</i> | (Blackwall, 1854) | 371 | (h)f | MFO |
| <i>Tenuiphantes mengei</i> | (Kulczynski, 1887) | 9 | h(f) | FOR |
| <i>Tenuiphantes tenebricola</i> | (Wider, 1834) | 61 | (h)f | MFO |
| <i>Tenuiphantes tenuis</i> | (Blackwall, 1852) | 256 | (x) | GRL |
| <i>Tenuiphantes zimmermanni</i> | (Bertkau, 1890) | 108 | (h)f | MFO |
| <i>Tetragnatha montana</i> | Simon, 1874 | 1 | h(f) | WFO |
| <i>Tetragnatha nigrita</i> | Lendl, 1886 | 1 | h(f) | VRB |
| <i>Tetragnatha obtusa</i> | C.L. Koch, 1837 | 1 | (x)f | AFO |
| <i>Tetragnatha pinicola</i> | L. Koch, 1870 | 3 | (x) | FIE |
| <i>Theridion varians</i> | Hahn, 1833 | 1 | (x) | ODH |
| <i>Thyreosthenius biovatus</i> | (O. P.-Cambridge, 1875) | 1 | x | DGL |
| <i>Tiso vagans</i> | (Blackwall, 1834) | 94 | (h) | PAS |
| <i>Trochosa ruricola</i> | (De Geer, 1778) | 1324 | eu | GRL |
| <i>Trochosa spinipalpis</i> | (F.O. P.-Cambridge, 1895) | 1 | h | WME |
| <i>Trochosa terricola</i> | Thorell, 1856 | 782 | (h) | FAL |
| <i>Troxochrus scabriculus</i> | (Westring, 1851) | 3 | x | DGL |
| <i>Walckenaeria acuminata</i> | Blackwall, 1833 | 1 | (h)f | MFO |

| | | | | |
|----------------------------------|-------------------------|-----|--------|-----|
| <i>Walckenaeria alticeps</i> | (Denis, 1952) | 4 | hf | WFO |
| <i>Walckenaeria antica</i> | (Wider, 1834) | 1 | (x)(f) | PFO |
| <i>Walckenaeria atrotibialis</i> | (O. P.-Cambridge, 1878) | 52 | (h)f | MFO |
| <i>Walckenaeria corniculans</i> | (O. P.-Cambridge, 1875) | 5 | (h)f | MFO |
| <i>Walckenaeria cucullata</i> | (C.L. Koch, 1836) | 3 | (h)f | MFO |
| <i>Walckenaeria cuspidata</i> | Blackwall, 1833 | 2 | hf | WFO |
| <i>Walckenaeria dysderoides</i> | (Wider, 1834) | 31 | (x)f | MFO |
| <i>Walckenaeria nudipalpis</i> | (Westring, 1851) | 56 | h(f) | OWH |
| <i>Walckenaeria obtusa</i> | Blackwall, 1836 | 15 | (h)f | MFO |
| <i>Walckenaeria unicornis</i> | O. P.-Cambridge, 1861 | 4 | h(f) | OWH |
| <i>Walckenaeria vigilax</i> | (Blackwall, 1853) | 98 | h | FIE |
| <i>Xysticus bifasciatus</i> | C.L. Koch, 1837 | 9 | x | FAL |
| <i>Xysticus cristatus</i> | (Clerck, 1757) | 138 | (x) | GRL |
| <i>Xysticus kochi</i> | Thorell, 1872 | 138 | (x) | OHA |
| <i>Xysticus lanio</i> | C.L. Koch, 1835 | 7 | (h)f | MFO |
| <i>Xysticus ulmi</i> | (Hahn, 1831) | 44 | (h) | OHA |
| <i>Zelotes clivicola</i> | (L. Koch, 1870) | 3 | (x)f | AFO |
| <i>Zelotes latreillei</i> | (Simon, 1878) | 14 | (x) | ODH |
| <i>Zelotes subterraneus</i> | (C.L. Koch, 1833) | 6 | (x)(f) | MFO |
| <i>Zora spinimana</i> | (Sundevall, 1833) | 12 | eu | NHP |
